


FORM PTO-1390 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER PF-0629 USN
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		U.S. APPLICATION NUMBER (U.S. Patent, PCT or CBP) TO BE ASSIGNED 097856679
INTERNATIONAL APPLICATION NO. PCT/US99/28013	INTERNATIONAL FILING DATE 23 November 1999	PRIORITY DATE CLAIMED 23 November 1998
TITLE OF INVENTION GTPASE ASSOCIATED PROTEINS		
APPLICANT(S) FOR DO/EO/US INCYTE PHARMACEUTICALS, INC.; HILLMAN, Jennifer L.; TANG, Y. Tom; BANDMAN, Olga; LAL, Preeti; YUE, Henry; LU, Dyung Aina M.; BAUGHN, Mariah R.; YANG, Junming; AZIMZAI, Yalda		
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information: 1. <input checked="" type="checkbox"/> This is the FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input type="checkbox"/> This is an express request to promptly begin national examination procedures (35 U.S.C. 371 (f)). 4. <input type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (PCT Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau) b. <input type="checkbox"/> has been communicated by the International Bureau. c. <input checked="" type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).		
Items 11 to 16 below concern document(s) or information included:		
11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.27 and 3.31 is included. 13. <input type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 14. <input type="checkbox"/> A substitute specification. 15. <input type="checkbox"/> A change of power of attorney and/or address letter. 16. <input checked="" type="checkbox"/> Other items or information: 1) Transmittal Letter (2 pp, in duplicate) 2) Return Postcard 3) Express Mail Label No.: <u>EL 856 153 928 US</u> 4) Request to Transfer		

097856679.153204

U.S. APPLICATION NO. <u>09/0856679</u> (37 CFR 1.5) TO BE DESIGNATED		INTERNATIONAL APPLICATION NO.: PCT/US99/28013		ATTORNEY'S DOCKET NUMBER PF-0629 USN	
17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO.....\$1000.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO.....\$860.00 <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO.....\$710.00 <input checked="" type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4).....\$690.00 <input type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4).....\$100.00					
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$690.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total Claims	20 =	0	X \$ 18.00	\$	
Independent Claims	2 =	0	X \$ 80.00	\$	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)				+ \$270.00	\$
TOTAL OF ABOVE CALCULATIONS =				\$690.00	
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.				\$	
SUBTOTAL =				\$690.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
TOTAL NATIONAL FEE =				\$690.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by the appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				\$	
TOTAL FEES ENCLOSED =				\$690.00	
				Amount to be Refunded:	\$
				Charged:	\$
a. <input type="checkbox"/> A check in the amount of \$_____ to cover the above fees is enclosed. b. <input checked="" type="checkbox"/> Please charge my Deposit Account No. <u>09-0108</u> in the amount of \$ <u>690.00</u> to cover the above fees. c. <input type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>09-0108</u> . A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO:					
INCYTE GENOMICS, INC. 3160 Porter Drive Palo Alto, CA 94304			 SIGNATURE		
NAME: Diana Hamlet-Cox					
REGISTRATION NUMBER: 33,302					
DATE: <u>22</u> May 2001					

090856679 "PCT/PTO"

GTPASE ASSOCIATED PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of GTPase associated proteins and to the use of these sequences in the diagnosis, treatment, and prevention of cell proliferative, autoimmune/inflammatory, and immune system disorders.

BACKGROUND OF THE INVENTION

Guanine nucleotide binding proteins (GTP-binding proteins) participate in a wide range of regulatory functions in all eukaryotic cells, including metabolism, cellular growth, differentiation, signal transduction, cytoskeletal organization, and intracellular vesicle transport and secretion. In higher organisms they are involved in signaling that regulates such processes as the immune response (Aussel, C. et al (1988) J. Immunol. 140:215-220), apoptosis, differentiation, and cell proliferation including oncogenesis (Dhanasekaran, N. et al. (1998) Oncogene 17:1383-1394). Exchange of bound GDP for GTP followed by hydrolysis of GTP to GDP provides the energy that enables GTP-binding proteins to alter their conformation and interact with other cellular components. The superfamily of GTP-binding proteins consists of several families and may be grouped as translational factors, heterotrimeric GTP-binding proteins involved in transmembrane signaling processes (also called G-proteins), and low molecular weight GTP-binding proteins including the proto-oncogene Ras proteins and products of rab, rap, rho, rac, smg21, smg25, YPT, SEC4, and ARF genes, and tubulins (Kaziro, Y. et al. (1991) Ann. Rev. Biochem. 60:349-400). In all cases, the GTPase activity is regulated through interactions with other proteins.

GTP-binding proteins involved in protein biosynthesis include initiation factor 2 (IF-2), elongation factor 2 (EF-Tu), and elongation factor G (EF-G), observed in prokaryotes; and initiation factor 2 (eIF-2), elongation factor 1 α (EF-1 α) and elongation factor 2 (EF-2) observed in eukaryotes (Kaziro, supra). IF-2 promotes the GTP-dependent binding of the tRNA to the small subunit of the ribosome, the step that initiates protein translation. Similarly, elongation factors promote the binding of tRNA and GTP and the displacement of GDP after hydrolysis as protein biosynthesis proceeds.

Heterotrimeric GTP-binding proteins are composed of 3 subunits (α , β and γ) which, in their inactive conformation, associate as a trimer at the inner face of the plasma membrane. G_α binds GDP or GTP and contains the GTPase activity. The $\beta\gamma$ complex enhances binding of G_α to a receptor. G_γ is necessary for the folding and activity of $G\beta$. (Neer, E.J. et al. (1994) Nature 371:297-300.) Multiple homologs of each subunit have been identified in mammalian tissues, and different combinations of subunits have specific functions and tissue specificities. (Spiegel, A.M. (1997) J.

Inher. Metab. Dis. 20:113-121.) G protein activity is triggered by seven-transmembrane cell surface receptors (G-protein coupled receptors) which respond to lipid analogs, amino acids and their derivatives, peptides, cytokines, and specialized stimuli such as light, taste, and odor. Activation of the receptor by its stimulus causes the replacement of the G protein-bound GDP with GTP. G α -GTP dissociates from the receptor/ $\beta\gamma$ complex and each of these separated components can interact with and regulate downstream effectors. The signaling stops when G α hydrolyzes its bound GTP to GDP and reassociates with the $\beta\gamma$ complex (Neer, supra).

The alpha subunits of heterotrimeric G proteins can be divided into four distinct classes. The α -s class is sensitive to ADP-ribosylation by pertussis toxin which uncouples the receptor:G-protein interaction. This uncoupling blocks signal transduction to receptors that decrease cAMP levels which normally regulate ion channels and activate phospholipases. The inhibitory α -i class is also susceptible to modification by pertussis toxin which prevents α -i from lowering cAMP levels. Two novel classes of α subunits refractory to pertussis toxin modification are α -q, which activates phospholipase C, and α -12, which has sequence homology with the *Drosophila* gene concertina and may contribute to the regulation of embryonic development (Simon, M.I. (1991) *Science* 252:802-808).

The mammalian G β and G γ subunits, each about 340 amino acids long, share more than 80% homology. The G β subunit (also called transducin) contains seven repeating units, each about 43 amino acids long. The activity of both subunits may be regulated by other proteins such as calmodulin and phosducin or the neural protein GAP 43 (D. Clapham and E. Neer, 1993, *Nature* 365:403-406). The β and γ subunits are tightly associated. The β subunit sequences are highly conserved between species, implying that they perform a fundamentally important role in the organization and function of G-protein linked systems (Van der Voorn L. (1992) *Febs. Lett.* 307 (2):131-134). They contain seven tandem repeats of the WD-repeat sequence motif, a motif found in many proteins with regulatory functions. WD-repeat proteins contain from four to eight copies of a loosely conserved repeat of approximately 40 amino acids which participates in protein-protein interactions. Mutations and variant expression of β transducin proteins are linked with various disorders. Mutations in LIS1, a subunit of the human platelet activating factor acetylhydrolase, cause Miller-Dieker lissencephaly. RACK1 binds activated protein kinase C, and RbAp48 binds retinoblastoma protein. CstF is required for polyadenylation of mammalian pre-mRNA *in vitro* and associates with subunits of cleavage-stimulating factor. Defects in the regulation of β -catenin contribute to the neoplastic transformation of human cells. The WD40 repeats of the human F-box protein β TrCP mediate binding to β -catenin, thus regulating the targeted degradation of β -catenin by

ubiquitin ligase (Neer, *supra*; Hart, M. et al (1999) *Curr. Biol.* 9:207-210). The γ subunit primary structures are more variable than those of the β subunits. They are often post-translationally modified by isoprenylation and carboxyl-methylation of a cysteine residue four amino acids from the C-terminus; this appears to be necessary for the interaction of the $\beta\gamma$ subunit with the membrane and

5 with other GTP-binding proteins. The $\beta\gamma$ subunit has been shown to modulate the activity of isoforms of adenylyl cyclase, phospholipase C, and some ion channels. It is involved in receptor phosphorylation via specific kinases, and has been implicated in the p21ras-dependent activation of the MAP kinase cascade and the recognition of specific receptors by GTP-binding proteins. (Clapham and Neer, *supra*).

10 G-proteins interact with a variety of effectors including adenylyl cyclase (Clapham and Neer, *supra*). The signaling pathway mediated by cAMP is mitogenic in hormone-dependent endocrine tissues such as adrenal cortex, thyroid, ovary, pituitary, and testes. Cancers in these tissues have been related to a mutationally activated form of a $G\alpha$, known as the gsp (Gs protein) oncogene (Dhanasekaran, *supra*). Another effector is phosphducin, a retinal phosphoprotein, which forms a

15 specific complex with retinal $G\beta$ and $G\gamma$ ($G\beta\gamma$) and modulates the ability of $G\beta\gamma$ to interact with retinal $G\alpha$ (Clapham and Neer, *supra*).

Irregularities in the GTP-binding protein signaling cascade may result in abnormal activation of leukocytes and lymphocytes, leading to the tissue damage and destruction seen in many inflammatory and autoimmune diseases such as rheumatoid arthritis, biliary cirrhosis, hemolytic

20 anemia, lupus erythematosus, and thyroiditis. Abnormal cell proliferation, including cyclic AMP stimulation of brain, thyroid, adrenal, and gonadal tissue proliferation is regulated by G proteins. Mutations in $G\alpha$ subunits have been found in growth-hormone-secreting pituitary somatotroph tumors, hyperfunctioning thyroid adenomas, and ovarian and adrenal neoplasms (Meij, J.T.A. (1996) *Mol. Cell. Biochem.* 157:31-38; Ausset, *supra*).

25 LMW GTP-binding proteins are GTPases which regulate cell growth, cell cycle control, protein secretion, and intracellular vesicle interaction. They consist of single polypeptides which, like the α subunit of the heterotrimeric GTP-binding proteins, are able to bind to and hydrolyze GTP, thus cycling between an inactive and an active state. LMW GTP-binding proteins respond to extracellular signals from receptors and activating proteins by transducing mitogenic signals involved

30 in various cell functions. The binding and hydrolysis of GTP regulates the response of LMW GTP-binding proteins and acts as an energy source during this process (Bokoch, G. M. and Der, C. J. (1993) *FASEB J.* 7:750-759).

At least sixty members of the LMW GTP-binding protein superfamily have been identified _

and are currently grouped into the ras, rho, arf, sar1, ran, and rab subfamilies. Activated ras genes were initially found in human cancers, and subsequent studies confirmed that ras function is critical in determining whether cells continue to grow or become differentiated. Ras1 and Ras2 proteins stimulate adenylate cyclase (Kaziro, supra), affecting a broad array of cellular processes. Stimulation of cell surface receptors activates Ras which, in turn, activates cytoplasmic kinases. These kinases translocate to the nucleus and activate key transcription factors that control gene expression and protein synthesis (Barbacid, M. (1987) *Ann. Rev Biochem.* 56:779-827, Treisman, R. (1994) *Curr. Opin.Genet. Dev.* 4:96-98). Other members of the LMW GTP-binding protein superfamily have roles in signal transduction that vary with the function of the activated genes and the locations of the GTP-binding proteins that initiate the activity. Rho GTP-binding proteins control signal transduction pathways that link growth factor receptors to actin polymerization, which is necessary for normal cellular growth and division. The rab, arf, and sar1 families of proteins control the translocation of vesicles to and from membranes for protein processing, localization, and secretion. Vesicle- and target- specific identifiers (v-SNAREs and t-SNAREs) bind to each other and dock the vesicle to the acceptor membrane. The budding process is regulated by the closely related ADP ribosylation factors (ARFs) and SAR proteins, while rab proteins allow assembly of SNARE complexes and may play a role in removal of defective complexes (J. Rothman and F. Wieland (1996) *Science* 272:227-234). Ran GTP-binding proteins are located in the nucleus of cells and have a key role in nuclear protein import, the control of DNA synthesis, and cell-cycle progression (Hall, A. (1990) *Science* 249:635-640; Barbacid, M. (1987) *Ann. Rev Biochem.* 56:779-827; Kistakis, N. (1998) *BioEssays* 20:495-504; and Sasaki, T. and Takai, Y. (1998) *Biochem. Biophys. Res. Commun.* 245:641-645).

The cycling of LMW GTP-binding proteins between the GTP-bound active form and the GDP-bound inactive form is regulated by additional proteins. Guanosine nucleotide exchange factors (GEFs) increase the rate of nucleotide dissociation by several orders of magnitude, thus facilitating release of GDP and loading with GTP. The best characterized is the mammalian homologue of the *Drosophila* Son-of-Sevenless protein. Certain Ras-family proteins are also regulated by guanine nucleotide dissociation inhibitors (GDIs), which inhibit GDP dissociation. The intrinsic rate of GTP hydrolysis of the LMW GTP-binding proteins is typically very slow, but it can be stimulated by several orders of magnitude by GTPase-activating proteins (GAPs) (Geyer, M. and Wittinghofer, A. (1997) *Curr. Opin. Struct. Biol.* 7:786-792). Both GEF and GAP activity may be controlled in response to extracellular stimuli and modulated by accessory proteins such as RalBP1 and POB1. Mutant Ras-family proteins, which bind but can not hydrolyze GTP, are permanently activated, and cause cell proliferation or cancer, as do GEFs that inappropriately activate LMW GTP-binding proteins, such as the human oncogene NET1, a Rho-GEF (Drivas, G. T. et al. (1990) *Mol. Cell. Biol.*

10:1793-1798; Alberts, A. S. and Treisman, R. (1998) EMBO J. 14:4075-4085).

A novel group of GTP-binding proteins is the GTP1/OBG family, which are found in species ranging from bacteria to yeast to humans. These proteins contain characteristic GTP-binding motifs and are similar to one another but do not show sequence homology to other GTP-binding proteins.

- 5 The exact functions of these proteins are as yet uncertain, but they have been shown to be important for regulation of cell differentiation and development (Okamoto, S. and Ochi, K. (1998). Mol. Microbiol 30:107-119; Sazaka, T. et al. (1992) Biochem. Biophys. Res. Commun. 189:363-370).

The discovery of new GTPase associated proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis,

- 10 prevention, and treatment of cell proliferative, autoimmune/inflammatory, and immune system disorders.

SUMMARY OF THE INVENTION

The invention features substantially purified polypeptides, GTPase associated proteins,

- 15 referred to collectively as "GTPAP" and individually as "GTPAP-1," "GTPAP-2," "GTPAP-3," "GTPAP-4," "GTPAP-5," "GTPAP-6," "GTPAP-7," "GTPAP-8," "GTPAP-9," "GTPAP-10," "GTPAP-11," "GTPAP-12," "GTPAP-13," "GTPAP-14," "GTPAP-15," "GTPAP-16," "GTPAP-17," "GTPAP-18," "GTPAP-19," "GTPAP-20," "GTPAP-21," "GTPAP-22," "GTPAP-23," "GTPAP-24," "GTPAP-25," "GTPAP-26," "GTPAP-27," "GTPAP-28," and "GTPAP-29." In one aspect, the
20 invention provides a substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also includes a polypeptide comprising an amino acid sequence that differs by one or more conservative amino acid substitutions from an amino acid sequence selected from the group consisting of SEQ ID NO:1-29.

- The invention further provides a substantially purified variant having at least 90% amino acid
25 identity to at least one of the amino acid sequences selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also includes an isolated and purified
30 polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof.

Additionally, the invention provides an isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The

invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof.

The invention also provides a method for detecting a polynucleotide in a sample containing nucleic acids, the method comprising the steps of: (a) hybridizing the complement of the polynucleotide sequence to at least one of the polynucleotides of the sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of a polynucleotide in the sample. In one aspect, the method further comprises amplifying the polynucleotide prior to hybridization.

The invention also provides an isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:30-58 and fragments thereof. The invention further provides an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide sequence selected from the group consisting of SEQ ID NO:30-58 and fragments thereof. The invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:30-58 and fragments thereof.

The invention further provides an expression vector containing at least a fragment of the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29. In another aspect, the expression vector is contained within a host cell.

The invention also provides a method for producing a polypeptide, the method comprising the steps of: (a) culturing the host cell containing an expression vector containing a polynucleotide of the invention under conditions suitable for the expression of the polypeptide; and (b) recovering the polypeptide from the host cell culture.

The invention also provides a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

The invention further includes a purified antibody which binds to a polypeptide selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also provides a purified agonist and a purified antagonist to the polypeptide.

The invention also provides a method for treating or preventing a disorder associated with decreased expression or activity of GTPAP, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially

purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

- The invention also provides a method for treating or preventing a disorder associated with increased expression or activity of GTPAP, the method comprising administering to a subject in need of such treatment an effective amount of an antagonist of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof.

BRIEF DESCRIPTION OF THE TABLES

- Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding GTPAP.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of GTPAP.

- Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding GTPAP were isolated.

- Table 5 shows the tools, programs, and algorithms used to analyze GTPAP, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

- Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

- It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same

meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"GTPAP" refers to the amino acid sequences of substantially purified GTPAP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of GTPAP. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of GTPAP either by directly interacting with GTPAP or by acting on components of the biological pathway in which GTPAP participates.

An "allelic variant" is an alternative form of the gene encoding GTPAP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding GTPAP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as GTPAP or a polypeptide with at least one functional characteristic of GTPAP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding GTPAP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding GTPAP. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent GTPAP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of GTPAP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged -

amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of GTPAP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of GTPAP either by directly interacting with GTPAP or by acting on components of the biological pathway in which GTPAP participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind GTPAP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition containing a nucleic acid sequence which is -

complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic GTPAP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding GTPAP or fragments of GTPAP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using the XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of one or more Incyte Clones and, in some cases, one or more public domain ESTs, using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that, when made, least interfere with the properties of the original protein, i.e., the structure and especially the function of the

protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
5	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
10	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
15	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
20	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "fragment" is a unique portion of GTPAP or the polynucleotide encoding GTPAP which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment

used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:30-58 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:30-58, for example, as distinct from any other sequence in the same genome. A fragment of SEQ ID NO:30-58 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:30-58 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:30-58 and the region of SEQ ID NO:30-58 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-29 is encoded by a fragment of SEQ ID NO:30-58. A fragment of SEQ ID NO:1-29 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-29. For example, a fragment of SEQ ID NO:1-29 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-29. The precise length of a fragment of SEQ ID NO:1-29 and the region of SEQ ID NO:1-29 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the

substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequence pairs.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bi2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

Penalty for mismatch: -2

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 11

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the hydrophobicity and acidity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term "humanized antibody" refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of identity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Generally, such wash temperatures are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY;

specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_{ot} or R_{ot} analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

The terms "element" and "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term "modulate" refers to a change in the activity of GTPAP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of GTPAP.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Probe" refers to nucleic acid sequences encoding GTPAP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

"Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis et al., 1990, PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5. 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to

5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, *supra*. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding GTPAP, or fragments thereof, or GTPAP itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA,-

RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

"Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may

have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human GTPase associated proteins (GTPAP), the polynucleotides encoding GTPAP, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative, autoimmune/inflammatory, and immune system disorders.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding GTPAP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each GTPAP were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. The Incyte clones in column 5 were used to assemble the consensus nucleotide sequence of each GTPAP and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical

methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding GTPAP. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:30-58 and to distinguish between SEQ ID NO:30-58 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express GTPAP as a fraction of total tissues expressing GTPAP. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing GTPAP as a fraction of total tissues expressing GTPAP. Column 5 lists the vectors used to subclone each cDNA library. Of particular note is the specific expression of SEQ ID NO:43 in only one library, a human testis tissue library; the specific expression of SEQ ID NO:49 in only 4 libraries, one of which is associated with cell proliferation and 3 of which are associated with inflammation; and the specific expression of SEQ ID NO:40 in only 5 libraries, 3 of which are associated with cell proliferation and one of which is associated with inflammation.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding GTPAP were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

The invention also encompasses GTPAP variants. A preferred GTPAP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the GTPAP amino acid sequence, and which contains at least one functional or structural characteristic of GTPAP.

The invention also encompasses polynucleotides which encode GTPAP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:30-58, which encodes GTPAP.

The invention also encompasses a variant of a polynucleotide sequence encoding GTPAP. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding GTPAP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:30-58 which has at least about 70%, or alternatively at least about 90%, or even at least about

95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:30-58. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of GTPAP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding GTPAP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring GTPAP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode GTPAP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring GTPAP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding GTPAP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding GTPAP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode GTPAP and GTPAP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding GTPAP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:30-58 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment

of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Perkin-Elmer). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Perkin-Elmer), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding GTPAP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060.) Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been

size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode GTPAP may be cloned in recombinant DNA molecules that direct expression of GTPAP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express GTPAP.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter GTPAP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In another embodiment, sequences encoding GTPAP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; and Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively, GTPAP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of GTPAP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY.)

In order to express a biologically active GTPAP, the nucleotide sequences encoding GTPAP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding GTPAP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding GTPAP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding GTPAP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding GTPAP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding GTPAP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or

tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding GTPAP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding GTPAP can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSORT1 plasmid (Life Technologies). Ligations of sequences encoding GTPAP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of GTPAP are needed, e.g. for the production of antibodies, vectors which direct high level expression of GTPAP may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of GTPAP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *supra*; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of GTPAP. Transcription of sequences encoding GTPAP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding GTPAP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader

sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses GTPAP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of GTPAP in cell lines is preferred. For example, sequences encoding GTPAP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *apv* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system.

(See, e.g., Rhodes, C.A. (1995) *Methods Mol. Biol.* 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding GTPAP is inserted within a marker gene sequence, transformed cells containing sequences encoding GTPAP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding GTPAP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding GTPAP and that express GTPAP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of GTPAP using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on GTPAP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding GTPAP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding GTPAP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for

ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding GTPAP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode GTPAP may be designed to contain signal sequences which direct secretion of GTPAP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and W138) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding GTPAP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric GTPAP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of GTPAP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the GTPAP encoding sequence and the heterologous protein sequence, so that GTPAP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10).

A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled GTPAP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ^{35}S -methionine.

Fragments of GTPAP may be produced not only by recombinant means, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the ABI 431A peptide synthesizer (Perkin-Elmer). Various fragments of GTPAP may be synthesized separately and then combined to produce the full length molecule.

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of GTPAP and GTPase associated proteins. In addition, the expression of GTPAP is closely associated with proliferating tissues associated with cancer and fetal development, inflamed tissues, and tissues involved in the immune response. Therefore, GTPAP appears to play a role in cell proliferative, autoimmune/inflammatory, and immune system disorders. In the treatment of disorders associated with increased GTPAP expression or activity, it is desirable to decrease the expression or activity of GTPAP. In the treatment of disorders associated with decreased GTPAP expression or activity, it is desirable to increase the expression or activity of GTPAP.

Therefore, in one embodiment, GTPAP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP. Examples of such disorders include, but are not limited to, a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis,

autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis,

5 glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner

10 syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and an immune system disorder, such as acquired immunodeficiency syndrome (AIDS), X-linked agammaglobinemia of Bruton, common variable immunodeficiency (CVI), DiGeorge's syndrome (thymic hypoplasia), thymic dysplasia, isolated IgA deficiency, severe combined immunodeficiency disease (SCID),

15 immunodeficiency with thrombocytopenia and eczema (Wiskott-Aldrich syndrome), Chediak-Higashi syndrome, chronic granulomatous diseases, hereditary angioneurotic edema, and immunodeficiency associated with Cushing's disease, leukemias such as multiple myeloma, and lymphomas such as Hodgkin's disease.

In another embodiment, a vector capable of expressing GTPAP or a fragment or derivative

20 thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified GTPAP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP including, but not

25 limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of GTPAP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP including, but not limited to, those listed above.

In a further embodiment, an antagonist of GTPAP may be administered to a subject to treat or

30 prevent a disorder associated with increased expression or activity of GTPAP. Examples of such disorders include, but are not limited to, those cell proliferative, autoimmune/inflammatory, and immune system disorders described above. In one aspect, an antibody which specifically binds GTPAP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for

bringing a pharmaceutical agent to cells or tissues which express GTPAP.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding GTPAP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of GTPAP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of GTPAP may be produced using methods which are generally known in the art. In particular, purified GTPAP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind GTPAP. Antibodies to GTPAP may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with GTPAP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to GTPAP have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of GTPAP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to GTPAP may be prepared using any technique which provides for

the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J.*

Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and

Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce GTPAP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for GTPAP may also be generated. For example, such fragments include, but are not limited to, $F(ab')_2$ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the $F(ab')_2$ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between GTPAP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering GTPAP epitopes is generally used, but a competitive binding assay may also be employed (Pound, *supra*).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for GTPAP. Affinity is expressed as an

association constant, K_a , which is defined as the molar concentration of GTPAP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple GTPAP epitopes, represents the average affinity, or avidity, of the antibodies for GTPAP. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular GTPAP epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the GTPAP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of GTPAP, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington, DC; Liddell, J.E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of GTPAP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding GTPAP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding GTPAP may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding GTPAP. Thus, complementary molecules or fragments may be used to modulate GTPAP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding GTPAP.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides

encoding GTPAP. (See, e.g., Sambrook, *supra*; Ausubel, 1995, *supra*.)

Genes encoding GTPAP can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding GTPAP. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding GTPAP. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may be employed. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, *Molecular and Immunologic Approaches*, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding GTPAP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques

for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding GTPAP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA
5 constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase
10 linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable
15 for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat.
20 Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical
25 or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of GTPAP, antibodies to GTPAP, and mimetics, agonists, antagonists, or inhibitors of GTPAP. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical
30 carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial,

intraductal, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration.

Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's

102250.695201
0356679.05201

solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of GTPAP, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example GTPAP or fragments thereof, antibodies of GTPAP, and agonists, antagonists or inhibitors of GTPAP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be

determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about $0.1 \mu\text{g}$ to $100,000 \mu\text{g}$, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind GTPAP may be used for the diagnosis of disorders characterized by expression of GTPAP, or in assays to monitor patients being treated with GTPAP or agonists, antagonists, or inhibitors of GTPAP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for GTPAP include methods which utilize the antibody and a label to detect GTPAP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring GTPAP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of GTPAP expression. Normal or standard values for GTPAP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to

GTPAP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of GTPAP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding GTPAP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of GTPAP may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess

expression of GTPAP, and to monitor regulation of GTPAP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding GTPAP or closely related molecules may be used to identify nucleic acid sequences which encode GTPAP. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding GTPAP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the GTPAP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:30-58 or from genomic sequences including promoters, enhancers, and introns of the GTPAP gene.

Means for producing specific hybridization probes for DNAs encoding GTPAP include the cloning of polynucleotide sequences encoding GTPAP or GTPAP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding GTPAP may be used for the diagnosis of disorders associated with expression of GTPAP. Examples of such disorders include, but are not limited to, a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and an immune system disorder, such as acquired immunodeficiency syndrome (AIDS), X-linked agammaglobinemia of Bruton, common variable immunodeficiency (CVI), DiGeorge's syndrome (thymic hypoplasia), thymic dysplasia, isolated IgA deficiency, severe combined immunodeficiency disease (SCID), immunodeficiency with thrombocytopenia and eczema (Wiskott-Aldrich syndrome), Chediak-Higashi syndrome, chronic granulomatous diseases, hereditary angioneurotic edema, and immunodeficiency associated with Cushing's disease, leukemias such as multiple myeloma, and lymphomas such as Hodgkin's disease. The polynucleotide sequences encoding GTPAP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered GTPAP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding GTPAP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding GTPAP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding GTPAP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of GTPAP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding GTPAP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding GTPAP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a

polynucleotide encoding GTPAP, or a fragment of a polynucleotide complementary to the polynucleotide encoding GTPAP, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

5 Methods which may also be used to quantify the expression of GTPAP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer of interest is
10 presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify
15 genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) *Proc. Natl. Acad. Sci.*
20 USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

In another embodiment of the invention, nucleic acid sequences encoding GTPAP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence.

25 The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355; Price, C.M. (1993) *Blood Rev.* 7:127-134; and Trask, B.J. (1991) *Trends Genet.* 7:149-154.)

30 Fluorescent *in situ* hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, *supra*, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the

location of the gene encoding GTPAP on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

5 In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides
10 valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention
15 may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, GTPAP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a
20 solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between GTPAP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are
25 synthesized on a solid substrate. The test compounds are reacted with GTPAP, or fragments thereof, and washed. Bound GTPAP is then detected by methods well known in the art. Purified GTPAP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

30 In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding GTPAP specifically compete with a test compound for binding GTPAP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with GTPAP.

In additional embodiments, the nucleotide sequences which encode GTPAP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. Nos. 60/109,592, 60/118,610, and 60/127,990 are hereby expressly incorporated by reference.

EXAMPLES

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the

recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-

1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), or pINCY (Incyte Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into competent E. coli cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

cDNA sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Perkin-Elmer) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer). Electrophoretic separation of cDNA sequencing reactions and detection of labeled

polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Perkin-Elmer) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example V.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene

families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:30-58. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding GTPAP occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories.

Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Extension of GTPAP Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:30-58 were produced by extension of

an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham

Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

In like manner, the nucleotide sequences of SEQ ID NO:30-58 are used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

VI. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:30-58 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 µCi of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁶ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and

compared.

VII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, *supra*.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

VIII. Complementary Polynucleotides

Sequences complementary to the GTPAP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring GTPAP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of GTPAP. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the GTPAP-encoding transcript.

IX. Expression of GTPAP

Expression and purification of GTPAP is achieved using bacterial or virus-based expression

systems. For expression of GTPAP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express GTPAP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of GTPAP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant *Autographica californica* nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding GTPAP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect *Spodoptera frugiperda* (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, GTPAP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from *Schistosoma japonicum*, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from GTPAP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10 and 16). Purified GTPAP obtained by these methods can be used directly in the following activity assay.

X. Demonstration of GTPAP Activity

The role of GTPAP can be assayed *in vitro* by monitoring the mobilization of Ca^{++} as part of the signal transduction pathway. (See, e.g., Grynkievich, G. et al. (1985) J. Biol. Chem. 260:3440; McColl, S. et al. (1993) J. Immunol. 150:4550-4555; and Aussel, C. et al. (1988) J. Immunol. 140-215.) The assay requires preloading neutrophils or T cells with a fluorescent dye such as FURA-2.

Upon binding Ca^{++} , FURA-2 exhibits an absorption shift that can be observed by scanning the excitation spectrum between 300 and 400 nm, while monitoring the emission at 510 nm. When the cells are exposed to one or more activating stimuli artificially (i.e., anti-CD3 antibody ligation of the T cell receptor) or physiologically (i.e., by allogeneic stimulation), Ca^{++} flux takes place. Ca^{++} flux results from the release of Ca^{++} from intracellular organelles or from Ca^{++} entry into the cell through activated Ca^{++} channels. This flux can be observed and quantified by assaying the cells in a fluorometer or fluorescence activated cell sorter. Measurements of Ca^{++} flux are compared between cells in their normal state and those preloaded with GTPAP. Increased mobilization attributable to increased GTPAP availability results in increased emission.

Alternatively, GTPAP activity is measured by quantifying the amount of a non-hydrolyzable GTP analogue, GTP γ S, bound over a 10 minute incubation period. Varying amounts of GTPAP are incubated at 30°C in 50mM Tris buffer, pH 7.5, containing 1mM dithiothreitol, 1mM EDTA and 1 μ M [35 S]GTP γ S. Samples are passed through nitrocellulose filters and washed twice with a buffer consisting of 50mM Tris-HCl, pH 7.8, 1mM NaN_3 , 10mM MgCl_2 , 1mM EDTA, 0.5mM dithiothreitol, 0.01mM PMSF, and 200mM NaCl. The filter-bound counts are measured by liquid scintillation to quantify the amount of bound [35 S]GTP γ S. GTPAP activity may also be measured as the amount of GTP hydrolysed over a 10 minute incubation period at 37°C. GTPAP is incubated in 50mM Tris-HCl buffer, pH 7.8, containing 1mM dithiothreitol, 2mM EDTA, 10 μ M [α - 32 P]GTP, and 1 μ M H-rab protein. GTPase activity is initiated by adding MgCl_2 to a final concentration of 10 mM. Samples are removed at various time points, mixed with an equal volume of ice-cold 0.5mM EDTA, and frozen. Aliquots are spotted onto polyethyleneimine-cellulose thin layer chromatography plates, which are developed in 1M LiCl, dried, and autoradiographed. The signal detected is proportional to GTPAP activity.

Alternatively, GTPAP activity may be demonstrated as the ability to interact with its associated G α or LMW GTPase in an in vitro binding assay. The candidate GTPases are expressed as fusion proteins with glutathione S-transferase (GST), and purified by affinity chromatography on glutathione-Sepharose. The GTPases are loaded with GDP by incubating 20 mM Tris buffer, pH 8.0, containing 100 mM NaCl, 2 mM EDTA, 5 mM MgCl_2 , 0.2 mM DTT, 100 μ M AMP-PNP and 10 μ M GDP at 30°C for 20 minutes. GTPAP is expressed as a FLAG fusion proteins in a baculovirus system. Extracts of these baculovirus cells containing GTPAP-FLAG fusion proteins are precleared with GST beads, then incubated with GST-GTPase fusion proteins. The complexes formed are precipitated by glutathione-Sepharose and separated by SDS-polyacrylamide gel electrophoresis. The separated proteins are blotted onto nitrocellulose membranes and probed with commercially available anti-

FLAG antibodies. GTPAP activity is proportional to the amount of GTPAP-FLAG fusion protein detected in the complex.

XI. Functional Assays

GTPAP function is assessed by expressing the sequences encoding GTPAP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT (Life Technologies) and pCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP: Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of GTPAP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding GTPAP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding GTPAP and other genes of interest can be analyzed by northern analysis or microarray techniques.

XII. Production of GTPAP Specific Antibodies

GTPAP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the GTPAP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-GTPAP activity by, for example, binding the peptide or GTPAP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIII. Purification of Naturally Occurring GTPAP Using Specific Antibodies

Naturally occurring or recombinant GTPAP is substantially purified by immunoaffinity chromatography using antibodies specific for GTPAP. An immunoaffinity column is constructed by covalently coupling anti-GTPAP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing GTPAP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of GTPAP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/GTPAP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and GTPAP is collected.

XIV. Identification of Molecules Which Interact with GTPAP

GTPAP, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled GTPAP, washed, and any wells with labeled GTPAP complex are assayed. Data obtained using different concentrations of GTPAP are used to calculate values for the number, affinity, and association of

GTPAP with the candidate molecules.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention.

Although the invention has been described in connection with certain embodiments, it should be

5 understood that the invention as claimed should not be unduly limited to such specific embodiments.

Indeed, various modifications of the described modes for carrying out the invention which are

obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	30	708398	SYNORAT04	568987X31 (MWLR3DT01), 708398X11, 708398X15, 708398X16, 708398X17, and 708398X21 (SYNORAT04), 2170523F6 (FENCNOT03), 3374750H1 (CONUTUT05)
2	31	1259937	MENITUT03	913652R6 (STONNOT02), 1259937F6 and 1259937H1 (MENITUT03), 1476721F1 (CORENOT02), 1729248F6 (BRSTTUT08), 2191963H1 (THVSTUT03), 3129757F6 (LUNGUTUT12), 3268746X15F1 (BRAINOT20), 3428891F6 (SKLNNOT04)
3	32	1452285	PENITUT01	1452285F6 and 1452285H1 (PENITUT01), 2605011H1 (LUNGUTUT07), 3505135H1 (ADRENUT11)
4	33	1812894	PROSTUT12	1812894H1, 1812894X12 and 1809113T6 (PROSTUT12), 190479F6 (OVARNOT07), 2232535X15F1 and 2232535X18F1 (PROSNOT16), 2267486X16C1 (UTRSNOT02), 2508562F6 (CONUTUT01)
5	34	3074884	BONEUNT01	225362F1 (PANCNOT01), 900707R1 (BRSTTUT03), 1339234F6 (COLINTUT03), 1759046R6 (PTFUNG03), 3074884H1 (BONEUNT01), SBA02767F1
6	35	3452277	UTRSNOT03	168453F6 (PROSNOT15), 1951534H1 (PTFUNG01), 3452277H1 (UTRSNOT03), 4092781T6 (BSCNSZT01), SBA01413F1, SBA03044F1, SBA01805F1
7	36	4203832	BRAITUT29	723394F1 (SYNOOAT01), 862290R1, and 862290T1 (BRAITUT03), 1560918F1 (SPLNNOT04), 3509241H1 (CONCNOT01), 4203832H1 (BRAITUT29)
8	37	104368	BMARNOT02	104368H1 (BMARNOT02), SAEA03574F1, SAEA01063F1, SAEA00392F1, SAEA02287F1
9	38	1441680	THYRNUT03	1441680F6, 1441680H1, and 1441680T6 (THYRNUT03), 1904222F6 (OVARNOT07), 2477983F6 (SMCANOT01)
10	39	1494955	PROSNON01	965986R1 (BRSTNOT05), 1429037F1 and 1429037T1 (SINTBST01), 1453487F6 (PENITUT01), 1486114H1 (CORENOT02), 1494955H1 (PROSNON01), 1995426R6 (BRSTTUT03), 2112074X18F1 and 2112348R6 (BRAITUT03)
11	40	1508161	LUNGNOT14	1508161F6 and 1508161H1 (LUNGNOT14), 3334303H1 (BRAIFET01), 4755656H1 (BRAHNOT01)
12	41	1811877	PROSTUT12	493795H1 (HWT2NOT01), 1573136H1 (LUNGNOT03), 1811877F6 and 1811877H1 (PROSTUT12), 1825223F6 (LSURNUT03), 2454143H1 (ENDANOT01), 2651022H1 (BLAUTUT08), 3487062H1 (EPIGNOT01), 4536331H1 (OVARNOT12), 4795253H1 (LIVRTUT09), 4854087H1 (TESTNOT10), 4906149H2 (TLYMNUT08), 5196386H1 (LUNLUTUT04)

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
13	42	1848674	LUNGFEF03	1574127F6, 3857867X306F1, and 3857867X313F1 (LNDNOT03), 1848674H1 (LUNGFEF03), 1877170F6 (LEUNNOT03), 2695307H1 (UTRSNOT12), 4148654H1 (SINITUT04), 4984182H1 (HELATXT05), 5288671H1 (LIVTUS02)
14	43	2012970	TESTNOT03	2012970H1, 2012970R6, 2012970X11F (TESTNOT03)
15	44	2254315	OVARTUT01	022341F1 (ADEINNOT01), 198478R6 (KIDNOT02), 2254315H1 (OVARTUT01), 2370170F6 (ADRENUT07), 245178F6 (ENDANOT01)
16	45	2415545	HNT3AZT01	775722H1 (COLANOT05), 870320R6 (LUNGAST01), 889023R1 (STOMTUT01), 895724R1 (BRSTNOT05), 1398541F1 (BRAITUT08), 1662585F6 (BRSTNOT09), 2415545H1 (HNT3AZT01), 2985066H1 (CARGDIT01), 3462702H1 (293TF2T01)
17	46	2707969	PONSZAT01	282552R1, 282552X23, and 282552X7 (CARDNOT01), 889783R1 (STOMTUT01), 1995451R6 (BRSTTUT03), 2707969H1 (PONSZAT01), SAAC00359R1.comp, SAAB00136R1, SAAC00330R1
18	47	2817769	BRSTNOT14	041660R1 (TBLYNOT01), 077378R1 (SYNORAB01), 740028R1 (PANCNOT04), 1593201P6 (BRAINOT14), 1924025R6 (BRSTTUT01), 2817769H1 (BRSTNOT14)
19	48	2817557	THIMFET03	473002F1 and 473002R1 (MMLRIDT01), 690999R6 (LUNGUT02), 997483R1 (KIDNOT01), 1430662P6 (SINVEST01), 1514017F1 (PANCUT01), 1740475R6 (HIPONON01), 2109547H1 (BRAITUT03), 2917557H1 (THIMFET03), 4309528H1 (BRAUNOT01), 4990135H1 (LIVTUT01)
20	49	3421335	UCMNOT04	777588R6 and 777588T6 (COLANOT05), 3421335H1 (UCMNOT04)
21	50	605761	BRSTTUT01	605761F1, 605761H1, and 605761R6 (BRSTTUT01), 1271131X15 (TESTTUT02), 1516985F1 (PANCUT01), 1524935H1 (UCMCL5T01), 2234646F6 (PANCUT02)
22	51	483862	HNT2RAT01	483862H1 and 483862R1 (HNT2RAT01), 1750781X305F1, 1750781X307D2 (LIVTUT01)
23	52	1256777	MENITUT03	264041R6 (HNT2AGT01), 826449R1 (PROSNOT06), 1256777H1 (MENITUT03), 2276061R6 (PROSNON01), 4614049H1 (BRAHNOT01)
24	53	2198779	SPLNPFET02	1557708F6 (BLAUTUT04), 1922490R6 (BRSTTUT01), 2198779H1 (SPLNPFET02), 2541193P7 (BONUT01), 3039254F6 (BRSTNOT16), 3057079H1 (LNDNOT08), 3105017H1 (COLNUCT03), 4239592H1 (SYNMDIT01), 5064513H1 (ARTFDT01)

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
25	54	2226116	SENVNOT01	1662607F6 (BRSTNOT09), 1662607T6 (BRSTNOT09), 2226116F6 (SENVNOT01), 2226116H1 (SENVNOT01), 2930011F6 (TYLNNOT04), 3015747T6 (MUSCNOT07), 4087670H1 (LIVNOT06)
26	55	2504472	CONUTUT01	420365P1 (BRSTNOT01), 762246R1 (BRAITUT02), 907754R2 (COLNNOT09), 1007508H1 (HEALDIT02), 1302342F6 (PLACNOT02), 1913887H1 (PROSTUT04), 2023822P6 (CONNOT01), 2023822X1R1 (CONNOT01), 2504472H1 (CONUTUT01), 2951618F6 (KIDNFET01)
27	56	3029920	HEARFET02	354846T6 (RATRNUT01), 418533R6 (BRSTNOT01), 935073R1 (CERVNOT01), 1340722P1 (COLNUT03), 1416203T6 (BRAINT02), 1524567F1 (UCMCLST01), 1773043H1 (MENTUN03), 2590310H2 (LUNGNOT22), 3029920H1 (HEARFET02), 4873053H1 (COLDNOT01), 5687696H1 (BRAINTUT01)
28	57	3332415	BRAIFET01	118166R1 (MUSCNOT01), 1257348H1 (MENITUT03), 1288237T6 (BRAINTUT1), 1335936F6 (COLANNOT13), 1452268H1 (PENITUT01), 1996016R6 (BRSTTUT03), 2116665R6 (BRSTTUT02), 2206894F6 (SINTFET03), 2540063H1 (BONRTUT01), 2808268H1 (BLADTUT08), 3086221H1 (HEAGNOT03), 3127508H1 (LUNGTFUT12), 3295812H1 (PLXJINT01), 3332415H1 (BRAIFET01), 3604705H1 (LUNGNOT30), 4821203H1 (PROSTUT17), 4970353H1 (KIDEUNC10), 5055775H1 (COLATUT01)
29	58	4031536	BRAINT03	029167X3 (SPLNFET01), 350137R1 (LVENNOT01), 408825X1 (EOSIHET02), 689446X23 (LUNGTFUT02), 1963062R6 (BRSTNOT04), 2288043R6 (BRAINO01), 4031536H1 (BRAINT03)

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
1	1002	T30 S224 T405 S499 T533 S558 S701 T737 T845 S864 S6 T152 T268 T412 T442 T464 T514 T528 T693 S814 S815 S823 T880 Y117 Y842 S21 S77 T86 S200 T246 T299 S77 S306 Y131	N446	G524-T531: ATP/GTP- binding site motif	GTP-binding protein [Mus musculus] g53169	BLAST MOTIFS
2	338		N244		CAMP- regulated Guanine nucleotide exchange factor [Rattus norvegicus] g4079657	BLAST
3	211	S159 S199	N33 N74	G16-T23: ATP/GTP- binding site motif	GTP-binding protein [Rattus norvegicus] g206543	BLAST MOTIFS PFAM BLOCKS PRINTS
4	516	T14 S42 T237 S270 S347 S360 T371 T395 T433 S500 T3 S13 S96 T316 S430			Fos-related antigen [Rattus norvegicus] g1016712 Rabaptin-4 [H. sapiens] g3832516	BLAST MOTIFS
5	445	T44 T114 T219 T297 S314 S341 S356 T412 T24 S72 T91 T328 T388 T394		G230-T237: ATP/GTP- binding site motif	GTP-binding protein [H. sapiens] g2765411	BLAST MOTIFS

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
6	445	S174 S202 S289 S29 S305 S323 T434 T11 T147 T197 T198 S270 S273 S371 S397 Y125	N73		Regulator of G-protein signaling-9 [H. sapiens] g3284012	BLAST
7	281	S182 S210 S254 S13 T56 S110 S182 S32 T46 S66 S177	N130 N181	G31-T38:ATP/GTP- binding site motif	Putative ras- like protein [H. sapiens] g4092830	MOTIFS PRINTS BLAST PFAM
8	301	S92 T2 T3 Y15 S18 S19 S20 S25 S97 T120 S165 S296 T94 S116 T120 S284		E47-G66, S116-E178, Y188-G272: Phosducin signature	Phosducin- like protein [Rattus rattus] g1323727	MOTIFS BLAST PRINTS
9	485	T6 Y57 S82 T91 S112 S187 T231 T257 S309 T6 T81 S132 S157 S210 S241 T462	N460	L49-S82: Beta G protein	Similar to WD domain Beta transducin- like protein [C. elegans] g5596646	MOTIFS BLAST PRINTS
10	447	S420 S94 T107 S118 T167 T179 T308 S390 S39 S58 T78 T113 S129 T160 T167 Y174 T199 S216 S291 T302 T323 T359 T384 S423 T438	N76 N92 N231 N289 N378 N421	M294-T308: Beta transducin	WS beta- transducin repeat protein [Homo sapiens] g4704417	MOTIFS BLAST
11	199	S90 T55 T140 S190		K6-E130: Ras Guanine exchange factor	Putative guanine nucleotide releasing factor [Drosophila affinis] g2981229	MOTIFS BLAST PFAM

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
12	694	S57 S67 S99 T150 T346 S416 S467 S500 T522 T684 S99 T156 S209 S285 T331 T360 T388 T430 T477 T650 T688		L10-I24, M96-L110: Beta transducin	Transducin- like protein [H. sapiens] g414536	MOTIFS BLAST
13	654	T10 S15 T49 S97 S102 S104 S112 S113 S377 S432 S638 T46 S54 S84 S97 T177 S217 T307 S401 S450 S504 T515 S546 T547 S561 Y618	N353 N362 N502	L197-F211: Beta transducin	Similar to the beta transducin family [C. elegans] g2315521	MOTIFS BLAST
14	180	S14		G23-S30: ATP-GTP binding site	Rab7C (small GTP binding protein) [Lotus japonicus] g1370186	MOTIFS BLAST
15	374	T100 T249 S260 T308 T328 S338 S351 S30 T73 T157 S237 T308	N114 N189 N222	G26-T33: ATP-GTP binding site	ATP(GTP)- binding protein [H. sapiens] g3646130	MOTIFS BLAST
16	649	S67 T344 S366 S63 S68 S75 S122 S177 S265 T282 T332 S373 S380 S563 T569 S634 S20 T94 S128 S314 T382 T385 T458 T559		F307-S544: Probable RABCAP domain	Similar to probable RABCAP [C. elegans] g3925265	MOTIFS BLAST PFAM
17	698	T244 S262 S17 T41 T42 T196 S206 S317 S479 S522 S556 T586 T680 T31 S95 T99 T140 T173 S257 T322 S374 T450 S568 T619	N171 N194 N685		Small GTP- binding protein associated protein [Mus musculus] g725274	MOTIFS BLAST

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
18	396	T325 S115 T133 S232 S275 T336 S22 T221 S232 T320	N60 N230 N286	G29-S36: ATP-GTP binding site	Putative GTP- binding protein [C. elegans] g3880615	MOTIFS BLAST
19	634	T197 S3 S5 S9 T14 S132 T197 T285 T553 T40 T56 S160 T189 S261 S582 Y20 Y396 Y419		G52-T59: ATP-GTP binding site	Putative GTP- binding protein [H. sapient] g3169010	MOTIFS BLAST
20	196	T60 S73 S90 S99 S73 S193		G19-T26: ATP-GTP binding site	Kidney injury associated protein H0052 Acc No W86322 ADP- ribosylation factor-like protein 3 [Rattus norvegicus] g560006	MOTIFS BLAST
21	446	T10 T24 T91 S122 T243 S263 S270 T305 S317 S325 T357 S372 T379 S100 S170 S223 T227 S285 T348	N79	L323-L337: Beta transducin	Putative WD40 repeat protein [A. thaliana] g4191784	MOTIFS BLAST
22	265	T184 T76 T137 S139 T161 T174 T183 S213	N159	L141, L148, L155 L: zipper gene regulatory motif	TipD: similar to beta transducin family [D. discoideum] g2407788	MOTIFS BLAST

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
23	185	T55 S111 S127 S148 S171 S14 S94 Y103		G10-T17: ATP/GTP binding site (P- loop) A4-S72: Ras domain		MOTIFS PFAM PRINTS
24	554	S388 T488 S30 S75 T111 S149 S220 S237 T255 S305 S325 T339 T359 S363 S509 S172 T195 S211 T378 T438 T470 Y203	N5	N297-D336, P345- D383, G481-Q519: Beta-transducin WD40 repeats	WD-repeat protein [Arabidopsis thaliana] g324603	BLAST MOTIFS PFAM PRINTS
25	434	S164 S341 T347 S36 S68 S92 T286 S364	N22 N383	G259-S266: ATP/GTP binding site (P- loop): G113-R433: GTP1/OBG domain	Predicted GTP binding protein [C. elegans] g3878629	BLAST MOTIFS PFAM BLOCKS PRINTS
26	826	S122 T243 T247 T427 S454 S519 T528 S623 S701 S715 S809 T88 S143 S266 T411 S505 S577 S603 T661 S735 T753 S791 T815	N23 N264 N576 N600 N789	R48-E91, L97-S143, F197 K237, V273- W319, W378-R416, W604 K642, A659- G697: Beta- transducin WD40 repeats	Predicted WD repeat protein [S. cerevisiae] F42935	BLAST MOTIFS PFAM PRINTS
27	618	T414 S59 T105 S126 T139 T143 S196 T203 S311 S325 T370 T390 S477 T483 S541 T583 T94 S148 T247 Y160 Y383 Y456	N118 N154 N346	G11-T18, G425-S432: ATP/GTP binding site (P-loop) R6-K187: Ras domain	GTP-binding protein APD08 [H. sapiens] Accession W5771	BLAST MOTIFS PFAM PRINTS

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
28	596	S17 S21 S50 S152 S153 T533 S539 T594 S36 S38 S80 T163 T169 S183 S211 T240 S306 T329 T417 S457 S508 T545 S45 T64 S88 T124 S139 S299 S451 S459 S528 S568 Y180 Y364		A178-L355: Rho- family guanine nucleotide exchange factor (RhoGEF) domain	Guanine nucleotide regulatory protein (NET1 homologue) [Mus musculus] g3834631	BLAST MOTIFS PFAM BLOCKS
29	589	T108 S20 T90 S127 S176 S278 S467 T521 S522 T189 S254 T284 T292 T321 T324 T345 T364 T423 S444 T484 T527	N572	L252-S289, G293- N329, G333-D369, G373-D409, E413- D449, G453-D489, G493-D532: Beta- transducin WD40 repeats R160-K206: F-box domain	SEM-10 [C.elegans] g2677836	BLAST MOTIFS PFAM PRINTS

Table 3

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
30	628-711	Reproductive (0.256) Nervous (0.154) Gastrointestinal (0.154)	Cell Proliferation (0.592) Inflammation (0.372)	PSPORT1
31	1094-1129	Reproductive (0.268) Cardiovascular (0.146) Nervous (0.146)	Cell Proliferation (0.731) Inflammation (0.219) Neurological (0.049)	pINCY
32	652-703	Cardiovascular (0.375) Reproductive (0.375) Dermatologic (0.125) Endocrine (0.125)	Cell Proliferation (0.875) Trauma (0.125)	pINCY
33	1224-1292	Reproductive (0.412) Gastrointestinal (0.147) Hematopoietic/Immune (0.147)	Cell Proliferation (0.647) Inflammation (0.264)	pINCY
34	16-65	Nervous (0.211) Reproductive (0.197) Gastrointestinal (0.169)	Cell Proliferation (0.507) Inflammation (0.352)	pINCY
35	947-1043	Reproductive (0.444) Nervous (0.333) Gastrointestinal (0.111) Urologic (0.111)	Cell Proliferation (0.667) Inflammation (0.111) Neurological (0.111)	pINCY
36	840-1001	Nervous (0.340) Reproductive (0.208) Gastrointestinal (0.151)	Cell Proliferation (0.641) Inflammation (0.302) Neurological (0.038)	pINCY

Table 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector	
				PBUESCRIPT	
37	507-551	Hematopoietic/Immune (0.269) Nervous (0.269) Reproductive (0.154)	Inflammation (0.423) Cell Proliferation (0.269)	PBUESCRIPT	
38	218-262	Cardiovascular (0.357) Nervous (0.214) Gastrointestinal (0.143)	Cell Proliferation (0.572) Inflammation (0.214)	PINCY	
39	164-208	Nervous (0.280) Reproductive (0.260) Developmental (0.120)	Cell Proliferation (0.740) Inflammation (0.180)	PSFORT1	
40	369-411	Cardiovascular (0.250) Developmental (0.250) Gastrointestinal (0.250)	Cell Proliferation (0.500) Inflammation (0.250)	PINCY	
41	272-316	Reproductive (0.392) Gastrointestinal (0.118) Hematopoietic/Immune (0.118)	Cell Proliferation (0.626) Inflammation (0.137)	PINCY	
42	664-708	Nervous (0.211) Reproductive (0.211) Cardiovascular (0.158)	Cell Proliferation (0.614) Inflammation (0.281)	PINCY	
43	226-270	Reproductive (1.000)	Inflammation (1.000)	PBUESCRIPT	
44	11-55	Reproductive (0.254) Gastrointestinal (0.206) Cardiovascular (0.159)	Cell Proliferation (0.698) Inflammation (0.206)	PSFORT1	

Table 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
45	637-681	Reproductive (0.281) Nervous (0.188) Gastrointestinal (0.156)	Cell Proliferation (0.781) Inflammation (0.234)	pINCY
46	1016-1060	Nervous (0.330) Reproductive (0.183) Hematopoietic/Immune (0.122)	Cell Proliferation (0.582) Inflammation (0.235)	pINCY
47	737-781	Nervous (0.218) Reproductive (0.188) Gastrointestinal (0.158)	Cell Proliferation (0.655) Inflammation (0.211)	pINCY
48	469-513	Reproductive (0.222) Hematopoietic/Immune (0.160) Nervous (0.160)	Cell Proliferation (0.543) Inflammation (0.272)	pINCY
49	226-270	Gastrointestinal (0.333) Hematopoietic/Immune (0.333) Reproductive (0.333)	Inflammation (1.000)	pINCY
50	456-500	Reproductive (0.289) Gastrointestinal (0.133) Hematopoietic/Immune (0.133)	Cell Proliferation (0.778) Inflammation (0.156)	PSFORT1
51	252-296	Nervous (0.500) Gastrointestinal (0.200) Cardiovascular (0.100)	Cell Proliferation (1.000) Inflammation (0.200)	PBLUESCRIPT
52	60-104	Nervous (0.326) Reproductive (0.326) Cardiovascular (0.152)	Cell proliferation (0.565) Inflammation (0.369)	pINCY

Table 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
53	488-532	Reproductive (0.232) Nervous (0.195) Hematopoietic/Immune (0.146)	Cell proliferation (0.622) Inflammation (0.427)	pINCY
54	686-730	Reproductive (0.250) Gastrointestinal (0.150) Hematopoietic/Immune (0.150)	Cell proliferation (0.700) Inflammation (0.400)	pINCY
55	543-587 1299-1343	Reproductive (0.282) Nervous (0.155) Gastrointestinal (0.146)	Cell proliferation (0.592) Inflammation (0.359)	pINCY
56	345-389 792-836	Nervous (0.268) Reproductive (0.169) Cardiovascular (0.113) Hematopoietic/Immune (0.113)	Cell proliferation (0.606) Inflammation (0.296)	pINCY
57	163-207	Reproductive (0.270) Gastrointestinal (0.189) Nervous (0.156)	Cell proliferation (0.705) Inflammation (0.254)	pINCY
58	381-425 726-770	Nervous (0.317) Reproductive (0.250) Gastrointestinal (0.117)	Cell proliferation (0.450) Inflammation (0.283)	pINCY

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
30	SYNORAT04	This library was constructed using RNA isolated from the wrist synovial membrane tissue of a 62-year-old female with rheumatoid arthritis.
31	MENITUT03	This library was constructed using RNA isolated from brain meningioma tissue removed from a 35-year-old female during excision of a cerebral meningial lesion. Pathology indicated a benign neoplasm in the right cerebellopontine angle of the brain. Patient history included hypothyroidism. Family history included myocardial infarction and breast cancer.
32	PENITUT01	This library was constructed using RNA isolated from tumor tissue removed from the penis of a 64-year-old male during penile amputation. Pathology indicated a fungating invasive grade 4 squamous cell carcinoma involving the inner wall of the foreskin and extending onto the glans penis. Patient history included benign neoplasm of the large bowel, atherosclerotic coronary artery disease, angina pectoris, gout, and obesity. Family history included malignant pharyngeal neoplasm, chronic lymphocytic leukemia, and chronic liver disease.
33	PROSTUT12	This library was constructed using RNA isolated from prostate tumor tissue removed from a 65-year-old male during a radical prostatectomy. Pathology indicated an adenocarcinoma (Gleason grade 2+2). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA).
34	BONEUNT01	This library was constructed using RNA isolated from Saos-2, a primary osteogenic sarcoma cell line (ATCC HTB-85) derived from an 11-year-old Caucasian female.
35	UTERSON03	This library was constructed from 6.4 million independent clones from a uterine library. RNA for these libraries was isolated from uterine myometrial tissue removed from a 41-year-old female during a vaginal hysterectomy with dilation and curettage. The endometrium was secretory and contained fragments of endometrial polyps. Benign endo- and ectocervical mucosa were identified in the endocervix. Pathology for the associated tumor tissue indicated uterine leiomyoma. The normalization and hybridization conditions were adapted from Soares et al. (Proc.Natl.Acad.Sci. USA (1994) 91:9928).
36	BRAITUT29	This library was constructed using RNA isolated from brain tumor tissue removed from the parietal lobe of a 43-year-old female during excision of a cerebral meningial lesion. Pathology indicated high grade glioma. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, benign hypertension, and hyperlipidemia.
37	BMARNOT02	This library was constructed using RNA isolated from the bone marrow of 24 male and female Caucasian donors, 16 to 70 years old. (RNA came from Clontech.)

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
38	THYRNOT03	This library was constructed using RNA isolated from thyroid tissue removed from the left thyroid of a 28-year-old Caucasian female during a complete thyroidectomy. Pathology indicated a small nodule of adenomatous hyperplasia present in the left thyroid. Pathology for the associated tumor tissue indicated dominant follicular adenoma, forming a well-encapsulated mass in the left thyroid.
39	PROSNON01	This normalized library was constructed from 4.4 million independent clones from a prostate library. Starting RNA was made from prostate tissue removed from a 28-year-old Caucasian male who died from a self-inflicted gunshot wound. The normalization and hybridization conditions were adapted from Soares, M.B. et al. (1994) Proc. Natl. Acad. Sci. USA 91:9228-9232, using a longer (19 hour) reannealing hybridization period.
40	LUNGNOT14	This library was constructed using RNA isolated from lung tissue removed from the left lower lobe of a 47-year-old Caucasian male during a segmental lung resection. Pathology for the associated tumor tissue indicated a grade 4 adenocarcinoma, and the parenchyma showed calcified granuloma. Patient history included benign hypertension and chronic obstructive pulmonary disease. Family history included type II diabetes and acute myocardial infarction.
41	PROSTUT12	This library was constructed using RNA isolated from prostate tumor tissue removed from a 65-year-old Caucasian male during a radical prostatectomy. Pathology indicated an adenocarcinoma (Gleason grade 2+2). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostatic specific antigen (PSA).
42	LUNGFET03	This library was constructed using RNA isolated from lung tissue removed from a Caucasian female fetus who died at 20 weeks' gestation.
43	TESTNOT03	This library was constructed using RNA isolated from testicular tissue removed from a 37-year-old Caucasian male, who died from liver disease. Patient history included cirrhosis, jaundice, and liver failure.
44	OVARTUT01	This library was constructed using RNA isolated from ovarian tumor tissue removed from a 43-year-old Caucasian female during removal of the fallopian tubes and ovaries. Pathology indicated grade 2 mucinous cystadenocarcinoma involving the entire left ovary. Patient history included mitral valve disorder, pneumonia, and viral hepatitis. Family history included atherosclerotic coronary artery disease, pancreatic cancer, stress reaction, cerebrovascular disease, breast cancer, and uterine cancer.
45	HNT3AZT01	This library was constructed using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated for three days with 0.35 micromolar 5-aza-2'-deoxycytidine (AZ).

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
46	PONS2AT01	This library was constructed using RNA isolated from diseased pons tissue from the brain of a 74-year-old Caucasian male who died from Alzheimer's disease.
47	BRSTNOT14	This library was constructed using RNA isolated from breast tissue obtained from a 62-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology for the associated tumor tissue indicated an invasive grade 3 (of 4), nuclear grade 3 (of 3) adenocarcinoma, ductal type. Patient history included a benign colon neoplasm, hyperlipidemia, cardiac dysrhythmia, and obesity. Family history included atherosclerotic coronary artery disease, myocardial infarction, colon cancer, ovarian cancer, lung cancer, and cerebrovascular disease.
48	THYMPET03	This library was constructed using RNA isolated from thymus tissue removed from a Caucasian male fetus.
49	UCHCNOT04	This library was constructed using RNA isolated from mononuclear cells obtained from the umbilical cord blood of multiple individuals of mixed age and sex. The cells were treated with G-CSF.
50	BRSTTUT01	This library was constructed using RNA isolated from breast tumor tissue removed from a 55-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated invasive grade 4 mammary adenocarcinoma of mixed lobular and ductal type, extensively involving the left breast. Family history included benign hypertension, atherosclerotic coronary artery disease, cerebrovascular disease, and depressive disorder.
51	HNT2PAT01	This library was constructed at Stratagene (STR037231), using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated with retinoic acid for 24 hours.
52	MENITUT03	This library was constructed using RNA isolated from brain meningioma tissue removed from a 35-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated a benign neoplasm in the right cerebellopontine angle of the brain. Patient history included hypothyroidism. Family history included myocardial infarction and breast cancer.
53	SPINFET02	This library was constructed using RNA isolated from spleen tissue removed from a Caucasian male fetus, who died at 23 weeks' gestation.
54	SENVNOT01	This library was constructed using RNA isolated from seminal vesicle tissue removed from a 58-year-old Caucasian male during radical prostatectomy. Pathology for the associated tumor tissue indicated adenocarcinoma (Gleason grade 3+2) of the prostate. Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA). Family history included a malignant breast neoplasm.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
55	CONUTUT01	This library was constructed using RNA isolated from sigmoid mesentery tumor tissue obtained from a 61-year-old female during a total abdominal hysterectomy and bilateral salpingo-oophorectomy with regional lymph node excision. Pathology indicated a metastatic grade 4 malignant mixed müllerian tumor present in the sigmoid mesentery at two sites.
56	HEARFET02	This library was constructed using RNA isolated from heart tissue removed from a Caucasian male fetus, who was stillborn at 23 weeks' gestation with a hypoplastic left heart.
57	BRAIFET01	This library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus, who was stillborn at 23 weeks' gestation with a hypoplastic left heart.
58	BRAINOT23	This library was constructed using RNA isolated from right temporal lobe tissue removed from a 45-year-old Black male during a brain lobectomy. Pathology for the associated tumor tissue indicated dysembryoplastic neuroepithelial tumor of the right temporal lobe. The right temporal region dura was consistent with calcifying pseudotumor of the neuraxis. The patient presented with convulsive intractable epilepsy, partial epilepsy, and memory disturbance. Patient history included obesity, meningitis, backache, unspecified sleep apnea, acute stress reaction, acquired knee deformity, and chronic sinusitis. Family history included obesity, benign hypertension, cirrhosis of the liver, alcohol abuse, hyperlipidemia, cerebrovascular disease, and type II diabetes.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.0E-6 Assembled ESTs: fasta Identity=95% or greater and Match length=200 bases or greater; fasta E value=1.0E-8 or less Full Length sequences: fasta score=100 or greater
BLIMPS	A Blocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Somnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Normalized quality scores>GCC-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <i>supra</i> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. A substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof.
2. A substantially purified variant having at least 90% amino acid sequence identity to the amino acid sequence of claim 1.
3. An isolated and purified polynucleotide encoding the polypeptide of claim 1.
4. An isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide of claim 3.
5. An isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide of claim 3.
6. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 3.
7. A method for detecting a polynucleotide, the method comprising the steps of:
 - (a) hybridizing the polynucleotide of claim 6 to at least one nucleic acid in a sample, thereby forming a hybridization complex; and
 - (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of the polynucleotide in the sample.
8. The method of claim 7 further comprising amplifying the polynucleotide prior to hybridization.
9. An isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:30-58 and fragments thereof.
10. An isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide of claim 9.

11. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 9.
12. An expression vector comprising at least a fragment of the polynucleotide of claim 3.
13. A host cell comprising the expression vector of claim 12.
14. A method for producing a polypeptide, the method comprising the steps of:
- a) culturing the host cell of claim 13 under conditions suitable for the expression of the polypeptide; and
 - b) recovering the polypeptide from the host cell culture.
15. A pharmaceutical composition comprising the polypeptide of claim 1 in conjunction with a suitable pharmaceutical carrier.
16. A purified antibody which specifically binds to the polypeptide of claim 1.
17. A purified agonist of the polypeptide of claim 1.
18. A purified antagonist of the polypeptide of claim 1.
19. A method for treating or preventing a disorder associated with decreased expression or activity of GTPAP, the method comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 15.
20. A method for treating or preventing a disorder associated with increased expression or activity of GTPAP, the method comprising administering to a subject in need of such treatment an effective amount of the antagonist of claim 18.

**DECLARATION AND POWER OF ATTORNEY FOR
UNITED STATES PATENT APPLICATION**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,
and

I believe that I am the original, first and sole inventor (if only one name is listed below)
or an original, first and joint inventor (if more than one name is listed below) of the subject
matter which is claimed and for which a United States patent is sought on the invention entitled

GTPASE ASSOCIATED PROTEINS

the specification of which:

 / is attached hereto.

 / was filed on _____ as application Serial No. _____ and if this box
contains an X /, was amended on _____.

 X / was filed as Patent Cooperation Treaty international application No. PCT/US99/28013
on 23 November, 1999, if this box contains an X /, was amended on under Patent Cooperation
Treaty Article 19 on _____ 2001, and if this box contains an X /, was amended on _____
_____.

I hereby state that I have reviewed and understand the contents of the above-identified
specification, including the claims, as amended by any amendment referred to above.

I acknowledge my duty to disclose information which is material to the examination of
this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim the benefit under Title 35, United States Code, §119 or §365(a)-(b) of any
foreign application(s) for patent or inventor's certificate indicated below and of any Patent
Cooperation Treaty international applications(s) designating at least one country other than the
United States indicated below and have also identified below any foreign application(s) for
patent or inventor's certificate and Patent Cooperation Treaty international application(s)
designating at least one country other than the United States for the same subject matter and
having a filing date before that of the application for said subject matter the priority of which is
claimed:

Country	Number	Filing Date	Priority Claimed
_____	_____	_____	// Yes // No
_____	_____	_____	// Yes // No

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

Application Serial No.	Filed	Status (Pending, Abandoned, Patented)
60/118,610	February 4, 1999	Expired
60/127,990	April 6, 1999	Expired
60/109,592	November 23, 1998	Expired

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in said prior application(s) in the manner required by the first paragraph of Title 35, United States Code §112, I acknowledge my duty to disclose material information as defined in Title 37 Code of Federal Regulations, §1.56(a) which occurred between the filing date(s) of the prior application(s) and the national or Patent Cooperation Treaty international filing date of this application:

Application Serial No.	Filed	Status (Pending, Abandoned, Patented)
_____	_____	_____

I hereby appoint the following:

Lucy J. Billings	Reg. No. <u>36,749</u>
Michael C. Cerrone	Reg. No. <u>39,132</u>
Diana Hamlet-Cox	Reg. No. <u>33,302</u>
Richard C. Ekstrom	Reg. No. <u>37,027</u>
Barrie D. Greene	Reg. No. <u>46,740</u>
Matthew R. Kaser	Reg. No. <u>44,817</u>
Lynn E. Murry	Reg. No. <u>42,918</u>
Shirley A. Recipon	Reg. No. <u>47,016</u>
Susan K. Sather	Reg. No. <u>44,316</u>
Michelle M. Stempien	Reg. No. <u>41,327</u>
David G. Streeter	Reg. No. <u>43,168</u>
Stephen Todd	Reg. No. <u>47,139</u>
Christopher Turner	Reg. No. <u>45,167</u>
P. Ben Wang	Reg. No. <u>41,420</u>

respectively and individually, as my patent attorneys and/or agents, with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent

and Trademark Office connected therewith. Please address all communications to:

LEGAL DEPARTMENT
INCYTE GENOMICS, INC.
3160 PORTER DRIVE, PALO ALTO, CA 94304

TEL: 650-855-0555 FAX: 650-849-8886 or 650-845-4166

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

**Sole Inventor or
First Joint Inventor:**

Full name:

JENNIFER L. HILLMAN

Signature:

Jennifer L. Hillman

Date:

February 16, 2001

Citizenship

United States *CA*

Residence:

Mountain View, California 94040

P.O. Address:

230 Monroe Drive, #17
Mountain View, California 94040

Second Joint Inventor:

Full name:

Y. TOM TANG

Signature:

Y. Tom Tang

Date:

February 27, 2001

Citizenship

People's Republic of China

Residence:

San Jose, California 95118 *CA*

P.O. Address:

4230 Ranwick Court
San Jose, California 95118

Third Joint Inventor:

Full name:

OLGA BANDMAN

Signature:

Olga Bandman

Date:

16 February, 2001

Citizenship

United States *CH*

Residence:

Mountain View, California 94043

P.O. Address:

366 Anna Avenue
Mountain View, California 94043

Fourth Joint Inventor:

Full name:

PREETI LAL

Signature:

Preeti Lal

Date:

FEBRUARY, 16, 2001

Citizenship

India *CH*

Residence:

Santa Clara, California 95056

P.O. Address:

P.O. Box 5142
Santa Clara California 95056

Fifth Joint Inventor:

Full name:

HENRY YUE

Signature:

Henry Yue

Date:

March 02, 2001 *CH*

Citizenship

United States

Residence:

Sunnyvale, California 94087

P.O. Address:

826 Lois Avenue
Sunnyvale, California 94087

09556679-052201

Sixth Joint Inventor:

Full name:

DYUNG AINA M. LU

Signature:

[Signature]

Date:

March 22, 2001

Citizenship

United States

Residence:

San Jose, California 95123

P.O. Address:

233 Coy Drive
San Jose, California 95123

Seventh Joint Inventor:

Full name:

MARIAH R. BAUGHN

Signature:

[Signature]

Date:

February 12, 2001

Citizenship

United States

Residence:

San Leandro, California 94577

P.O. Address:

14244 Santiago Road
San Leandro California 94577

Eighth Joint Inventor:

Full name:

JUNMING YANG

Signature:

[Signature]

Date:

February 27, 2001

Citizenship

China

Residence:

San Jose, California 95129

P.O. Address:

7125 Bark Lane
San Jose, California 95129

09856679.052201

Ninth Joint Inventor:

9W

Full name:

YALDA AZIMZAI

Signature:

Yalda Azimzai

Date:

February 02, 2001

Citizenship

United States CA

Residence:

Castro Valley, California 94552

P.O. Address:

5518 Boulder Canyon Drive
Castro Valley, California 94552

09856679-052201

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.

HILLMAN, Jennifer L.

TANG, Y. Tom

BANDMAN, Olga

LAL, Preeti

YUE, Henry

LU, Dyung Aina M.

BAUGHN, Mariah R.

YANG, Junming

AZIMZAI, Yalda

<120> GTPASE ASSOCIATED PROTEINS

<130> PF-0629 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/109,592; 60/118,610; 60/127,990

<151> 1998-11-23; 1999-02-04; 1999-04-06

<160> 58

<170> PERL Program

<210> 1

<211> 1002

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 708398CD1

<400> 1

```

Met Pro Ser Lys Phe Ser Cys Arg Gln Leu Arg Glu Ala Gly Gln
  1          5          10          15
Cys Phe Glu Ser Phe Leu Val Val Arg Gly Leu Asp Met Glu Thr
  20          25          30
Asp Arg Glu Arg Leu Arg Thr Ile Tyr Asn Arg Asp Phe Lys Ile
  35          40          45
Ser Phe Gly Thr Pro Ala Pro Gly Phe Ser Ser Met Leu Tyr Gly
  50          55          60
Met Lys Ile Ala Asn Leu Ala Tyr Val Thr Lys Thr Arg Val Arg
  65          70          75
Phe Phe Arg Leu Asp Arg Trp Ala Asp Val Arg Phe Pro Glu Lys
  80          85          90
Arg Arg Met Lys Leu Gly Ser Asp Ile Ser Lys His His Lys Ser
  95          100         105
Leu Leu Ala Lys Ile Phe Tyr Asp Arg Ala Glu Tyr Leu His Gly
  110         115         120
Lys His Gly Val Asp Val Glu Val Gln Gly Pro His Glu Ala Arg

```

	125		130		135
Asp Gly Gln Leu	Leu Ile Arg Leu	Asp Leu Asn Arg Lys	Glu Val		
	140		145		150
Leu Thr Leu Arg	Leu Arg Asn Gly Gly	Thr Gln Ser Val	Thr Leu		
	155		160		165
Thr His Leu Phe	Pro Leu Cys Arg Thr	Pro Gln Phe Ala	Phe Tyr		
	170		175		180
Asn Glu Asp Gln	Glu Leu Pro Cys Pro	Leu Gly Pro Gly	Glu Cys		
	185		190		195
Tyr Glu Leu His	Val His Cys Lys Thr	Ser Phe Val Gly	Tyr Phe		
	200		205		210
Pro Ala Thr Val	Leu Trp Glu Leu Leu	Gly Pro Gly Glu	Ser Gly		
	215		220		225
Ser Glu Gly Ala	Gly Thr Phe Tyr Ile	Ala Arg Phe Leu	Ala Ala		
	230		235		240
Val Ala His Ser	Pro Leu Ala Ala Gln	Leu Lys Pro Met	Thr Pro		
	245		250		255
Phe Lys Arg Thr	Arg Ile Thr Gly Asn	Pro Val Val Thr	Asn Arg		
	260		265		270
Ile Glu Glu Gly	Glu Arg Pro Asp Arg	Ala Lys Gly Tyr	Asp Leu		
	275		280		285
Glu Leu Ser Met	Ala Leu Gly Thr Tyr	Tyr Pro Pro Pro	Arg Leu		
	290		295		300
Arg Gln Leu Leu	Pro Met Leu Leu Gln	Gly Thr Ser Ile	Phe Thr		
	305		310		315
Ala Pro Lys Glu	Ile Ala Glu Ile Lys	Ala Gln Leu Glu	Thr Ala		
	320		325		330
Leu Lys Trp Arg	Asn Tyr Glu Val Lys	Leu Arg Leu Leu	Leu His		
	335		340		345
Leu Glu Glu Leu	Gln Met Glu His Asp	Ile Arg His Tyr	Asp Leu		
	350		355		360
Glu Ser Val Pro	Met Thr Trp Asp Pro	Val Asp Gln Asn	Pro Arg		
	365		370		375
Leu Leu Thr Leu	Glu Val Pro Gly Val	Thr Glu Ser Arg	Pro Ser		
	380		385		390
Val Leu Arg Gly	Asp His Leu Phe Ala	Leu Leu Ser Ser	Glu Thr		
	395		400		405
His Gln Glu Asp	Pro Ile Thr Tyr Lys	Gly Phe Val His	Lys Val		
	410		415		420
Glu Leu Asp Arg	Val Lys Leu Ser Phe	Ser Met Ser Leu	Leu Ser		
	425		430		435
Arg Phe Val Asp	Gly Leu Thr Phe Lys	Val Asn Phe Thr	Phe Asn		
	440		445		450
Arg Gln Pro Leu	Arg Val Gln His Arg	Ala Leu Glu Leu	Thr Gly		
	455		460		465
Arg Trp Leu Leu	Trp Pro Met Leu Phe	Pro Val Ala Pro	Arg Asp		
	470		475		480
Val Pro Leu Leu	Pro Ser Asp Val Lys	Leu Lys Leu Tyr	Asp Arg		
	485		490		495
Ser Leu Glu Ser	Asn Pro Glu Gln Leu	Gln Ala Met Arg	His Ile		
	500		505		510
Val Thr Gly Thr	Thr Arg Pro Ala Pro	Tyr Ile Ile Phe	Gly Pro		
	515		520		525
Pro Gly Thr Gly	Lys Thr Val Thr Leu	Val Glu Ala Ile	Lys Gln		
	530		535		540

Val Val Lys His Leu Pro Lys Ala His Ile Leu Ala Cys Ala Pro
 545 550 555
 Ser Asn Ser Gly Ala Asp Leu Leu Cys Gln Arg Leu Arg Val His
 560 565 570
 Leu Pro Ser Ser Ile Tyr Arg Leu Leu Ala Pro Ser Arg Asp Ile
 575 580 585
 Arg Met Val Pro Glu Asp Ile Lys Pro Cys Cys Asn Trp Asp Ala
 590 595 600
 Lys Lys Gly Glu Tyr Val Phe Pro Ala Lys Lys Lys Leu Gln Glu
 605 610 615
 Tyr Arg Val Leu Ile Thr Thr Leu Ile Thr Ala Gly Arg Leu Val
 620 625 630
 Ser Ala Gln Phe Pro Ile Asp His Phe Thr His Ile Phe Ile Asp
 635 640 645
 Glu Ala Gly His Cys Met Glu Pro Glu Ser Leu Val Ala Ile Ala
 650 655 660
 Gly Leu Met Glu Val Lys Glu Thr Gly Asp Pro Gly Gly Gln Leu
 665 670 675
 Val Leu Ala Gly Asp Pro Arg Gln Leu Gly Pro Val Leu Arg Ser
 680 685 690
 Pro Leu Thr Gln Lys His Gly Leu Gly Tyr Ser Leu Leu Glu Arg
 695 700 705
 Leu Leu Ile Tyr Asn Ser Leu Tyr Lys Lys Gly Pro Asp Gly Tyr
 710 715 720
 Asp Pro Gln Phe Ile Thr Lys Leu Leu Arg Asn Tyr Arg Ser His
 725 730 735
 Pro Thr Ile Leu Asp Ile Pro Asn Gln Leu Tyr Tyr Glu Gly Glu
 740 745 750
 Leu Gln Ala Cys Ala Asp Val Val Asp Arg Glu Arg Phe Cys Arg
 755 760 765
 Trp Ala Gly Leu Pro Arg Gln Gly Phe Pro Ile Ile Phe His Gly
 770 775 780
 Val Met Gly Lys Asp Glu Arg Glu Gly Asn Ser Pro Ser Phe Phe
 785 790 795
 Asn Pro Glu Glu Ala Ala Thr Val Thr Ser Tyr Leu Lys Leu Leu
 800 805 810
 Leu Ala Pro Ser Ser Lys Lys Gly Lys Ala Arg Leu Ser Pro Arg
 815 820 825
 Ser Val Gly Val Ile Ser Pro Tyr Arg Lys Gln Val Glu Lys Ile
 830 835 840
 Arg Tyr Cys Ile Thr Lys Leu Asp Arg Glu Leu Arg Gly Leu Asp
 845 850 855
 Asp Ile Lys Asp Leu Lys Val Gly Ser Val Glu Glu Phe Gln Gly
 860 865 870
 Gln Glu Arg Ser Val Ile Leu Ile Ser Thr Val Arg Ser Ser Gln
 875 880 885
 Ser Phe Val Gln Leu Asp Leu Asp Phe Asn Leu Gly Phe Leu Lys
 890 895 900
 Asn Pro Lys Arg Phe Asn Val Ala Val Thr Arg Ala Lys Ala Leu
 905 910 915
 Leu Ile Ile Val Gly Asn Pro Leu Leu Leu Gly His Asp Pro Asp
 920 925 930
 Trp Lys Val Phe Leu Glu Phe Cys Lys Glu Asn Gly Gly Tyr Thr
 935 940 945
 Gly Cys Pro Phe Pro Ala Lys Leu Asp Leu Gln Gln Gly Gln Asn

	950		955		960
Leu Leu Gln Gly	Leu Ser Lys Leu Ser	Pro Ser Thr Ser Gly	Pro		
	965		970		975
His Ser His Asp	Tyr Leu Pro Gln Glu	Arg Glu Gly Glu Gly	Gly		
	980		985		990
Leu Ser Leu Gln	Val Glu Pro Glu Trp	Arg Asn Glu			
	995		1000		

<210> 2
 <211> 338
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1259937CD1

<400> 2
 Met Ala Ala Leu Ala Gln Glu Asp Gly Trp Thr Lys Gly Gln Val
 1 5 10 15
 Leu Val Lys Val Asn Ser Ala Gly Asp Ala Ile Gly Leu Gln Pro
 20 25 30
 Asp Ala Arg Gly Val Ala Thr Ser Leu Gly Leu Asn Glu Arg Leu
 35 40 45
 Phe Val Val Asn Pro Gln Glu Val His Glu Leu Ile Pro His Pro
 50 55 60
 Asp Gln Leu Gly Pro Thr Val Gly Ser Ala Glu Gly Leu Asp Leu
 65 70 75
 Val Ser Ala Lys Asp Leu Ala Gly Gln Leu Thr Asp His Asp Trp
 80 85 90
 Ser Leu Phe Asn Ser Ile His Gln Val Glu Leu Ile His Tyr Val
 95 100 105
 Leu Gly Pro Gln His Leu Arg Asp Val Thr Thr Ala Asn Leu Glu
 110 115 120
 Arg Phe Met Arg Arg Phe Asn Glu Leu Gln Tyr Trp Val Ala Thr
 125 130 135
 Glu Leu Cys Leu Cys Pro Val Pro Gly Pro Arg Ala Gln Leu Leu
 140 145 150
 Arg Lys Phe Ile Lys Leu Ala Ala His Leu Lys Glu Gln Lys Asn
 155 160 165
 Leu Asn Ser Phe Phe Ala Val Met Phe Gly Leu Ser Asn Ser Ala
 170 175 180
 Ile Ser Arg Leu Ala His Thr Trp Glu Arg Leu Pro His Lys Val
 185 190 195
 Arg Lys Leu Tyr Ser Ala Leu Glu Arg Leu Leu Asp Pro Ser Trp
 200 205 210
 Asn His Arg Val Tyr Arg Leu Ala Leu Ala Lys Leu Ser Pro Pro
 215 220 225
 Val Ile Pro Phe Met Pro Leu Leu Leu Lys Asp Met Thr Phe Ile
 230 235 240
 His Glu Gly Asn His Thr Leu Val Glu Asn Leu Ile Asn Phe Glu
 245 250 255
 Lys Met Arg Met Met Ala Arg Ala Ala Arg Met Leu His His Cys
 260 265 270

Arg Ser His Asn Pro Val Pro Leu Ser Pro Leu Arg Ser Arg Val
 275 285
 Ser His Leu His Glu Asp Ser Gln Val Ala Arg Ile Ser Thr Cys
 290 295 300
 Ser Glu Gln Ser Leu Ser Thr Arg Ser Pro Ala Ser Thr Trp Ala
 305 310 315
 Tyr Val Gln Gln Leu Lys Val Ile Asp Asn Gln Arg Glu Leu Ser
 320 325 330
 Arg Leu Ser Arg Glu Leu Glu Pro
 335

<210> 3

<211> 211

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1452285CD1

<400> 3

Met Gln Ala Pro His Lys Glu His Leu Tyr Lys Leu Leu Val Ile
 1 5 10 15
 Gly Asp Leu Gly Val Gly Lys Thr Ser Ile Ile Lys Arg Tyr Val
 20 25 30
 His Gln Asn Phe Ser Ser His Tyr Arg Ala Thr Ile Gly Val Asp
 35 40 45
 Phe Ala Leu Lys Val Leu His Trp Asp Pro Glu Thr Val Val Arg
 50 55 60
 Leu Gln Leu Trp Asp Ile Ala Gly Gln Glu Arg Phe Gly Asn Met
 65 70 75
 Thr Arg Val Tyr Tyr Arg Glu Ala Met Gly Ala Phe Ile Val Phe
 80 85 90
 Asp Val Thr Arg Pro Ala Thr Phe Glu Ala Val Ala Lys Trp Lys
 95 100 105
 Asn Asp Leu Asp Ser Lys Leu Ser Leu Pro Asn Gly Lys Pro Val
 110 115 120
 Ser Val Val Leu Leu Ala Asn Lys Cys Asp Gln Gly Lys Asp Val
 125 130 135
 Leu Met Asn Asn Gly Leu Lys Met Asp Gln Phe Cys Lys Glu His
 140 145 150
 Gly Phe Val Gly Trp Phe Glu Thr Ser Ala Lys Glu Asn Ile Asn
 155 160 165
 Ile Asp Glu Ala Ser Arg Cys Leu Val Lys His Ile Leu Ala Asn
 170 175 180
 Glu Cys Asp Leu Met Glu Ser Ile Glu Pro Asp Val Val Lys Pro
 185 190 195
 His Leu Thr Ser Thr Lys Val Ala Ser Cys Ser Gly Cys Ala Lys
 200 205 210
 Ser

<210> 4

<211> 516

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1812894CD1

<400> 4

```

Met Glu Thr Met Lys Ala Val Ala Glu Val Ser Glu Ser Thr Lys
 1          5          10          15
Ala Glu Ala Val Ala Ala Val Gln Arg Gln Cys Gln Glu Glu Val
 20          25          30
Ala Ser Leu Gln Ala Ile Leu Lys Asp Ser Ile Ser Ser Tyr Glu
 35          40          45
Ala Gln Ile Thr Ala Leu Lys Gln Glu Arg Gln Gln Gln Gln Gln
 50          55          60
Asp Cys Glu Glu Lys Glu Arg Glu Leu Gly Arg Leu Lys Gln Leu
 65          70          75
Leu Ser Arg Ala Tyr Pro Leu Asp Ser Leu Glu Lys Gln Met Glu
 80          85          90
Lys Ala His Glu Asp Ser Glu Lys Leu Arg Glu Ile Val Leu Pro
 95          100          105
Met Glu Lys Glu Ile Glu Glu Leu Lys Ala Lys Leu Leu Arg Ala
110          115          120
Glu Glu Leu Ile Gln Glu Ile Gln Arg Arg Pro Arg His Ala Pro
125          130          135
Ser Leu His Gly Ser Thr Glu Leu Leu Pro Leu Ser Arg Asp Pro
140          145          150
Ser Pro Pro Leu Glu Pro Leu Glu Glu Leu Ser Gly Asp Gly Gly
155          160          165
Pro Ala Ala Glu Ala Phe Ala His Asn Cys Asp Asp Ser Ala Ser
170          175          180
Ile Ser Ser Phe Ser Leu Gly Gly Gly Val Gly Ser Ser Ser Ser
185          190          195
Leu Pro Gln Ser Arg Gln Gly Leu Ser Pro Glu Gln Glu Glu Thr
200          205          210
Ala Ser Leu Val Ser Thr Gly Thr Leu Val Pro Glu Gly Ile Tyr
215          220          225
Leu Pro Pro Pro Gly Tyr Gln Leu Val Pro Asp Thr Gln Trp Glu
230          235          240
Gln Leu Gln Thr Glu Gly Arg Gln Leu Gln Lys Asp Leu Glu Ser
245          250          255
Val Ser Arg Glu Arg Asp Glu Leu Gln Glu Gly Leu Arg Arg Ser
260          265          270
Asn Glu Asp Cys Ala Lys Gln Met Gln Val Leu Leu Ala Gln Val
275          280          285
Gln Asn Ser Glu Gln Leu Leu Arg Thr Leu Gln Gly Thr Val Ser
290          295          300
Gln Ala Gln Glu Arg Val Gln Leu Gln Met Ala Glu Leu Val Thr
305          310          315
Thr His Lys Cys Leu His His Glu Val Lys Arg Leu Asn Glu Glu
320          325          330
Asn Gln Gly Leu Arg Ala Glu Gln Leu Pro Ser Ser Ala Pro Gln
335          340          345
Gly Ser Gln Gln Glu Gln Gly Glu Glu Glu Ser Leu Pro Ser Ser

```

350 355 360
 Val Pro Glu Leu Gln Gln Leu Leu Cys Cys Thr Arg Gln Glu Ala
 365 370 375
 Arg Ala Arg Leu Gln Ala Gln Glu His Gly Ala Glu Arg Leu Arg
 380 385 390
 Ile Glu Ile Val Thr Leu Arg Glu Ala Leu Glu Glu Glu Thr Val
 395 400 405
 Ala Arg Ala Ser Leu Glu Gly Gln Leu Arg Val Gln Arg Glu Glu
 410 415 420
 Thr Glu Val Leu Glu Ala Ser Leu Cys Ser Leu Arg Thr Glu Met
 425 430 435
 Glu Arg Val Gln Gln Glu Gln Ser Lys Ala Gln Leu Pro Asp Leu
 440 445 450
 Leu Ser Glu Gln Arg Ala Lys Val Leu Arg Leu Gln Ala Glu Leu
 455 460 465
 Glu Thr Ser Glu Gln Val Gln Arg Asp Phe Val Arg Leu Ser Gln
 470 475 480
 Ala Leu Gln Val Arg Leu Glu Arg Ile Arg Gln Ala Glu Thr Leu
 485 490 495
 Glu Gln Val Arg Ser Ile Met Asp Glu Ala Pro Leu Thr Asp Val
 500 505 510
 Arg Asp Ile Lys Asp Thr
 515

<210> 5

<211> 445

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3074884CD1

<400> 5

Met Pro Glu Asp Ala Asp Glu Asn Ala Glu Glu Glu Leu Leu Arg
 1 5 10 15
 Gly Glu Pro Leu Leu Pro Ala Gly Thr Gln Arg Val Cys Leu Val
 20 25 30
 His Pro Asp Val Lys Trp Gly Pro Gly Lys Ser Gln Met Thr Arg
 35 40 45
 Ala Glu Trp Gln Val Ala Glu Ala Thr Ala Leu Val His Thr Leu
 50 55 60
 Asp Gly Trp Ser Val Val Gln Thr Met Val Val Ser Thr Lys Thr
 65 70 75
 Pro Asp Arg Lys Leu Ile Phe Gly Lys Gly Asn Phe Glu His Leu
 80 85 90
 Thr Glu Lys Ile Arg Gly Ser Pro Asp Val Thr Cys Val Phe Leu
 95 100 105
 Asn Val Glu Arg Met Ala Ala Pro Thr Lys Lys Glu Leu Glu Ala
 110 115 120
 Ala Trp Gly Val Glu Val Phe Asp Arg Phe Thr Val Val Leu His
 125 130 135
 Ile Phe Arg Cys Asn Ala Arg Thr Lys Glu Ala Arg Leu Gln Val
 140 145 150

Ala Leu Ala Glu Met Pro Leu His Arg Ser Asn Leu Lys Arg Asp
 155 160 165
 Val Ala His Leu Tyr Arg Gly Val Gly Ser Arg Tyr Ile Met Gly
 170 175 180
 Ser Gly Glu Ser Phe Met Gln Leu Gln Gln Arg Leu Leu Arg Glu
 185 190 195
 Lys Glu Ala Lys Ile Arg Lys Ala Leu Asp Arg Leu Arg Lys Lys
 200 205 210
 Arg His Leu Leu Arg Arg Gln Arg Thr Arg Arg Glu Phe Pro Val
 215 220 225
 Ile Ser Val Val Gly Tyr Thr Asn Cys Gly Lys Thr Thr Leu Ile
 230 235 240
 Lys Ala Leu Thr Gly Asp Ala Ala Ile Gln Pro Arg Asp Gln Leu
 245 250 255
 Phe Ala Thr Leu Asp Val Thr Ala His Ala Gly Thr Leu Pro Ser
 260 265 270
 Arg Met Thr Val Leu Tyr Val Asp Thr Ile Gly Phe Leu Ser Gln
 275 280 285
 Leu Pro His Gly Leu Ile Glu Ser Phe Ser Ala Thr Leu Glu Asp
 290 295 300
 Val Ala His Ser Asp Leu Ile Leu His Val Arg Asp Val Ser His
 305 310 315
 Pro Glu Ala Glu Leu Gln Lys Cys Ser Val Leu Ser Thr Leu Arg
 320 325 330
 Gly Leu Gln Leu Pro Ala Pro Leu Leu Asp Ser Met Val Glu Val
 335 340 345
 His Asn Lys Val Asp Leu Val Pro Gly Tyr Ser Pro Thr Glu Pro
 350 355 360
 Asn Val Val Pro Val Ser Ala Leu Arg Gly His Gly Leu Gln Glu
 365 370 375
 Leu Lys Ala Glu Leu Asp Ala Ala Val Leu Lys Ala Thr Gly Arg
 380 385 390
 Gln Ile Leu Thr Leu Arg Val Arg Leu Ala Gly Ala Gln Leu Ser
 395 400 405
 Trp Leu Tyr Lys Glu Ala Thr Val Gln Glu Val Asp Val Ile Pro
 410 415 420
 Glu Asp Gly Ala Ala Asp Val Arg Val Ile Ile Ser Asn Ser Ala
 425 430 435
 Tyr Gly Lys Phe Arg Lys Leu Phe Pro Gly
 440 445

<210> 6

<211> 445

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3452277CD1

<400> 6

Met Tyr Tyr Gln Gln Ala Leu Met Arg Ser Thr Val Lys Ser Ser
 1 5 10 15
 Val Ser Leu Gly Gly Ile Val Lys Tyr Ser Glu Gln Phe Ser Ser

	20		25		30
Asn Asp Ala Ile Met Ser Gly Cys Leu Pro Ser Asn Pro Trp Ile					
	35		40		45
Thr Asp Asp Thr Gln Phe Trp Asp Leu Asn Ala Lys Leu Val Glu					
	50		55		60
Ile Pro Thr Lys Met Arg Val Glu Arg Trp Ala Phe Asn Phe Ser					
	65		70		75
Glu Leu Ile Arg Asp Pro Lys Gly Arg Gln Ser Phe Gln Tyr Phe					
	80		85		90
Leu Lys Lys Glu Phe Ser Gly Glu Asn Leu Gly Phe Trp Glu Ala					
	95		100		105
Cys Glu Asp Leu Lys Tyr Gly Asp Gln Ser Lys Val Lys Glu Lys					
	110		115		120
Ala Glu Glu Ile Tyr Lys Leu Phe Leu Ala Pro Gly Ala Arg Arg					
	125		130		135
Trp Ile Asn Ile Asp Gly Lys Thr Met Asp Ile Thr Val Lys Gly					
	140		145		150
Leu Lys His Pro His Arg Tyr Val Leu Asp Ala Ala Gln Thr His					
	155		160		165
Ile Tyr Met Leu Met Lys Lys Asp Ser Tyr Ala Arg Tyr Leu Lys					
	170		175		180
Ser Pro Ile Tyr Lys Asp Met Leu Ala Lys Ala Ile Glu Pro Gln					
	185		190		195
Glu Thr Thr Lys Lys Ser Ser Thr Leu Pro Phe Met Arg Arg His					
	200		205		210
Leu Arg Ser Ser Pro Ser Pro Val Ile Leu Arg Gln Leu Glu Glu					
	215		220		225
Glu Ala Lys Ala Arg Glu Ala Ala Asn Thr Val Asp Ile Thr Gln					
	230		235		240
Pro Gly Gln His Met Ala Pro Ser Pro His Leu Thr Val Tyr Thr					
	245		250		255
Gly Thr Cys Met Pro Pro Ser Pro Ser Ser Pro Phe Ser Ser Ser					
	260		265		270
Cys Arg Ser Pro Arg Lys Pro Phe Ala Ser Pro Ser Arg Phe Ile					
	275		280		285
Arg Arg Pro Ser Thr Thr Ile Cys Pro Ser Pro Ile Arg Val Ala					
	290		295		300
Leu Glu Ser Ser Ser Gly Leu Glu Gln Lys Gly Glu Cys Ser Gly					
	305		310		315
Ser Met Ala Pro Arg Gly Pro Ser Val Thr Glu Ser Ser Glu Ala					
	320		325		330
Ser Leu Asp Thr Ser Trp Pro Arg Ser Arg Pro Arg Ala Pro Pro					
	335		340		345
Lys Ala Arg Met Ala Leu Ser Phe Ser Arg Phe Leu Arg Arg Gly					
	350		355		360
Cys Leu Ala Ser Pro Val Phe Ala Arg Leu Ser Pro Lys Cys Pro					
	365		370		375
Ala Val Ser His Gly Arg Val Gln Pro Leu Gly Asp Val Gly Gln					
	380		385		390
Gln Leu Pro Arg Leu Lys Ser Lys Arg Val Ala Asn Phe Phe Gln					
	395		400		405
Ile Lys Met Asp Val Pro Thr Gly Ser Gly Thr Cys Leu Met Asp					
	410		415		420
Ser Glu Asp Ala Gly Thr Gly Glu Ser Gly Asp Arg Ala Thr Glu					
	425		430		435

Lys Glu Val Ile Cys Pro Trp Glu Ser Leu
440 445

<210> 7
<211> 281
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 4203832CD1

<400> 7
Met Lys Leu Ala Ala Met Ile Lys Lys Met Cys Pro Ser Asp Ser
1 5 10 15
Glu Leu Ser Ile Pro Ala Lys Asn Cys Tyr Arg Met Val Ile Leu
20 25 30
Gly Ser Ser Lys Val Gly Lys Thr Ala Ile Val Ser Arg Phe Leu
35 40 45
Thr Gly Arg Phe Glu Asp Ala Tyr Thr Pro Thr Ile Glu Asp Phe
50 55 60
His Arg Lys Phe Tyr Ser Ile Arg Gly Glu Val Tyr Gln Leu Asp
65 70 75
Ile Leu Asp Thr Ser Gly Asn His Pro Phe Pro Ala Met Arg Cys
80 85 90
Leu Ser Ile Leu Thr Gly Asp Val Phe Ile Leu Val Phe Ser Leu
95 100 105
Asp Asn Arg Asp Ser Phe Glu Glu Val Gln Arg Leu Arg Gln Gln
110 115 120
Ile Leu Asp Thr Lys Ser Cys Leu Lys Asn Lys Thr Lys Glu Asn
125 130 135
Val Asp Val Pro Leu Val Ile Cys Gly Asn Lys Gly Asp Arg Asp
140 145 150
Phe Tyr Arg Glu Val Asp Gln Arg Glu Ile Glu Gln Leu Val Gly
155 160 165
Asp Asp Pro Gln Arg Cys Ala Tyr Phe Glu Ile Ser Ala Lys Lys
170 175 180
Asn Ser Ser Leu Asp Gln Met Phe Arg Ala Leu Phe Ala Met Ala
185 190 195
Lys Leu Pro Ser Glu Met Ser Pro Asp Leu His Arg Lys Val Ser
200 205 210
Val Gln Tyr Cys Asp Val Leu His Lys Lys Ala Leu Arg Asn Lys
215 220 225
Lys Leu Leu Arg Ala Gly Ser Gly Gly Gly Gly Gly Asp Pro Gly
230 235 240
Asp Ala Phe Gly Ile Val Ala Pro Phe Ala Arg Arg Pro Ser Val
245 250 255
His Ser Asp Leu Met Tyr Ile Arg Glu Lys Ala Ser Ala Gly Ser
260 265 270
Gln Ala Lys Asp Lys Glu Arg Cys Val Ile Ser
275 280

<210> 8
<211> 301
<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 104368CD1

<400> 8

```

Met Thr Thr Leu Asp Asp Lys Leu Leu Gly Glu Lys Leu Gln Tyr
 1          5          10          15
Tyr Tyr Ser Ser Ser Glu Asp Glu Asp Ser Asp His Glu Asp Lys
 20          25          30
Asp Arg Gly Arg Cys Ala Pro Ala Ser Ser Ser Val Pro Ala Glu
 35          40          45
Ala Glu Leu Ala Gly Glu Gly Ile Ser Val Asn Thr Gly Pro Lys
 50          55          60
Gly Val Ile Asn Asp Trp Arg Arg Phe Lys Gln Leu Glu Thr Glu
 65          70          75
Gln Arg Glu Glu Gln Cys Arg Glu Met Glu Arg Leu Ile Lys Lys
 80          85          90
Leu Ser Met Thr Cys Arg Ser His Leu Asp Glu Glu Glu Glu Gln
 95          100         105
Gln Lys Gln Lys Asp Leu Gln Glu Lys Ile Ser Gly Lys Met Thr
 110         115         120
Leu Lys Glu Phe Ala Ile Met Asn Glu Asp Gln Asp Asp Glu Glu
 125         130         135
Phe Leu Gln Gln Tyr Arg Lys Gln Arg Met Glu Glu Met Arg Gln
 140         145         150
Gln Leu His Lys Gly Pro Gln Phe Lys Gln Val Phe Glu Ile Ser
 155         160         165
Ser Gly Glu Gly Phe Leu Asp Met Ile Asp Lys Glu Gln Lys Ser
 170         175         180
Ile Val Ile Met Val His Ile Tyr Glu Asp Gly Ile Pro Gly Thr
 185         190         195
Glu Ala Met Asn Gly Cys Met Ile Cys Leu Ala Ala Glu Tyr Pro
 200         205         210
Ala Val Lys Phe Cys Lys Val Lys Ser Ser Val Ile Gly Ala Ser
 215         220         225
Ser Gln Phe Thr Arg Asn Ala Leu Pro Ala Leu Leu Ile Tyr Lys
 230         235         240
Gly Gly Glu Leu Ile Gly Asn Phe Val Arg Val Thr Asp Gln Leu
 245         250         255
Gly Asp Asp Phe Phe Ala Val Asp Leu Glu Ala Phe Leu Gln Glu
 260         265         270
Phe Gly Leu Leu Pro Glu Lys Glu Val Leu Val Leu Thr Ser Val
 275         280         285
Arg Asn Ser Ala Thr Cys His Ser Glu Asp Ser Asp Leu Glu Ile
 290         295         300
Asp

```

<210> 9

<211> 485

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1441680CD1

<400> 9

```

Met Arg Ala Val Leu Thr Trp Arg Asp Lys Ala Glu His Cys Ile
 1          5          10          15
Asn Asp Ile Ala Phe Lys Pro Asp Gly Thr Gln Leu Ile Leu Ala
 20          25          30
Ala Gly Ser Arg Leu Leu Val Tyr Asp Thr Ser Asp Gly Thr Leu
 35          40          45
Leu Gln Pro Leu Lys Gly His Lys Asp Thr Val Tyr Cys Val Ala
 50          55          60
Tyr Ala Lys Asp Gly Lys Arg Phe Ala Ser Gly Ser Ala Asp Lys
 65          70          75
Ser Val Ile Ile Trp Thr Ser Lys Leu Glu Gly Ile Leu Lys Tyr
 80          85          90
Thr His Asn Asp Ala Ile Gln Cys Val Ser Tyr Asn Pro Ile Thr
 95          100          105
His Gln Leu Ala Ser Cys Ser Ser Ser Asp Phe Gly Leu Trp Ser
 110          115          120
Pro Glu Gln Lys Ser Val Ser Lys His Lys Ser Ser Ser Lys Ile
 125          130          135
Ile Cys Cys Ser Trp Thr Asn Asp Gly Gln Tyr Leu Ala Leu Gly
 140          145          150
Met Phe Asn Gly Ile Ile Ser Ile Arg Asn Lys Asn Gly Glu Glu
 155          160          165
Lys Val Lys Ile Glu Arg Pro Gly Gly Ser Leu Ser Pro Ile Trp
 170          175          180
Ser Ile Cys Trp Asn Pro Ser Arg Glu Glu Arg Asn Asp Ile Leu
 185          190          195
Ala Val Ala Asp Trp Gly Gln Lys Val Ser Phe Tyr Gln Leu Ser
 200          205          210
Gly Lys Gln Ile Gly Lys Asp Arg Ala Leu Asn Phe Asp Pro Cys
 215          220          225
Cys Ile Ser Tyr Phe Thr Lys Gly Glu Tyr Ile Leu Leu Gly Gly
 230          235          240
Ser Asp Lys Gln Val Ser Leu Phe Thr Lys Asp Gly Val Arg Leu
 245          250          255
Gly Thr Val Gly Glu Gln Asn Ser Trp Val Trp Thr Cys Gln Ala
 260          265          270
Lys Pro Asp Ser Asn Tyr Val Val Val Gly Cys Gln Asp Gly Thr
 275          280          285
Ile Ser Phe Tyr Gln Leu Ile Phe Ser Thr Val His Gly Val Tyr
 290          295          300
Lys Asp Arg Tyr Ala Tyr Arg Asp Ser Met Thr Asp Val Ile Val
 305          310          315
Gln His Leu Ile Thr Glu Gln Lys Val Arg Ile Lys Cys Lys Glu
 320          325          330
Leu Val Lys Lys Ile Ala Ile Tyr Arg Asn Arg Leu Ala Ile Gln
 335          340          345
Leu Pro Glu Lys Ile Leu Ile Tyr Glu Leu Tyr Ser Glu Asp Leu
 350          355          360

```

Ser Asp Met His Tyr Arg Val Lys Glu Lys Ile Ile Lys Lys Phe
 365 370
 Glu Cys Asn Leu Leu Val Val Cys Ala Asn His Ile Ile Leu Cys
 380 385
 Gln Glu Lys Arg Leu Gln Cys Leu Ser Phe Ser Gly Val Lys Glu
 395 400
 Arg Glu Trp Gln Met Glu Ser Leu Ile Arg Tyr Ile Lys Val Ile
 410 415
 Gly Gly Pro Pro Gly Arg Glu Gly Leu Leu Val Gly Leu Lys Lys
 425 430
 Met Tyr Leu Leu Val Tyr Ser Phe Ile Leu Ile Val Lys Asp Tyr
 440 445
 Phe Ser Leu Ser Thr Asp Val Leu Gly Asn Leu Thr Trp Lys His
 455 460
 Val Cys Lys Lys His Tyr Trp Val Phe His Leu Phe Ser Trp Tyr
 470 475
 Tyr Ile Phe Val Gln
 485

<210> 10

<211> 447

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1494955CD1

<400> 10

Met Glu Leu Ser Gln Met Ser Glu Leu Met Gly Leu Ser Val Leu
 1 5 10 15
 Leu Gly Leu Leu Ala Leu Met Ala Thr Ala Ala Val Ala Arg Gly
 20 25 30
 Trp Leu Arg Ala Gly Glu Glu Arg Ser Gly Arg Pro Ala Cys Gln
 35 40 45
 Lys Ala Asn Gly Phe Pro Pro Asp Lys Ser Ser Gly Ser Lys Lys
 50 55 60
 Gln Lys Gln Tyr Gln Arg Ile Arg Lys Glu Lys Pro Gln Gln His
 65 70 75
 Asn Phe Thr His Arg Leu Leu Ala Ala Ala Leu Lys Ser His Ser
 80 85 90
 Gly Asn Ile Ser Cys Met Asp Phe Ser Ser Asn Gly Lys Tyr Leu
 95 100 105
 Ala Thr Cys Ala Asp Asp Arg Thr Ile Arg Ile Trp Ser Thr Lys
 110 115 120
 Asp Phe Leu Gln Arg Glu His Arg Ser Met Arg Ala Asn Val Glu
 125 130 135
 Leu Asp His Ala Thr Leu Val Arg Phe Ser Pro Asp Cys Arg Ala
 140 145 150
 Phe Ile Val Trp Leu Ala Asn Gly Asp Thr Leu Arg Val Phe Lys
 155 160 165
 Met Thr Lys Arg Glu Asp Gly Gly Tyr Thr Phe Thr Ala Thr Pro
 170 175 180
 Glu Asp Phe Pro Lys Lys His Lys Ala Pro Val Ile Asp Ile Gly

	185		190		195
Ile Ala Asn Thr	Gly Lys Phe Ile Met	Thr Ala Ser Ser Asp Thr			
	200		205		210
Thr Val Leu Ile	Trp Ser Leu Lys Gly	Gln Val Leu Ser Thr Ile			
	215		220		225
Asn Thr Asn Gln	Met Asn Asn Thr His	Ala Ala Val Ser Pro Cys			
	230		235		240
Gly Arg Phe Val	Ala Ser Cys Gly Phe Thr	Pro Asp Val Lys Val			
	245		250		255
Trp Glu Val Cys	Phe Gly Lys Lys Gly	Glu Phe Gln Glu Val Val			
	260		265		270
Arg Ala Phe Glu	Leu Lys Gly His Ser	Ala Ala Val His Ser Phe			
	275		280		285
Ala Phe Ser Asn	Asp Ser Arg Arg Met	Ala Ser Val Ser Lys Asp			
	290		295		300
Gly Thr Trp Lys	Leu Trp Asp Thr Asp	Val Glu Tyr Lys Lys Lys			
	305		310		315
Gln Asp Pro Tyr	Leu Leu Lys Thr Gly	Arg Phe Glu Glu Ala Ala			
	320		325		330
Gly Ala Ala Pro	Cys Arg Leu Ala Leu	Ser Pro Asn Ala Gln Val			
	335		340		345
Leu Ala Leu Ala	Ser Gly Ser Ser Ile	His Leu Tyr Asn Thr Arg			
	350		355		360
Arg Gly Glu Lys	Glu Glu Cys Phe Glu	Arg Val His Gly Glu Cys			
	365		370		375
Ile Ala Asn Leu	Ser Phe Asp Ile Thr	Gly Arg Phe Leu Ala Ser			
	380		385		390
Cys Gly Asp Arg	Ala Val Arg Leu Phe	His Asn Thr Pro Gly His			
	395		400		405
Arg Ala Met Val	Glu Glu Met Gln Gly	His Leu Lys Arg Ala Ser			
	410		415		420
Asn Glu Ser Thr	Arg Gln Arg Leu Gln	Gln Gln Leu Thr Gln Ala			
	425		430		435
Gln Glu Thr Leu	Lys Ser Leu Gly Ala	Leu Lys Lys			
	440		445		

<210> 11

<211> 199

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1508161CD1

<400> 11

Met Pro Val Lys	Lys Lys His Arg Ala	Arg Met Ile Glu Tyr Phe
1	5	10
Ile Asp Val Ala	Arg Glu Cys Phe Asn	Ile Gly Asn Phe Asn Ser
	20	25
Leu Met Ala Ile	Ile Ser Gly Met Asn	Met Ser Pro Val Ser Arg
	35	40
Leu Lys Lys Thr	Trp Ala Lys Val Lys	Thr Ala Lys Phe Asp Ile
	50	55
		60

Leu Glu His Gln Met Asp Pro Ser Ser Asn Phe Tyr Asn Tyr Arg
 65 70
 Thr Ala Leu Arg Gly Ala Ala Gln Arg Ser Leu Thr Ala His Ser
 80 85
 Ser Arg Glu Lys Ile Val Ile Pro Phe Phe Ser Leu Leu Ile Lys
 95 100
 Asp Ile Tyr Phe Leu Asn Glu Gly Cys Ala Asn Arg Leu Pro Asn
 110 115
 Gly His Val Asn Phe Glu Lys Phe Trp Glu Leu Ala Lys Gln Val
 125 130
 Ser Glu Phe Met Thr Trp Lys Gln Val Glu Cys Pro Phe Glu Arg
 140 145
 Asp Arg Lys Ile Leu Gln Tyr Leu Leu Thr Val Pro Val Phe Ser
 155 160
 Glu Asp Ala Leu Tyr Leu Ala Ser Tyr Glu Ser Glu Gly Pro Glu
 170 175
 Asn His Ile Glu Lys Asp Arg Trp Lys Ser Leu Arg Ser Ser Leu
 185 190
 Leu Gly Arg Val 195

<210> 12

<211> 694

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1811877CD1

<400> 12

Met Ala Phe Asp Pro Thr Ser Thr Leu Leu Ala Thr Gly Gly Cys
 1 5 10
 Asp Gly Ala Val Arg Val Trp Asp Ile Val Arg His Tyr Gly Thr
 20 25 30
 His His Phe Arg Gly Ser Pro Gly Val Val His Leu Val Ala Phe
 35 40 45
 His Pro Asp Pro Thr Arg Leu Leu Leu Phe Ser Ser Ala Thr Asp
 50 55 60
 Ala Ala Ile Arg Val Trp Ser Leu Gln Asp Arg Ser Cys Leu Ala
 65 70 75
 Val Leu Thr Ala His Tyr Ser Ala Val Thr Ser Leu Ala Phe Ser
 80 85 90
 Ala Asp Gly His Thr Met Leu Ser Ser Gly Arg Asp Lys Ile Cys
 95 100 105
 Ile Ile Trp Asp Leu Gln Ser Cys Gln Ala Thr Arg Thr Val Pro
 110 115 120
 Val Phe Glu Ser Val Glu Ala Ala Val Leu Leu Pro Glu Glu Pro
 125 130 135
 Val Ser Gln Leu Gly Val Lys Ser Pro Gly Leu Tyr Phe Leu Thr
 140 145 150
 Ala Gly Asp Gln Gly Thr Leu Arg Val Trp Glu Ala Ala Ser Gly
 155 160 165
 Gln Cys Val Tyr Thr Gln Ala Gln Pro Pro Gly Pro Gly Gln Glu
 170 175 180

Leu Thr His Cys Thr Leu Ala His Thr Ala Gly Val Val Leu Thr
 185 195
 Ala Thr Ala Asp His Asn Leu Leu Leu Tyr Glu Ala Arg Ser Leu
 200 210
 Arg Leu Gln Lys Gln Phe Ala Gly Tyr Ser Glu Glu Val Leu Asp
 215 225
 Val Arg Phe Leu Gly Pro Glu Asp Ser His Val Val Val Ala Ser
 230 240
 Asn Ser Pro Cys Leu Lys Val Phe Glu Leu Gln Thr Ser Ala Cys
 245 255
 Gln Ile Leu His Gly His Thr Asp Ile Val Leu Ala Leu Asp Val
 260 270
 Phe Arg Lys Gly Trp Leu Phe Ala Ser Cys Ala Lys Asp Gln Ser
 275 285
 Val Arg Ile Trp Arg Met Asn Lys Ala Gly Gln Val Met Cys Val
 290 300
 Ala Gln Gly Ser Gly His Thr His Ser Val Gly Thr Val Cys Cys
 305 315
 Ser Arg Leu Lys Glu Ser Phe Leu Val Thr Gly Ser Gln Asp Cys
 320 330
 Thr Val Lys Leu Trp Pro Leu Pro Lys Ala Leu Leu Ser Lys Asn
 335 345
 Thr Ala Pro Asp Asn Gly Pro Ile Leu Leu Gln Ala Gln Thr Thr
 350 360
 Gln Arg Cys His Asp Lys Asp Ile Asn Ser Val Ala Ile Ala Pro
 365 375
 Asn Asp Lys Leu Leu Ala Thr Gly Ser Gln Asp Arg Thr Ala Lys
 380 390
 Leu Trp Ala Leu Pro Gln Cys Gln Leu Leu Gly Val Phe Ser Gly
 395 405
 His Arg Arg Gly Leu Trp Cys Val Gln Phe Ser Pro Met Asp Gln
 410 420
 Val Leu Ala Thr Ala Ser Ala Asp Gly Thr Ile Lys Leu Trp Ala
 425 435
 Leu Gln Asp Phe Ser Cys Leu Lys Thr Phe Glu Gly His Asp Ala
 440 450
 Ser Val Leu Lys Val Ala Phe Val Ser Arg Gly Thr Gln Leu Leu
 455 465
 Ser Ser Gly Ser Asp Gly Leu Val Lys Leu Trp Thr Ile Lys Asn
 470 480
 Asn Glu Cys Val Arg Thr Leu Asp Ala His Glu Asp Lys Val Trp
 485 495
 Gly Leu His Cys Ser Arg Leu Asp Asp His Ala Leu Thr Gly Ala
 500 510
 Ser Asp Ser Arg Val Ile Leu Trp Lys Asp Val Thr Glu Ala Glu
 515 525
 Gln Ala Glu Glu Gln Ala Arg Gln Glu Glu Gln Val Val Arg Gln
 530 540
 Gln Glu Leu Asp Asn Leu Leu His Glu Lys Arg Tyr Leu Arg Ala
 545 555
 Leu Gly Leu Ala Ile Ser Leu Asp Arg Pro His Thr Val Leu Thr
 560 570
 Val Ile Gln Ala Ile Arg Arg Asp Pro Glu Ala Cys Glu Lys Leu
 575 585
 Glu Ala Thr Met Leu Arg Leu Arg Arg Asp Gln Lys Glu Ala Leu

	590		595		600
Leu Arg Phe Cys Val Thr Trp Asn Thr Asn Ser Arg His Cys His					
	605		610		615
Glu Ala Gln Ala Val Leu Gly Val Leu Leu Arg Arg Glu Ala Pro					
	620		625		630
Glu Glu Leu Leu Ala Tyr Glu Gly Val Arg Ala Ala Leu Glu Ala					
	635		640		645
Leu Leu Pro Tyr Thr Glu Arg His Phe Gln Arg Leu Ser Arg Thr					
	650		655		660
Leu Gln Ala Ala Ala Phe Leu Asp Phe Leu Trp His Asn Met Lys					
	665		670		675
Leu Pro Val Pro Ala Ala Ala Pro Thr Pro Trp Glu Thr His Lys					
	680		685		690
Gly Ala Leu Pro					

<210> 13

<211> 654

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1848674CD1

<400> 13

Met Glu Arg Ser Gly Pro Ser Glu Val Thr Gly Ser Asp Ala Ser	
1 5	10 15
Gly Pro Asp Pro Gln Leu Ala Val Thr Met Gly Phe Thr Gly Phe	
20 25	30 35
Gly Lys Lys Ala Arg Thr Phe Asp Leu Glu Ala Met Phe Glu Gln	
35 40	45 50
Thr Arg Arg Thr Ala Val Glu Arg Ser Arg Lys Thr Leu Glu Ala	
50 55	60 65
Arg Glu Lys Glu Glu Glu Met Asn Arg Glu Lys Glu Leu Arg Arg	
65 70	75 80
Gln Asn Glu Asp Ile Glu Pro Thr Ser Ser Arg Ser Asn Val Val	
80 85	90 95
Arg Asp Cys Ser Lys Ser Ser Ser Arg Asp Thr Ser Ser Ser Glu	
95 100	105 110
Ser Glu Gln Ser Ser Asp Ser Ser Asp Glu Leu Ile Gly Pro	
110 115	120 125
Pro Leu Pro Pro Lys Met Val Gly Lys Pro Val Asn Phe Met Glu	
125 130	135 140
Glu Asp Ile Leu Gly Pro Leu Pro Pro Pro Leu Asn Glu Glu Glu	
140 145	150 155
Glu Glu Ala Glu Glu Glu Glu Glu Glu Glu Glu Glu Asn	
155 160	165 170
Pro Val His Lys Ile Pro Asp Ser His Glu Ile Thr Leu Lys His	
170 175	180 185
Gly Thr Lys Thr Val Ser Ala Leu Gly Leu Asp Pro Ser Gly Ala	
185 190	195 200
Arg Leu Val Thr Gly Gly Tyr Asp Tyr Asp Val Lys Phe Trp Asp	
200 205	210 215
Phe Ala Gly Met Asp Ala Ser Phe Lys Ala Phe Arg Ser Leu Gln	

	215	220	225
Pro Cys Glu Cys His Gln Ile Lys Ser		Leu Gln Tyr Ser Asn Thr	
	230	235	240
Gly Asp Met Ile Leu Val Val Ser Gly Ser		Ser Gln Ala Lys Val	
	245	250	255
Ile Asp Arg Asp Gly Phe Glu Val Met Glu		Cys Ile Lys Gly Asp	
	260	265	270
Gln Tyr Ile Val Asp Met Ala Asn Thr Lys		Gly His Thr Ala Met	
	275	280	285
Leu His Thr Gly Ser Trp His Pro Lys Ile		Lys Gly Glu Phe Met	
	290	295	300
Thr Cys Ser Asn Asp Ala Thr Val Arg Thr		Trp Glu Val Glu Asn	
	305	310	315
Pro Lys Lys Gln Lys Ser Val Phe Lys Pro		Arg Thr Met Gln Gly	
	320	325	330
Lys Lys Val Ile Pro Thr Thr Cys Thr Tyr		Ser Arg Asp Gly Asn	
	335	340	345
Leu Ile Ala Ala Ala Cys Gln Asn Gly Ser		Ile Gln Ile Trp Asp	
	350	355	360
Arg Asn Leu Thr Val His Pro Lys Phe His		Tyr Lys Gln Ala His	
	365	370	375
Asp Ser Gly Thr Asp Thr Ser Cys Val Thr		Phe Ser Tyr Asp Gly	
	380	385	390
Asn Val Leu Ala Ser Arg Gly Gly Asp Asp		Ser Leu Lys Leu Trp	
	395	400	405
Asp Ile Arg Gln Phe Asn Lys Pro Leu Phe		Ser Ala Ser Gly Leu	
	410	415	420
Pro Thr Met Phe Pro Met Thr Asp Cys Cys		Phe Ser Pro Asp Asp	
	425	430	435
Lys Leu Ile Val Thr Gly Thr Ser Ile Gln		Arg Gly Cys Gly Ser	
	440	445	450
Gly Lys Leu Val Phe Phe Glu Arg Arg Thr		Phe Gln Arg Val Tyr	
	455	460	465
Glu Ile Asp Ile Thr Asp Ala Ser Val Val		Arg Cys Leu Trp His	
	470	475	480
Pro Lys Leu Asn Gln Ile Met Val Gly Thr		Gly Asn Gly Leu Ala	
	485	490	495
Lys Val Tyr Tyr Asp Pro Asn Lys Ser Gln		Arg Gly Ala Lys Leu	
	500	505	510
Cys Val Val Lys Thr Gln Arg Lys Ala Lys		Gln Ala Glu Thr Leu	
	515	520	525
Thr Gln Asp Tyr Ile Ile Thr Pro His Ala		Leu Pro Met Phe Arg	
	530	535	540
Glu Pro Arg Gln Arg Ser Thr Arg Lys Gln		Leu Glu Lys Asp Arg	
	545	550	555
Leu Asp Pro Leu Lys Ser His Lys Pro Glu		Pro Pro Val Ala Gly	
	560	565	570
Pro Gly Arg Gly Gly Arg Val Gly Thr His		Gly Gly Thr Leu Ser	
	575	580	585
Ser Tyr Ile Val Lys Asn Ile Ala Leu Asp		Lys Thr Asp Asp Ser	
	590	595	600
Asn Pro Arg Glu Ala Ile Leu Arg His Ala		Lys Ala Ala Glu Asp	
	605	610	615
Ser Pro Tyr Trp Val Ser Pro Ala Tyr Ser		Lys Thr Gln Pro Lys	
	620	625	630

Thr Met Phe Ala Gln Val Glu Ser Asp Asp Glu Glu Ala Lys Asn
 635 640 645
 Glu Pro Glu Trp Lys Lys Arg Lys Ile
 650

<210> 14
 <211> 180
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2012970CD1

<400> 14
 Met Glu Ala Asn Met Pro Lys Arg Lys Glu Pro Gly Arg Ser Leu
 1 5 10 15
 Arg Ile Lys Val Ile Ser Met Gly Asn Ala Glu Val Gly Lys Ser
 20 25 30
 Cys Ile Ile Lys Arg Tyr Cys Glu Lys Arg Phe Val Ser Lys Tyr
 35 40 45
 Leu Ala Thr Ile Gly Ile Asp Tyr Gly Val Thr Lys Val His Val
 50 55 60
 Arg Asp Arg Glu Ile Lys Val Asn Ile Phe Asp Met Ala Gly His
 65 70 75
 Pro Phe Phe Tyr Glu Val Arg Asn Glu Phe Tyr Lys Asp Thr Gln
 80 85 90
 Gly Val Ile Leu Val Tyr Asp Val Gly Gln Lys Asp Ser Phe Asp
 95 100 105
 Ala Leu Asp Ala Trp Leu Ala Glu Met Lys Gln Glu Leu Gly Pro
 110 115 120
 His Gly Asn Met Glu Asn Ile Ile Phe Val Val Cys Ala Asn Lys
 125 130 135
 Ile Asp Cys Thr Lys His Arg Cys Val Asp Glu Ser Glu Gly Arg
 140 145 150
 Leu Trp Ala Glu Ser Lys Gly Phe Leu Tyr Phe Glu Thr Ser Ala
 155 160 165
 Gln Thr Gly Glu Gly Ile Asn Glu Met Phe Gln Ile His Leu Gly
 170 175 180

<210> 15
 <211> 374
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2254315CD1

<400> 15
 Met Ala Ala Ser Ala Ala Ala Glu Leu Gln Ala Ser Gly Gly
 1 5 10 15
 Pro Arg His Pro Val Cys Leu Leu Val Leu Gly Met Ala Gly Ser

	20		25		30
Gly Lys Thr Thr	Phe Val Gln Arg Leu	Thr Gly His Leu His	Ala		
	35		40		45
Gln Gly Thr Pro	Pro Tyr Val Ile Asn	Leu Asp Pro Ala Val His			
	50		55		60
Glu Val Pro Phe	Pro Ala Asn Ile Asp	Ile Arg Asp Thr Val Lys			
	65		70		75
Tyr Lys Glu Val	Met Lys Gln Tyr Gly	Leu Gly Pro Asn Gly Gly			
	80		85		90
Ile Val Thr Ser	Leu Asn Leu Phe Ala	Thr Arg Phe Asp Gln Val			
	95		100		105
Met Lys Phe Ile	Glu Lys Ala Gln Asn	Met Ser Lys Tyr Val Leu			
	110		115		120
Ile Asp Thr Pro	Gly Gln Ile Glu Val	Phe Thr Trp Ser Ala Ser			
	125		130		135
Gly Thr Ile Ile	Thr Glu Ala Leu Ala	Ser Ser Phe Pro Thr Val			
	140		145		150
Val Ile Tyr Val	Met Asp Thr Ser Arg	Ser Thr Asn Pro Val Thr			
	155		160		165
Phe Met Ser Asn	Met Leu Tyr Ala Cys	Ser Ile Leu Tyr Lys Thr			
	170		175		180
Lys Leu Pro Phe	Ile Val Val Met Asn	Lys Thr Asp Ile Ile Asp			
	185		190		195
His Ser Phe Ala	Val Glu Trp Met Gln	Asp Phe Glu Ala Phe Gln			
	200		205		210
Asp Ala Leu Asn	Gln Glu Thr Thr Tyr	Val Ser Asn Leu Thr Arg			
	215		220		225
Ser Met Ser Leu	Val Leu Asp Glu Phe	Tyr Ser Ser Leu Arg Val			
	230		235		240
Val Gly Val Ser	Ala Val Leu Gly Thr	Gly Leu Asp Glu Leu Phe			
	245		250		255
Val Gln Val Thr	Ser Ala Ala Glu Glu	Tyr Glu Arg Glu Tyr Arg			
	260		265		270
Pro Glu Tyr Glu	Arg Leu Lys Lys Ser	Leu Ala Asn Ala Glu Ser			
	275		280		285
Gln Gln Gln Arg	Glu Gln Leu Glu Arg	Leu Arg Lys Asp Met Gly			
	290		295		300
Ser Val Ala Leu	Asp Ala Gly Thr Ala	Lys Asp Ser Leu Ser Pro			
	305		310		315
Val Leu His Pro	Ser Asp Leu Ile Leu	Thr Arg Gly Thr Leu Asp			
	320		325		330
Glu Glu Asp Glu	Glu Ala Asp Ser Asp	Thr Asp Asp Ile Asp His			
	335		340		345
Arg Val Thr Glu	Glu Ser His Glu Glu	Pro Ala Phe Gln Asn Phe			
	350		355		360
Met Gln Glu Ser	Met Ala Gln Tyr Trp	Lys Arg Asn Asn Lys			
	365		370		

<210> 16

<211> 649

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No. 2415545CD1

<400> 16

Met	Glu	Gly	Ala	Gly	Tyr	Arg	Val	Val	Phe	Glu	Lys	Gly	Gly	Val	
1				5					10					15	
Tyr	Leu	His	Thr	Ser	Ala	Lys	Lys	Tyr	Gln	Asp	Arg	Asp	Ser	Leu	
				20					25					30	
Ile	Ala	Gly	Val	Ile	Arg	Val	Val	Glu	Lys	Asp	Asn	Asp	Val	Leu	
				35					40					45	
Leu	His	Trp	Ala	Pro	Val	Glu	Glu	Ala	Gly	Asp	Ser	Thr	Gln	Ile	
				50					55					60	
Leu	Phe	Ser	Lys	Lys	Asp	Ser	Ser	Gly	Gly	Asp	Ser	Cys	Ala	Ser	
				65					70					75	
Glu	Glu	Glu	Pro	Thr	Phe	Asp	Pro	Gly	Tyr	Glu	Pro	Asp	Trp	Ala	
				80					85					90	
Val	Ile	Ser	Thr	Val	Arg	Pro	Gln	Pro	Cys	His	Ser	Glu	Pro	Thr	
				95					100					105	
Arg	Gly	Ala	Glu	Pro	Ser	Cys	Pro	Gln	Gly	Ser	Trp	Ala	Phe	Ser	
				110					115					120	
Val	Ser	Leu	Gly	Glu	Leu	Lys	Ser	Ile	Arg	Arg	Ser	Lys	Pro	Gly	
				125					130					135	
Leu	Ser	Trp	Ala	Tyr	Leu	Val	Leu	Val	Thr	Gln	Ala	Gly	Gly	Ser	
				140					145					150	
Leu	Pro	Ala	Leu	His	Phe	His	Arg	Gly	Gly	Thr	Arg	Ala	Leu	Leu	
				155					160					165	
Arg	Val	Leu	Ser	Arg	Tyr	Leu	Leu	Leu	Ala	Ser	Ser	Pro	Gln	Asp	
				170					175					180	
Ser	Arg	Leu	Tyr	Leu	Val	Phe	Pro	His	Asp	Ser	Ser	Ala	Leu	Ser	
				185					190					195	
Asn	Ser	Phe	His	His	Leu	Gln	Leu	Phe	Asp	Gln	Asp	Ser	Ser	Asn	
				200					205					210	
Val	Val	Ser	Arg	Phe	Leu	Gln	Asp	Pro	Tyr	Ser	Thr	Thr	Phe	Ser	
				215					220					225	
Ser	Phe	Ser	Arg	Val	Thr	Asn	Phe	Phe	Arg	Gly	Ala	Leu	Gln	Pro	
				230					235					240	
Gln	Pro	Glu	Gly	Ala	Ala	Ser	Asp	Leu	Pro	Pro	Pro	Pro	Asp	Asp	
				245					250					255	
Glu	Pro	Glu	Pro	Gly	Phe	Glu	Val	Ile	Ser	Cys	Val	Glu	Leu	Gly	
				260					265					270	
Pro	Arg	Pro	Thr	Val	Glu	Arg	Gly	Pro	Pro	Val	Thr	Glu	Glu	Glu	
				275					280					285	
Trp	Ala	Arg	His	Val	Gly	Pro	Glu	Gly	Arg	Leu	Gln	Gln	Val	Pro	
				290					295					300	
Glu	Leu	Lys	Asn	Arg	Ile	Phe	Ser	Gly	Gly	Leu	Ser	Pro	Ser	Leu	
				305					310					315	
Arg	Arg	Glu	Ala	Trp	Lys	Phe	Leu	Leu	Gly	Tyr	Leu	Ser	Trp	Glu	
				320					325					330	
Gly	Thr	Ala	Glu	Glu	His	Lys	Ala	His	Ile	Arg	Lys	Lys	Thr	Asp	
				335					340					345	
Glu	Tyr	Phe	Arg	Met	Lys	Leu	Gln	Trp	Lys	Ser	Val	Ser	Pro	Glu	
				350					355					360	
Gln	Glu	Arg	Arg	Asn	Ser	Leu	Leu	His	Gly	Tyr	Arg	Ser	Leu	Ile	
				365					370					375	
Glu	Arg	Asp	Val	Ser	Arg	Thr	Asp	Arg	Thr	Asn	Lys	Phe	Tyr	Glu	

380 385 390
 Gly Pro Glu Asn Pro Gly Leu Gly Leu Leu Asn Asp Ile Leu Leu
 395 400 405
 Thr Tyr Cys Met Tyr His Phe Asp Leu Gly Tyr Val Gln Gly Met
 410 415 420
 Ser Asp Leu Leu Ser Pro Ile Leu Tyr Val Ile Gln Asn Glu Val
 425 430 435
 Asp Ala Phe Trp Cys Phe Cys Gly Phe Met Glu Leu Val Gln Gly
 440 445 450
 Asn Phe Glu Glu Ser Gln Glu Thr Met Lys Arg Gln Leu Gly Arg
 455 460 465
 Leu Leu Leu Leu Leu Arg Val Leu Asp Pro Leu Leu Cys Asp Phe
 470 475 480
 Leu Asp Ser Gln Asp Ser Gly Ser Leu Cys Phe Cys Phe Arg Trp
 485 490 495
 Leu Leu Ile Trp Phe Lys Arg Glu Phe Pro Phe Pro Asp Val Leu
 500 505 510
 Arg Leu Trp Glu Val Leu Trp Thr Gly Leu Pro Gly Pro Asn Leu
 515 520 525
 His Leu Leu Val Ala Cys Ala Ile Leu Asp Met Glu Arg Asp Thr
 530 535 540
 Leu Met Leu Ser Gly Phe Gly Ser Asn Glu Ile Leu Lys His Ile
 545 550 555
 Asn Glu Leu Thr Met Lys Leu Ser Val Glu Asp Val Leu Thr Arg
 560 565 570
 Ala Glu Ala Leu His Arg Gln Leu Thr Ala Cys Thr Arg Ala Ala
 575 580 585
 Pro Gln Arg Ala Gly Asp Pro Gly Ala Gly Pro Ala Thr Gln Ser
 590 595 600
 Pro Thr Ala Pro Arg Pro Pro Pro Pro Arg Cys Leu Cys Thr Pro
 605 610 615
 Thr Arg Ala Pro Pro Thr Pro Pro Pro Ser Thr Asp Thr Ala Pro
 620 625 630
 Gln Pro Asp Ser Ser Leu Glu Ile Leu Pro Glu Glu Glu Asp Glu
 635 640 645
 Gly Ala Asp Ser

<210> 17
 <211> 698
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2707969CD1

<400> 17
 Met Cys His Asp Asp Asp Lys Asp Pro Val Leu Arg Val Phe
 1 5 10 15
 Asp Ser Arg Val Asp Lys Ile Arg Leu Leu Asn Val Arg Thr Pro
 20 25 30
 Thr Leu Arg Thr Ser Met Tyr Gln Lys Cys Thr Thr Val Asp Glu
 35 40 45
 Ala Glu Lys Ala Ile Glu Leu Arg Leu Ala Lys Ile Asp His Thr

	50		55		60
Ala Ile His Pro	His Leu Leu Asp Met	Lys Ile Gly Gln Gly	Lys		
	65		70		75
Tyr Glu Pro Gly	Phe Phe Pro Lys Leu Gln	Ser Asp Val Leu Ser			
	80		85		90
Thr Gly Pro Ala	Ser Asn Lys Trp Thr Lys	Arg Asn Ala Pro Ala			
	95		100		105
Gln Trp Arg Arg	Lys Asp Arg Gln Lys Gln	His Thr Glu His Leu			
	110		115		120
Arg Leu Asp Asn	Asp Gln Arg Glu Lys Tyr	Ile Gln Glu Ala Arg			
	125		130		135
Thr Met Gly Ser	Thr Ile Arg Gln Pro Lys	Leu Ser Asn Leu Ser			
	140		145		150
Pro Ser Val Ile	Ala Gln Thr Asn Trp Lys	Phe Val Glu Gly Leu			
	155		160		165
Leu Lys Glu Cys	Arg Asn Lys Thr Lys Arg	Met Leu Val Glu Lys			
	170		175		180
Met Gly Arg Glu	Ala Val Glu Leu Gly His	Gly Glu Val Asn Ile			
	185		190		195
Thr Gly Val Glu	Glu Asn Thr Leu Ile Ala	Ser Leu Cys Asp Leu			
	200		205		210
Leu Glu Arg Ile	Trp Ser His Gly Leu Gln	Val Lys Gln Gly Lys			
	215		220		225
Ser Ala Leu Trp	Ser His Leu Leu His Tyr	Gln Asp Asn Arg Gln			
	230		235		240
Arg Lys Leu Thr	Ser Gly Ser Leu Ser Thr	Ser Gly Ile Leu Leu			
	245		250		255
Asp Ser Glu Arg	Arg Lys Ser Asp Ala Ser	Ser Leu Met Pro Pro			
	260		265		270
Leu Arg Ile Ser	Leu Ile Gln Asp Met Arg	His Ile Gln Asn Ile			
	275		280		285
Gly Glu Ile Lys	Thr Asp Val Gly Lys Ala	Arg Ala Trp Val Arg			
	290		295		300
Leu Ser Met Glu	Lys Lys Leu Leu Ser Arg	His Leu Lys Gln Leu			
	305		310		315
Leu Ser Asp His	Glu Leu Thr Lys Lys Leu	Tyr Lys Arg Tyr Ala			
	320		325		330
Phe Leu Arg Cys	Asp Asp Glu Lys Glu Gln	Phe Leu Tyr His Leu			
	335		340		345
Leu Ser Phe Asn	Ala Val Asp Tyr Phe Cys	Phe Thr Asn Val Phe			
	350		355		360
Thr Thr Ile Leu	Ile Pro Tyr His Ile Leu	Ile Val Pro Ser Lys			
	365		370		375
Lys Leu Gly Gly	Ser Met Phe Thr Ala Asn	Pro Trp Ile Cys Ile			
	380		385		390
Ser Gly Glu Leu	Gly Glu Thr Gln Ile Met	Gln Ile Pro Arg Asn			
	395		400		405
Val Leu Glu Met	Thr Phe Glu Cys Gln Asn	Leu Gly Lys Leu Thr			
	410		415		420
Thr Val Gln Ile	Gly His Asp Asn Ser Gly	Leu Tyr Ala Lys Trp			
	425		430		435
Leu Val Glu Tyr	Val Met Val Arg Asn Glu	Ile Thr Gly His Thr			
	440		445		450
Tyr Lys Phe Pro	Cys Gly Arg Trp Leu Gly	Lys Gly Met Asp Asp			
	455		460		465

Gly Ser Leu Glu Arg Ile Leu Val Gly Glu Leu Leu Thr Ser Gln
 470 475
 Pro Glu Val Asp Glu Arg Pro Cys Arg Thr Pro Pro Leu Gln Gln
 485 490
 Ser Pro Ser Val Ile Arg Arg Leu Val Thr Ile Ser Pro Asn Asn
 500 505
 Lys Pro Lys Leu Asn Thr Gly Gln Ile Gln Glu Ser Ile Gly Glu
 515 520
 Ala Val Asn Gly Ile Val Lys His Phe His Lys Pro Glu Lys Glu
 530 535
 Arg Gly Ser Leu Thr Leu Leu Leu Cys Gly Glu Cys Gly Leu Val
 545 550
 Ser Ala Leu Glu Gln Ala Phe Gln His Gly Phe Lys Ser Pro Arg
 560 565
 Leu Phe Lys Asn Val Phe Ile Trp Asp Phe Leu Glu Lys Ala Gln
 575 580
 Thr Tyr Tyr Glu Thr Leu Glu Lys Asn Glu Val Val Pro Glu Glu
 590 595
 Asn Trp His Thr Arg Ala Arg Asn Phe Cys Arg Phe Val Thr Ala
 605 610
 Ile Asn Asn Thr Pro Arg Asn Ile Gly Lys Asp Gly Lys Phe Gln
 620 625
 Met Leu Val Cys Leu Gly Ala Arg Asp His Leu Leu His His Trp
 635 640
 Ile Ala Leu Leu Ala Asp Cys Pro Ile Thr Ala His Met Tyr Glu
 650 655
 Asp Val Ala Leu Ile Lys Asp His Thr Leu Val Asn Ser Leu Ile
 665 670
 Arg Val Leu Gln Thr Leu Gln Glu Phe Asn Ile Thr Leu Glu Thr
 680 685
 Ser Leu Val Lys Gly Ile Asp Ile
 695

<210> 18

<211> 396

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2817769CD1

<400> 18

Met Pro Pro Lys Lys Gly Gly Asp Gly Ile Lys Pro Pro Pro Ile
 1 5 10 15
 Ile Gly Arg Phe Gly Thr Ser Leu Lys Ile Gly Ile Val Gly Leu
 20 25 30
 Pro Asn Val Gly Lys Ser Thr Phe Phe Asn Val Leu Thr Asn Ser
 35 40 45
 Gln Ala Ser Ala Glu Asn Phe Pro Phe Cys Thr Ile Asp Pro Asn
 50 55 60
 Glu Ser Arg Val Pro Val Pro Asp Glu Arg Phe Asp Phe Leu Cys
 65 70 75
 Gln Tyr His Lys Pro Ala Ser Lys Ile Pro Ala Phe Leu Asn Val
 80 85 90

Val Asp Ile Ala Gly Leu Val Lys Gly Ala His Asn Gly Gln Gly
 95 100
 Leu Gly Asn Ala Phe Leu Ser His Ile Ser Ala Cys Asp Gly Ile
 110 115
 Phe His Leu Thr Arg Ala Phe Glu Asp Asp Asp Ile Thr His Val
 125 130
 Glu Gly Ser Val Asp Pro Ile Arg Asp Ile Glu Ile Ile His Glu
 140 145
 Glu Leu Gln Leu Lys Asp Glu Glu Met Ile Gly Pro Ile Ile Asp
 155 160
 Lys Leu Glu Lys Val Ala Val Arg Gly Gly Asp Lys Lys Leu Lys
 170 175
 Pro Glu Tyr Asp Ile Met Cys Lys Val Lys Ser Trp Val Ile Asp
 185 190
 Gln Lys Lys Pro Val Arg Phe Tyr His Asp Trp Asn Asp Lys Glu
 200 205
 Ile Glu Val Leu Asn Lys His Leu Phe Leu Thr Ser Lys Pro Met
 215 220
 Val Tyr Leu Val Asn Leu Ser Glu Lys Asp Tyr Ile Arg Lys Lys
 230 235
 Asn Lys Trp Leu Ile Lys Ile Lys Glu Trp Val Asp Lys Tyr Asp
 245 250
 Pro Gly Ala Leu Val Ile Pro Phe Ser Gly Ala Leu Glu Leu Lys
 260 265
 Leu Gln Glu Leu Ser Ala Glu Glu Arg Gln Lys Tyr Leu Glu Ala
 275 280
 Asn Met Thr Gln Ser Ala Leu Pro Lys Ile Ile Lys Ala Gly Phe
 290 295
 Ala Ala Leu Gln Leu Glu Tyr Phe Phe Thr Ala Gly Pro Asp Glu
 305 310
 Val Arg Ala Trp Thr Ile Arg Lys Gly Thr Lys Ala Pro Gln Ala
 320 325
 Ala Gly Lys Ile His Thr Asp Phe Glu Lys Gly Phe Ile Met Ala
 335 340
 Glu Val Met Lys Tyr Glu Asp Phe Lys Glu Glu Gly Ser Glu Asn
 350 355
 Ala Val Lys Ala Ala Gly Lys Tyr Arg Gln Gln Gly Arg Asn Tyr
 365 370
 Ile Val Glu Asp Gly Asp Ile Ile Phe Phe Lys Phe Asn Thr Pro
 380 385
 Gln Gln Pro Lys Lys Lys
 395

<210> 19

<211> 634

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2917557CD1

<400> 19

Met Ser Ser Asp Ser Glu Tyr Asp Ser Asp Asp Asp Arg Thr Lys

1	5	10	15
Glu Glu Arg Ala	Tyr Asp Lys Ala Lys	Arg Arg Ile Glu Lys Arg	
	20	25	30
Arg Leu Glu His	Ser Lys Asn Val Asn Thr	Glu Lys Leu Arg Ala	
	35	40	45
Pro Ile Ile Cys	Val Leu Gly His Val Asp	Thr Gly Lys Thr Lys	
	50	55	60
Ile Leu Asp Lys	Leu Arg His Thr His Val	Gln Asp Gly Glu Ala	
	65	70	75
Gly Gly Ile Thr	Gln Gln Ile Gly Ala Thr	Asn Val Pro Leu Glu	
	80	85	90
Ala Ile Asn Glu	Gln Thr Lys Met Ile Lys	Asn Phe Asp Arg Glu	
	95	100	105
Asn Val Arg Ile	Pro Gly Met Leu Ile Ile	Asp Thr Pro Gly His	
	110	115	120
Glu Ser Phe Ser	Asn Leu Arg Asn Arg Gly	Ser Ser Leu Cys Asp	
	125	130	135
Ile Ala Ile Leu	Val Val Asp Ile Met His	Gly Leu Glu Pro Gln	
	140	145	150
Thr Ile Glu Ser	Ile Asn Leu Leu Lys Ser	Lys Lys Cys Pro Phe	
	155	160	165
Ile Val Ala Leu	Asn Lys Ile Asp Arg Leu	Tyr Asp Trp Lys Lys	
	170	175	180
Ser Pro Asp Ser	Asp Val Ala Ala Thr Leu	Lys Lys Gln Lys Lys	
	185	190	195
Asn Thr Lys Asp	Glu Phe Glu Glu Arg Ala	Lys Ala Ile Ile Val	
	200	205	210
Glu Phe Ala Gln	Gln Gly Leu Asn Ala Ala	Leu Phe Tyr Glu Asn	
	215	220	225
Lys Asp Pro Arg	Thr Phe Val Ser Leu Val	Pro Thr Ser Ala His	
	230	235	240
Thr Gly Asp Gly	Met Gly Ser Leu Ile Tyr	Leu Leu Val Glu Leu	
	245	250	255
Thr Gln Thr Met	Leu Ser Lys Arg Leu Ala	His Cys Glu Glu Leu	
	260	265	270
Arg Ala Gln Val	Met Glu Val Lys Ala Leu	Pro Gly Met Gly Thr	
	275	280	285
Thr Ile Asp Val	Ile Leu Ile Asn Gly Arg	Leu Lys Glu Gly Asp	
	290	295	300
Thr Ile Ile Val	Pro Gly Val Glu Gly Pro	Ile Val Thr Gln Ile	
	305	310	315
Arg Gly Leu Leu	Leu Pro Pro Pro Met Lys	Glu Leu Arg Val Lys	
	320	325	330
Asn Gln Tyr Glu	Lys His Lys Glu Val Glu	Ala Ala Gln Gly Val	
	335	340	345
Lys Ile Leu Gly	Lys Asp Leu Glu Lys Thr	Leu Ala Gly Leu Pro	
	350	355	360
Leu Leu Val Ala	Tyr Lys Glu Asp Glu Ile	Pro Val Leu Lys Asp	
	365	370	375
Glu Leu Ile His	Glu Leu Lys Gln Thr Leu	Asn Ala Ile Lys Leu	
	380	385	390
Glu Glu Lys Gly	Val Tyr Val Gln Ala Ser	Thr Leu Gly Ser Leu	
	395	400	405
Glu Ala Leu Leu	Glu Phe Leu Lys Thr Ser	Glu Val Pro Tyr Ala	
	410	415	420

Gly Ile Asn Ile Gly Pro Val His Lys Lys Asp Val Met Lys Ala
 425 430 435
 Ser Val Met Leu Glu His Asp Pro Gln Tyr Ala Val Ile Leu Ala
 440 445 450
 Phe Asp Val Arg Ile Glu Arg Asp Ala Gln Glu Met Ala Asp Ser
 455 460 465
 Leu Gly Val Arg Ile Phe Ser Ala Glu Ile Ile Tyr His Leu Phe
 470 475 480
 Asp Ala Phe Thr Lys Tyr Arg Gln Asp Tyr Lys Lys Gln Lys Gln
 485 490 495
 Glu Glu Phe Lys His Ile Ala Val Phe Pro Cys Lys Ile Lys Ile
 500 505 510
 Leu Pro Gln Tyr Ile Phe Asn Ser Arg Asp Pro Ile Val Met Gly
 515 520 525
 Val Thr Val Glu Ala Gly Gln Val Lys Gln Gly Thr Pro Met Cys
 530 535 540
 Val Pro Ser Lys Asn Phe Val Asp Ile Gly Ile Val Thr Ser Ile
 545 550 555
 Glu Ile Asn His Lys Gln Val Asp Val Ala Lys Lys Gly Gln Glu
 560 565 570
 Val Cys Val Lys Ile Glu Pro Ile Pro Gly Glu Ser Pro Lys Met
 575 580 585
 Phe Gly Arg His Phe Glu Ala Thr Asp Ile Leu Val Ser Lys Ile
 590 595 600
 Ser Arg Gln Ser Ile Asp Ala Leu Lys Asp Trp Phe Arg Asp Glu
 605 610 615
 Met Gln Lys Ser Asp Trp Gln Leu Ile Val Glu Leu Lys Lys Val
 620 625 630
 Phe Glu Ile Ile

<210> 20

<211> 196

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3421335CD1

<400> 20

Met Gly Ser Val Asn Ser Arg Gly His Lys Ala Glu Ala Gln Val
 1 5 10 15
 Val Met Met Gly Leu Asp Ser Ala Gly Lys Thr Thr Leu Leu Tyr
 20 25 30
 Lys Leu Lys Gly His Gln Leu Val Glu Thr Leu Pro Thr Val Gly
 35 40 45
 Phe Asn Val Glu Pro Leu Lys Ala Pro Gly His Val Ser Leu Thr
 50 55 60
 Leu Trp Asp Val Gly Gly Gln Ala Pro Leu Arg Ala Ser Trp Lys
 65 70 75
 Asp Tyr Leu Glu Gly Thr Asp Ile Leu Val Tyr Val Leu Asp Ser
 80 85 90
 Thr Asp Glu Ala Arg Leu Pro Glu Ser Ala Ala Glu Leu Thr Glu
 95 100 105

Val Leu Asn Asp	Pro Asn Met Ala Gly	Val Pro Phe Leu Val Leu
110	115	120
Ala Asn Lys Gln Glu Ala Pro Asp Ala	Leu Pro Leu Leu Lys Ile	
125	130	135
Arg Asn Arg Leu Ser Leu Glu Arg Phe	Gln Asp His Cys Trp Glu	
140	145	150
Leu Arg Gly Cys Ser Ala Leu Thr Gly	Glu Gly Leu Pro Glu Ala	
155	160	165
Leu Gln Ser Leu Trp Ser Leu Leu Lys	Ser Arg Ser Cys Met Cys	
170	175	180
Leu Gln Ala Arg Ala His Gly Ala Glu	Arg Gly Asp Ser Lys Arg	
185	190	195
Ser		

<210> 21

<211> 446

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 605761CD1

<400> 21

Met Ala Ala Arg Lys	Gly Arg Arg Arg Thr	Cys Glu Thr Gly Glu
1	5	10
Pro Met Glu Ala Glu	Ser Gly Asp Thr Ser	Ser Glu Gly Pro Ala
20	25	30
Gln Val Tyr Leu Pro	Gly Arg Gly Pro Pro	Leu Arg Glu Gly Glu
35	40	45
Glu Leu Val Met Asp	Glu Glu Ala Tyr Val	Leu Tyr His Arg Ala
50	55	60
Gln Thr Gly Ala Pro	Cys Leu Ser Phe Asp	Ile Val Arg Asp His
65	70	75
Leu Gly Asp Asn Arg	Thr Glu Leu Pro Leu	Thr Leu Tyr Leu Cys
80	85	90
Ala Gly Thr Gln Ala	Glu Ser Ala Gln Ser	Asn Arg Leu Met Met
95	100	105
Leu Arg Met His Asn	Leu His Gly Thr Lys	Pro Pro Pro Ser Glu
110	115	120
Gly Ser Asp Glu Glu	Glu Glu Glu Glu Asp	Glu Glu Asp Glu Glu
125	130	135
Glu Arg Lys Pro Gln	Leu Glu Leu Ala Met	Val Pro His Tyr Gly
140	145	150
Gly Ile Asn Arg Val	Arg Val Ser Trp Leu	Gly Glu Glu Pro Val
155	160	165
Ala Gly Val Trp Ser	Glu Lys Gly Gln Val	Glu Val Phe Ala Leu
170	175	180
Arg Arg Leu Leu Gln	Val Val Glu Glu Pro	Gln Ala Leu Ala Ala
185	190	195
Phe Leu Arg Asp Glu	Gln Ala Gln Met Lys	Pro Ile Phe Ser Phe
200	205	210
Ala Gly His Met Gly	Glu Gly Phe Ala Leu	Asp Trp Ser Pro Arg
215	220	225

Val Thr Gly Arg Leu Leu Thr Gly Asp Cys Gln Lys Asn Ile His
 230 240
 Leu Trp Thr Pro Thr Asp Gly Gly Ser Trp His Val Asp Gln Arg
 245 250
 Pro Phe Val Gly His Thr Arg Ser Val Glu Asp Leu Gln Trp Ser
 260 265
 Pro Thr Glu Asn Thr Val Phe Ala Ser Cys Ser Ala Asp Ala Ser
 275 280
 Ile Arg Ile Trp Asp Ile Arg Ala Ala Pro Ser Lys Ala Cys Met
 290 295
 Leu Thr Thr Ala Thr Ala His Asp Gly Asp Val Asn Val Ile Ser
 305 310
 Trp Ser Arg Arg Glu Pro Phe Leu Leu Ser Gly Gly Asp Asp Gly
 320 325
 Ala Leu Lys Ile Trp Asp Leu Arg Gln Phe Lys Ser Gly Ser Pro
 335 340
 Val Ala Thr Phe Lys Gln His Val Ala Pro Val Thr Ser Val Glu
 350 355
 Trp His Pro Gln Asp Ser Gly Val Phe Ala Ala Ser Gly Ala Asp
 365 370
 His Gln Ile Thr Gln Trp Asp Leu Ala Val Glu Arg Asp Pro Glu
 380 385
 Ala Gly Asp Val Glu Ala Asp Pro Gly Leu Ala Asp Leu Pro Gln
 395 400
 Gln Leu Leu Phe Val His Gln Gly Glu Thr Glu Leu Lys Glu Leu
 410 415
 His Trp His Pro Gln Cys Pro Gly Leu Leu Val Ser Thr Ala Leu
 425 430
 Ser Gly Phe Thr Ile Phe Arg Thr Ile Ser Val
 440 445

<210> 22

<211> 265

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 483862CD1

<400> 22

Met Ser Ser Gly Leu Arg Ala Ala Asp Phe Pro Arg Trp Lys Arg
 1 5 10 15
 His Ile Ser Glu Gln Leu Arg Arg Arg Asp Arg Leu Gln Arg Gln
 20 25 30
 Ala Phe Glu Glu Ile Ile Leu Gln Tyr Asn Lys Leu Leu Glu Lys
 35 40 45
 Ser Asp Leu His Ser Val Leu Ala Gln Lys Leu Gln Ala Glu Lys
 50 55 60
 His Asp Val Pro Asn Arg His Glu Ile Ser Pro Gly His Asp Gly
 65 70 75
 Thr Trp Asn Asp Asn Gln Leu Gln Glu Met Ala Gln Leu Arg Ile
 80 85 90
 Lys His Gln Glu Glu Leu Thr Glu Leu His Lys Lys Arg Gly Glu

	95		100		105
Leu	Ala	Gln	Leu	Val	Ile
	110		115		120
Asp	Arg	Glu	Met	Gln	Met
	125		130		135
Gln	Thr	Ile	Ser	Asp	Leu
	140		145		150
Lys	Leu	Cys	Asp	Leu	Glu
	155		160		165
Tyr	Asp	Ala	Leu	Gln	Ile
	170		175		180
Arg	Lys	Thr	Thr	Glu	Asn
	185		190		195
Ala	Glu	Lys	Ala	Gln	Glu
	200		205		210
Lys	Asp	Ser	Arg	Arg	Gln
	215		220		225
Glu	Ala	Ala	Lys	Glu	Pro
	230		235		240
Glu	Val	Ile	Val	Asp	Glu
	245		250		255
Pro	Val	Arg	Ala	Ile	Ser
	260		265		

<210> 23

<211> 185

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1256777CD1

<400> 23

Met	Leu	Lys	Ala	Lys	Ile	Leu	Phe	Val	Gly	Pro	Cys	Glu	Ser	Gly
1		5							10					15
Lys	Thr	Val	Leu	Ala	Asn	Phe	Leu	Thr	Glu	Ser	Ser	Asp	Ile	Thr
		20							25					30
Glu	Tyr	Ser	Pro	Thr	Gln	Gly	Val	Arg	Ile	Leu	Glu	Phe	Glu	Asn
		35							40					45
Pro	His	Val	Thr	Ser	Asn	Asn	Lys	Gly	Thr	Gly	Cys	Glu	Phe	Glu
		50							55					60
Leu	Trp	Asp	Cys	Gly	Gly	Asp	Ala	Lys	Phe	Glu	Ser	Cys	Trp	Pro
		65							70					75
Ala	Leu	Met	Lys	Asp	Ala	His	Gly	Val	Val	Ile	Val	Phe	Asn	Ala
		80							85					90
Asp	Ile	Pro	Ser	His	Arg	Lys	Glu	Met	Glu	Met	Trp	Tyr	Ser	Cys
		95							100					105
Phe	Val	Gln	Gln	Pro	Ser	Leu	Gln	Asp	Thr	Gln	Cys	Met	Leu	Ile
		110							115					120
Ala	His	His	Lys	Pro	Gly	Ser	Gly	Asp	Asp	Lys	Gly	Ser	Leu	Ser
		125							130					135
Leu	Ser	Pro	Pro	Leu	Asn	Lys	Leu	Lys	Leu	Val	His	Ser	Asn	Leu
		140							145					150

Glu Asp Asp Pro Glu Glu Ile Arg Met Glu Phe Ile Lys Tyr Leu
 155 160
 Lys Ser Ile Ile Asn Ser Met Ser Glu Ser Arg Asp Arg Glu Glu
 170 175 180
 Met Ser Ile Met Thr
 185

<210> 24
 <211> 554
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2198779CD1

<400> 24
 Met Gly Ser Arg Asn Ser Ser Ser Ala Gly Ser Gly Ser Gly Asp
 1 5 10 15
 Pro Ser Glu Gly Leu Pro Arg Arg Gly Ala Gly Leu Arg Arg Ser
 20 25 30
 Glu Glu Glu Glu Glu Glu Asp Glu Asp Val Asp Leu Ala Gln Val
 35 40 45
 Leu Ala Tyr Leu Leu Arg Arg Gly Gln Val Arg Leu Val Gln Gly
 50 55 60
 Gly Gly Ala Ala Asn Leu Gln Phe Ile Gln Ala Leu Leu Asp Ser
 65 70 75
 Glu Glu Glu Asn Asp Arg Ala Trp Asp Gly Arg Leu Gly Asp Arg
 80 85 90
 Tyr Asn Pro Pro Val Asp Ala Thr Pro Asp Thr Arg Glu Leu Glu
 95 100 105
 Phe Asn Glu Ile Lys Thr Gln Val Glu Leu Ala Thr Gly Gln Leu
 110 115 120
 Gly Leu Arg Arg Ala Ala Gln Lys His Ser Phe Pro Arg Met Leu
 125 130 135
 His Gln Arg Glu Arg Gly Leu Cys His Arg Gly Ser Phe Ser Leu
 140 145 150
 Gly Glu Gln Ser Arg Val Ile Ser His Phe Leu Pro Asn Asp Leu
 155 160 165
 Gly Phe Thr Asp Ser Tyr Ser Gln Lys Ala Phe Cys Gly Ile Tyr
 170 175 180
 Ser Lys Asp Gly Gln Ile Phe Met Ser Ala Cys Gln Asp Gln Thr
 185 190 195
 Ile Arg Leu Tyr Asp Cys Arg Tyr Gly Arg Phe Arg Lys Phe Lys
 200 205 210
 Ser Ile Lys Ala Arg Asp Val Gly Trp Ser Val Leu Asp Val Ala
 215 220 225
 Phe Thr Pro Asp Gly Asn His Phe Leu Tyr Ser Ser Trp Ser Asp
 230 235 240
 Tyr Ile His Ile Cys Asn Ile Tyr Gly Glu Gly Asp Thr His Thr
 245 250 255
 Ala Leu Asp Leu Arg Pro Asp Glu Arg Arg Phe Ala Val Phe Ser
 260 265 270
 Ile Ala Val Ser Ser Asp Gly Arg Glu Val Leu Gly Gly Ala Asn
 275 280 285

Asp Gly Cys Leu Tyr Val Phe Asp Arg Glu Gln Asn Arg Arg Thr
 290 295 300
 Leu Gln Ile Glu Ser His Glu Asp Asp Val Asn Ala Val Ala Phe
 305 310 315
 Ala Asp Ile Ser Ser Gln Ile Leu Phe Ser Gly Gly Asp Asp Ala
 320 325 330
 Ile Cys Lys Val Trp Asp Arg Arg Thr Met Arg Glu Asp Asp Pro
 335 340 345
 Lys Pro Val Gly Ala Leu Ala Gly His Gln Asp Gly Ile Thr Phe
 350 355 360
 Ile Asp Ser Lys Gly Asp Ala Arg Tyr Leu Ile Ser Asn Ser Lys
 365 370 375
 Asp Gln Thr Ile Lys Leu Trp Asp Ile Arg Arg Phe Ser Ser Arg
 380 385 390
 Glu Gly Met Glu Ala Ser Arg Gln Ala Ala Thr Gln Gln Asn Trp
 395 400 405
 Asp Tyr Arg Trp Gln Gln Val Pro Lys Lys Gly Phe Thr Leu His
 410 415 420
 Pro Tyr Pro Ala Trp Arg Lys Leu Lys Leu Pro Gly Asp Ser Ser
 425 430 435
 Leu Met Thr Tyr Arg Gly His Gly Val Leu His Thr Leu Ile Arg
 440 445 450
 Cys Arg Phe Ser Pro Ile His Ser Thr Gly Gln Gln Phe Ile Tyr
 455 460 465
 Ser Gly Cys Ser Thr Gly Lys Val Val Tyr Asp Leu Leu Ser
 470 475 480
 Gly His Ile Val Lys Lys Leu Thr Asn His Lys Ala Cys Val Arg
 485 490 495
 Asp Val Ser Trp His Pro Phe Glu Glu Lys Ile Val Ser Ser Ser
 500 505 510
 Trp Asp Gly Asn Leu Arg Leu Trp Gln Tyr Arg Gln Ala Glu Tyr
 515 520 525
 Phe Gln Asp Asp Met Pro Glu Ser Glu Glu Cys Ala Ser Ala Pro
 530 535 540
 Ala Pro Val Pro Gln Ser Ser Thr Pro Phe Ser Ser Pro Gln
 545 550

<210> 25

<211> 434

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2226116CD1

<400> 25

Met Arg Pro Ser Ser Ser Val Ser Val Ser Cys Pro Ala Leu Asn
 1 5 10 15
 Gln Val Ser His Phe Ala Asn Leu Thr Ser Val Gly Ala Met Ala
 20 25 30
 Pro Ala Arg Cys Phe Ser Ala Arg Leu Arg Thr Val Phe Gln Gly
 35 40 45
 Val Gly His Trp Ala Leu Ser Thr Trp Ala Gly Leu Lys Pro Ser

Arg Leu Leu Pro Gln Arg Ala Ser Pro Arg Leu Leu Ser Val Gly	50	55	60
65	70	75	
Arg Ala Asp Leu Ala Lys His Gln Glu Leu Pro Gly Lys Lys Leu	80	85	90
Leu Ser Glu Lys Lys Leu Lys Arg Tyr Phe Val Asp Tyr Arg Arg	95	100	105
Val Leu Val Cys Gly Gly Asn Gly Gly Ala Gly Ala Ser Cys Phe	110	115	120
His Ser Glu Pro Arg Lys Glu Phe Gly Gly Pro Asp Gly Gly Asp	125	130	135
Gly Gly Asn Gly Gly His Val Ile Leu Arg Val Asp Gln Gln Val	140	145	150
Lys Ser Leu Ser Ser Val Leu Ser Arg Tyr Gln Gly Phe Ser Gly	155	160	165
Glu Asp Gly Gly Ser Lys Asn Cys Phe Gly Arg Ser Gly Ala Val	170	175	180
Leu Tyr Ile Arg Val Pro Val Gly Thr Leu Val Lys Glu Gly Gly	185	190	195
Arg Val Val Ala Asp Leu Ser Cys Val Gly Asp Glu Tyr Ile Ala	200	205	210
Ala Leu Gly Gly Ala Gly Gly Lys Gly Asn Arg Phe Phe Leu Ala	215	220	225
Asn Asn Asn Arg Ala Pro Val Thr Cys Thr Pro Gly Gln Pro Gly	230	235	240
Gln Gln Arg Val Leu His Leu Glu Leu Lys Thr Val Ala His Ala	245	250	255
Gly Met Val Gly Phe Pro Asn Ala Gly Lys Ser Ser Leu Leu Arg	260	265	270
Ala Ile Ser Asn Ala Arg Pro Ala Val Ala Ser Tyr Pro Phe Thr	275	280	285
Thr Leu Lys Pro His Val Gly Ile Val His Tyr Glu Gly His Leu	290	295	300
Gln Ile Ala Val Ala Asp Ile Pro Gly Ile Ile Arg Gly Ala His	305	310	315
Gln Asn Arg Gly Leu Gly Ser Ala Phe Leu Arg His Ile Glu Arg	320	325	330
Cys Arg Phe Leu Leu Phe Val Val Asp Leu Ser Gln Pro Glu Pro	335	340	345
Trp Thr Gln Val Asp Asp Leu Lys Tyr Glu Leu Glu Met Tyr Glu	350	355	360
Lys Gly Leu Ser Ala Arg Pro His Ala Ile Val Ala Asn Lys Ile	365	370	375
Asp Leu Pro Glu Ala Gln Ala Asn Leu Ser Gln Leu Arg Asp His	380	385	390
Leu Gly Gln Glu Val Ile Val Leu Ser Ala Leu Thr Gly Glu Asn	395	400	405
Leu Glu Gln Leu Leu His Leu Lys Val Leu Tyr Asp Ala Tyr	410	415	420
Ala Glu Ala Glu Leu Gly Gln Gly Arg Gln Pro Leu Arg Trp	425	430	

<210> 26

<211> 826

<212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2504472CD1

<400> 26
 Met Val Ala Pro Val Leu Glu Thr Ser His Val Phe Cys Cys Pro
 1 5 10 15
 Asn Arg Val Arg Gly Val Leu Asn Trp Ser Ser Gly Pro Arg Gly
 20 25 30
 Leu Leu Ala Phe Gly Thr Ser Cys Ser Val Val Leu Tyr Asp Pro
 35 40 45
 Leu Lys Arg Val Val Val Thr Asn Leu Asn Gly His Thr Ala Arg
 50 55 60
 Val Asn Cys Ile Gln Trp Ile Cys Lys Gln Asp Gly Ser Pro Ser
 65 70 75
 Thr Glu Leu Val Ser Gly Gly Ser Asp Asn Gln Val Ile His Trp
 80 85 90
 Glu Ile Glu Asp Asn Gln Leu Leu Lys Ala Val His Leu Gln Gly
 95 100 105
 His Glu Gly Pro Val Tyr Ala Val His Ala Val Tyr Gln Arg Arg
 110 115 120
 Thr Ser Asp Pro Ala Leu Cys Thr Leu Ile Val Ser Ala Ala Ala
 125 130 135
 Asp Ser Ala Val Arg Leu Trp Ser Lys Lys Gly Pro Glu Val Met
 140 145 150
 Cys Leu Gln Thr Leu Asn Phe Gly Asn Gly Phe Ala Leu Ala Leu
 155 160 165
 Cys Leu Ser Phe Leu Pro Asn Thr Asp Val Pro Ile Leu Ala Cys
 170 175 180
 Gly Asn Asp Asp Cys Arg Ile His Ile Phe Ala Gln Gln Asn Asp
 185 190 195
 Gln Phe Gln Lys Val Leu Ser Leu Cys Gly His Glu Asp Trp Ile
 200 205 210
 Arg Gly Val Glu Trp Ala Ala Phe Gly Arg Asp Leu Phe Leu Ala
 215 220 225
 Ser Cys Ser Gln Asp Cys Leu Ile Arg Ile Trp Lys Leu Tyr Ile
 230 235 240
 Lys Ser Thr Ser Leu Glu Thr Gln Asp Asp Asp Asn Ile Arg Leu
 245 250 255
 Lys Glu Asn Thr Phe Thr Ile Glu Asn Glu Ser Val Lys Ile Ala
 260 265 270
 Phe Ala Val Thr Leu Glu Thr Val Leu Ala Gly His Glu Asn Trp
 275 280 285
 Val Asn Ala Val His Trp Gln Pro Val Phe Tyr Lys Asp Gly Val
 290 295 300
 Leu Gln Gln Pro Val Arg Leu Leu Ser Ala Ser Met Asp Lys Thr
 305 310 315
 Met Ile Leu Trp Ala Pro Asp Glu Glu Ser Gly Val Trp Leu Glu
 320 325 330
 Gln Val Arg Val Gly Glu Val Gly Gly Asn Thr Leu Gly Phe Tyr
 335 340 345
 Asp Cys Gln Phe Asn Glu Asp Gly Ser Met Ile Ile Ala His Ala

	350		355		360
Phe His Gly Ala	Leu His Leu Trp Lys	Gln Asn Thr Val Asn	Pro		
	365		370		375
Arg Glu Trp Thr	Pro Glu Ile Val Ile	Ser Gly His Phe Asp	Gly		
	380		385		390
Val Gln Asp Leu	Val Trp Asp Pro Glu	Gly Glu Phe Ile Ile	Thr		
	395		400		405
Val Gly Thr Asp	Gln Thr Thr Arg Leu	Phe Ala Pro Trp Lys	Arg		
	410		415		420
Lys Asp Gln Ser	Gln Val Thr Trp His	Glu Ile Ala Arg Pro	Gln		
	425		430		435
Ile His Gly Tyr	Asp Leu Lys Cys Leu	Ala Met Ile Asn Arg	Phe		
	440		445		450
Gln Phe Val Ser	Gly Ala Asp Glu Lys	Val Leu Arg Val Phe	Ser		
	455		460		465
Ala Pro Arg Asn	Phe Val Glu Asn Phe	Cys Ala Ile Thr Gly	Gln		
	470		475		480
Ser Leu Asn His	Val Leu Cys Asn Gln	Asp Ser Asp Leu Pro	Glu		
	485		490		495
Gly Ala Thr Val	Pro Ala Leu Gly Leu	Ser Asn Lys Ala Val	Phe		
	500		505		510
Gln Gly Asp Ile	Ala Ser Gln Pro Ser	Asp Glu Glu Glu Leu	Leu		
	515		520		525
Thr Ser Thr Gly	Phe Glu Tyr Gln Gln	Val Ala Phe Gln Pro	Ser		
	530		535		540
Ile Leu Thr Glu	Pro Pro Thr Glu Asp	His Leu Leu Gln Asn	Thr		
	545		550		555
Leu Trp Pro Glu	Val Gln Lys Leu Tyr	Gly His Gly Tyr Glu	Ile		
	560		565		570
Phe Cys Val Thr	Cys Asn Ser Ser Lys	Thr Leu Leu Ala Ser	Ala		
	575		580		585
Cys Lys Ala Ala	Lys Lys Glu His Ala	Ala Ile Ile Leu Trp	Asn		
	590		595		600
Thr Thr Ser Trp	Lys Gln Val Gln Asn	Leu Val Phe His Ser	Leu		
	605		610		615
Thr Val Thr Gln	Met Ala Phe Ser Pro	Asn Glu Lys Phe Leu	Leu		
	620		625		630
Ala Val Ser Arg	Asp Arg Thr Trp Ser	Leu Trp Lys Lys Gln	Asp		
	635		640		645
Thr Ile Ser Pro	Glu Phe Glu Pro Val	Phe Ser Leu Phe Ala	Phe		
	650		655		660
Thr Asn Lys Ile	Thr Ser Val His Ser	Arg Ile Ile Trp Ser	Cys		
	665		670		675
Asp Trp Ser Pro	Asp Ser Lys Tyr Phe	Phe Thr Gly Ser Arg	Asp		
	680		685		690
Lys Lys Val Val	Val Trp Gly Glu Cys	Asp Ser Thr Asp Asp	Cys		
	695		700		705
Ile Glu His Asn	Ile Gly Pro Cys Ser	Ser Val Leu Asp Val	Gly		
	710		715		720
Gly Ala Val Thr	Ala Val Ser Val Cys	Pro Val Leu His Pro	Ser		
	725		730		735
Gln Arg Tyr Val	Val Ala Val Gly Leu	Glu Cys Gly Lys Ile	Cys		
	740		745		750
Leu Tyr Thr Trp	Lys Lys Thr Asp Gln	Val Pro Glu Ile Asn	Asp		
	755		760		765


```

Trp Thr His Cys Val Glu Thr Ser Gln Ser Gln Ser His Thr Leu
770 775 780
Ala Ile Arg Lys Leu Cys Trp Lys Asn Cys Ser Gly Lys Thr Glu
785 790 795
Gln Lys Glu Ala Glu Gly Ala Glu Trp Leu His Phe Ala Ser Cys
800 805 810
Gly Glu Asp His Thr Val Lys Ile His Arg Val Asn Lys Cys Ala
815 820 825
Leu

```

<210> 27

<211> 618

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3029920CD1

<400> 27

```

Met Lys Lys Asp Val Arg Ile Leu Leu Val Gly Glu Pro Arg Val
1 5 10 15
Gly Lys Thr Ser Leu Ile Met Ser Leu Val Ser Glu Glu Phe Pro
20 25 30
Glu Glu Val Pro Pro Arg Ala Glu Glu Ile Thr Ile Pro Ala Asp
35 40 45
Val Thr Pro Glu Arg Val Pro Thr His Ile Val Asp Tyr Ser Glu
50 55 60
Ala Glu Gln Ser Ser Asp Glu Gln Leu His Gln Glu Ile Ser Gln Ala
65 70 75
Asn Val Ile Cys Ile Val Tyr Ala Val Asn Asn Lys His Ser Ile
80 85 90
Asp Lys Val Thr Ser Arg Trp Ile Pro Leu Ile Asn Glu Arg Thr
95 100 105
Asp Lys Asp Ser Arg Leu Pro Leu Ile Leu Val Gly Asn Lys Ser
110 115 120
Asp Leu Val Glu Tyr Ser Ser Met Glu Thr Ile Leu Pro Ile Met
125 130 135
Asn Gln Tyr Thr Glu Ile Glu Thr Cys Val Glu Cys Ser Ala Lys
140 145 150
Asn Leu Lys Asn Ile Ser Glu Leu Phe Tyr Tyr Ala Gln Lys Ala
155 160 165
Val Leu His Pro Thr Gly Pro Leu Tyr Cys Pro Glu Glu Lys Glu
170 175 180
Met Lys Pro Ala Cys Ile Lys Ala Leu Thr Arg Ile Phe Lys Ile
185 190 195
Ser Asp Gln Asp Asn Asp Gly Thr Leu Asn Asp Ala Glu Leu Asn
200 205 210
Phe Phe Gln Arg Ile Cys Phe Asn Thr Pro Leu Ala Pro Gln Ala
215 220 225
Leu Glu Asp Val Lys Asn Val Val Arg Lys His Ile Ser Asp Gly
230 235 240
Val Ala Asp Ser Gly Leu Thr Leu Lys Gly Phe Leu Phe Leu His
245 250 255

```

Thr Leu Phe Ile Gln Arg Gly Arg His Glu Thr Thr Trp Thr Val
 260 265
 Leu Arg Arg Phe Gly Tyr Asp Asp Asp Leu Asp Leu Thr Pro Glu
 275 280
 Tyr Leu Phe Pro Leu Leu Lys Ile Pro Pro Asp Cys Thr Thr Glu
 290 295
 Leu Asn His His Ala Tyr Leu Phe Leu Gln Ser Thr Phe Asp Lys
 305 310
 His Asp Leu Asp Arg Asp Cys Ala Leu Ser Pro Asp Glu Leu Lys
 320 325
 Asp Leu Phe Lys Val Phe Pro Tyr Ile Pro Trp Gly Pro Asp Val
 335 340
 Asn Asn Thr Val Cys Thr Asn Glu Arg Gly Trp Ile Thr Tyr Gln
 350 355
 Gly Phe Leu Ser Gln Trp Thr Leu Thr Thr Tyr Leu Asp Val Gln
 365 370
 Arg Cys Leu Glu Tyr Leu Gly Tyr Leu Gly Tyr Ser Ile Leu Thr
 380 385
 Glu Gln Glu Ser Gln Ala Ser Ala Val Thr Val Thr Arg Asp Lys
 395 400
 Lys Ile Asp Leu Gln Lys Lys Gln Thr Gln Arg Asn Val Phe Arg
 410 415
 Cys Asn Val Ile Gly Val Lys Asn Cys Gly Lys Ser Gly Val Leu
 425 430
 Gln Ala Leu Leu Gly Arg Asn Leu Met Arg Gln Lys Lys Ile Arg
 440 445
 Glu Asp His Lys Ser Tyr Tyr Ala Ile Asn Thr Val Tyr Val Tyr
 455 460
 Gly Gln Glu Lys Tyr Leu Leu Leu His Asp Ile Ser Glu Ser Glu
 470 475
 Phe Leu Thr Glu Ala Glu Ile Ile Cys Asp Val Val Cys Leu Val
 485 490
 Tyr Asp Val Ser Asn Pro Lys Ser Phe Glu Tyr Cys Ala Arg Ile
 500 505
 Phe Lys Gln His Phe Met Asp Ser Arg Ile Pro Cys Leu Ile Val
 515 520
 Ala Ala Lys Ser Asp Leu His Glu Val Lys Gln Glu Tyr Ser Ile
 530 535
 Ser Pro Thr Asp Phe Cys Arg Lys His Lys Met Pro Pro Pro Gln
 545 550
 Ala Phe Thr Cys Asn Thr Ala Asp Ala Pro Ser Lys Asp Ile Phe
 560 565
 Val Lys Leu Thr Thr Met Ala Met Tyr Pro His Val Thr Gln Ala
 575 580
 Asp Leu Lys Ser Ser Thr Phe Trp Leu Arg Ala Ser Phe Gly Ala
 590 595
 Thr Val Phe Ala Val Leu Gly Phe Ala Met Tyr Lys Ala Leu Leu
 605 610
 Lys Gln Arg

<210> 28
 <211> 596
 <212> PRT
 <213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3332415CD1

<400> 28

Met	Glu	Pro	Glu	Leu	Asp	Ala	Gln	Lys	Gln	Pro	Arg	Pro	Arg	Arg		
1				5					10					15		
Arg	Ser	Arg	Arg	Ala	Ser	Gly	Leu	Ser	Thr	Glu	Gly	Ala	Thr	Gly		
				20					25					30		
Pro	Ser	Ala	Asp	Thr	Ser	Gly	Ser	Glu	Leu	Asp	Gly	Arg	Cys	Ser		
				35					40					45		
Leu	Arg	Arg	Gly	Ser	Ser	Phe	Thr	Phe	Leu	Thr	Pro	Gly	Pro	Asn		
				50					55					60		
Trp	Asp	Phe	Thr	Leu	Lys	Arg	Lys	Arg	Arg	Glu	Lys	Asp	Asp	Asp		
				65					70					75		
Val	Val	Ser	Leu	Ser	Ser	Leu	Asp	Leu	Lys	Glu	Pro	Ser	Asn	Lys		
				80					85					90		
Arg	Val	Arg	Pro	Leu	Ala	Arg	Val	Thr	Ser	Leu	Ala	Asn	Leu	Ile		
				95					100					105		
Ser	Pro	Val	Arg	Asn	Gly	Ala	Val	Arg	Arg	Phe	Gly	Gln	Thr	Ile		
				110					115					120		
Gln	Ser	Phe	Thr	Leu	Arg	Gly	Asp	His	Arg	Ser	Pro	Ala	Ser	Ala		
				125					130					135		
Gln	Lys	Phe	Ser	Ser	Arg	Ser	Thr	Val	Pro	Thr	Pro	Ala	Lys	Arg		
				140					145					150		
Arg	Ser	Ser	Ala	Leu	Trp	Ser	Glu	Met	Leu	Asp	Ile	Thr	Met	Lys		
				155					160					165		
Glu	Ser	Leu	Thr	Thr	Arg	Glu	Ile	Arg	Arg	Gln	Glu	Ala	Ile	Tyr		
				170					175					180		
Glu	Met	Ser	Arg	Gly	Glu	Gln	Asp	Leu	Ile	Glu	Asp	Leu	Lys	Leu		
				185					190					195		
Ala	Arg	Lys	Ala	Tyr	His	Asp	Pro	Met	Leu	Lys	Leu	Ser	Ile	Met		
				200					205					210		
Ser	Glu	Glu	Glu	Leu	Thr	His	Ile	Phe	Gly	Asp	Leu	Asp	Ser	Tyr		
				215					220					225		
Ile	Pro	Leu	His	Glu	Asp	Leu	Leu	Thr	Arg	Ile	Gly	Glu	Ala	Thr		
				230					235					240		
Lys	Pro	Asp	Gly	Thr	Val	Glu	Gln	Ile	Gly	His	Ile	Leu	Val	Ser		
				245					250					255		
Trp	Leu	Pro	Arg	Leu	Asn	Ala	Tyr	Arg	Gly	Tyr	Cys	Ser	Asn	Gln		
				260					265					270		
Leu	Ala	Ala	Lys	Ala	Leu	Leu	Asp	Gln	Lys	Lys	Gln	Asp	Pro	Arg		
				275					280					285		
Val	Gln	Asp	Phe	Leu	Gln	Arg	Cys	Leu	Glu	Ser	Pro	Phe	Ser	Arg		
				290					295					300		
Lys	Leu	Asp	Leu	Trp	Ser	Phe	Leu	Asp	Ile	Pro	Arg	Ser	Arg	Leu		
				305					310					315		
Val	Lys	Tyr	Pro	Leu	Leu	Lys	Glu	Ile	Leu	Lys	His	Thr	Pro			
				320					325					330		
Lys	Glu	His	Pro	Asp	Val	Gln	Leu	Leu	Glu	Asp	Ala	Ile	Leu	Ile		
				335					340					345		
Ile	Gln	Gly	Val	Leu	Ser	Asp	Ile	Asn	Leu	Lys	Lys	Gly	Glu	Ser		
				350					355					360		
Glu	Cys	Gln	Tyr	Tyr	Ile	Asp	Lys	Leu	Glu	Tyr	Leu	Asp	Glu	Lys		
				365					370					375		

Gln Arg Asp Pro Arg Ile Glu Ala Ser Lys Val Leu Leu Cys His
 380 385 390
 Gly Glu Leu Arg Ser Lys Ser Gly His Lys Leu Tyr Ile Phe Leu
 395 400 405
 Phe Gln Asp Ile Leu Val Leu Thr Arg Pro Val Thr Arg Asn Glu
 410 415 420
 Arg His Ser Tyr Gln Val Tyr Arg Gln Pro Ile Pro Val Gln Glu
 425 430 435
 Leu Val Leu Glu Asp Leu Gln Asp Gly Asp Val Arg Met Gly Gly
 440 445 450
 Ser Phe Arg Gly Ala Phe Ser Asn Ser Glu Lys Ala Lys Asn Ile
 455 460 465
 Phe Arg Ile Arg Phe His Asp Pro Ser Pro Ala Gln Ser His Thr
 470 475 480
 Leu Gln Ala Asn Asp Val Phe His Lys Gln Gln Trp Phe Asn Cys
 485 490 495
 Ile Arg Ala Ala Ile Ala Pro Phe Gln Ser Ala Gly Ser Pro Pro
 500 505 510
 Glu Leu Gln Gly Leu Pro Glu Leu His Glu Cys Glu Gly Asn
 515 520 525
 His Pro Ser Ala Arg Lys Leu Thr Ala Gln Arg Arg Ala Ser Thr
 530 535 540
 Val Ser Ser Val Thr Gln Val Glu Val Asp Glu Asn Ala Tyr Arg
 545 550 555
 Cys Gly Ser Gly Met Gln Met Ala Glu Asp Ser Lys Ser Leu Lys
 560 565 570
 Thr His Gln Thr Gln Pro Gly Ile Arg Arg Ala Arg Asp Lys Ala
 575 580 585
 Leu Ser Gly Gly Lys Arg Lys Glu Thr Leu Val
 590 595

<210> 29
 <211> 589
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4031536CD1

<400> 29
 Met Ser Lys Pro Gly Lys Pro Thr Leu Asn His Gly Leu Val Pro
 1 5 10 15
 Val Asp Leu Lys Ser Ala Lys Glu Pro Leu Pro His Gln Thr Val
 20 25 30
 Met Arg Ile Phe Ser Ile Ser Ile Ile Ala Gln Gly Leu Pro Phe
 35 40 45
 Cys Arg Arg Arg Met Lys Arg Lys Leu Asp His Gly Ser Glu Val
 50 55 60
 Arg Ser Phe Ser Leu Gly Lys Lys Pro Cys Lys Val Ser Glu Tyr
 65 70 75
 Thr Ser Thr Thr Gly Leu Val Pro Cys Ser Ala Thr Pro Thr Thr
 80 85 90
 Phe Gly Asp Leu Arg Ala Ala Asn Gly Gln Gly Gln Gln Arg Arg

	95	100	105
Arg Ile Thr Ser	Val Gln Pro Pro Thr	Gly Leu Gln Glu Trp	Leu
	110	115	120
Lys Met Phe Gln	Ser Trp Ser Gly Pro	Glu Lys Leu Leu Ala	Leu
	125	130	135
Asp Glu Leu Ile	Asp Ser Cys Glu Pro Thr	Gln Val Lys His	Met
	140	145	150
Met Gln Val Ile	Glu Pro Gln Phe Gln Arg	Asp Phe Ile Ser	Leu
	155	160	165
Leu Pro Lys Glu	Leu Ala Leu Tyr Val Leu	Ser Phe Leu Glu	Pro
	170	175	180
Lys Asp Leu Leu	Gln Ala Ala Gln Thr Cys	Arg Tyr Trp Arg	Ile
	185	190	195
Leu Ala Glu Asp	Asn Leu Leu Trp Arg Glu	Lys Cys Lys Glu	Glu
	200	205	210
Gly Ile Asp Glu	Pro Leu His Ile Lys Arg	Arg Lys Val Ile	Lys
	215	220	225
Pro Gly Phe Ile	His Ser Pro Trp Lys Ser	Ala Tyr Ile Arg	Gln
	230	235	240
His Arg Ile Asp	Thr Asn Trp Arg Arg Gly	Glu Leu Lys Ser	Pro
	245	250	255
Lys Val Leu Lys	Gly His Asp Asp His Val	Ile Thr Cys Leu	Gln
	260	265	270
Phe Cys Gly Asn	Arg Ile Val Ser Gly Ser	Asp Asp Asn Thr	Leu
	275	280	285
Lys Val Trp Ser	Ala Val Thr Gly Lys Cys	Leu Arg Thr Leu	Val
	290	295	300
Gly His Thr Gly	Gly Val Trp Ser Ser Gln	Met Arg Asp Asn	Ile
	305	310	315
Ile Ile Ser Gly	Ser Thr Asp Arg Thr Leu	Lys Val Trp Asn	Ala
	320	325	330
Glu Thr Gly Glu	Cys Ile His Thr Leu Tyr	Gly His Thr Ser	Thr
	335	340	345
Val Arg Cys Met	His Leu His Glu Lys Arg	Val Val Ser Gly	Ser
	350	355	360
Arg Asp Ala Thr	Leu Arg Val Trp Asp Ile	Glu Thr Gly Gln	Cys
	365	370	375
Leu His Val Leu	Met Gly His Val Ala Ala	Val Arg Cys Val	Gln
	380	385	390
Tyr Asp Gly Arg	Arg Val Val Ser Gly Ala	Tyr Asp Phe Met	Val
	395	400	405
Lys Val Trp Asp	Pro Glu Thr Glu Thr Cys	Leu His Thr Leu	Gln
	410	415	420
Gly His Thr Asn	Arg Val Tyr Ser Leu Gln	Phe Asp Gly Ile	His
	425	430	435
Val Val Ser Gly	Ser Leu Asp Thr Ser Ile	Arg Val Trp Asp	Val
	440	445	450
Glu Thr Gly Asn	Cys Ile His Thr Leu Thr	Gly His Gln Ser	Leu
	455	460	465
Thr Ser Gly Met	Glu Leu Lys Asp Asn Ile	Leu Val Ser Gly	Asn
	470	475	480
Ala Asp Ser Thr	Val Lys Ile Trp Asp Ile	Lys Thr Gly Gln	Cys
	485	490	495
Leu Gln Thr Leu	Gln Gly Pro Asn Lys His	Gln Ser Ala Val	Thr
	500	505	510

Cys Leu Gln Phe Asn Lys Asn Phe Val Ile Thr Ser Ser Asp Asp
 515 520
 Gly Thr Val Lys Leu Trp Asp Leu Lys Thr Gly Glu Phe Ile Arg
 530 535 540
 Asn Leu Val Thr Leu Glu Ser Gly Gly Ser Gly Gly Val Val Trp
 545 550 555
 Arg Ile Arg Ala Ser Asn Thr Lys Leu Val Cys Ala Val Gly Ser
 560 565 570
 Arg Asn Gly Thr Glu Glu Thr Lys Leu Leu Val Leu Asp Phe Asp
 575 580 585
 Val Asp Met Lys

<210> 30

<211> 3375

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 708398CB1

<400> 30

gggggagaagg gagccgcag gatcagggggt cagagttagg gggcttcctt cctgtctcac 60
 ctcacatctca gggccgccaa ctccagctg cagcggcgag ttccagtttc atttccacg 120
 acctctctgc ctgggcccgc agccgcgcgc gcgatgccca gtaagtccag ctgcggcgag 180
 ctccgggagg cgggccagtg ttccgagagt ttcttggtcg ttccgggact ggacatggag 240
 acagatcgcg agcggctcg gaccatttat aaccgcgact tcaagatcag ctttgggacc 300
 cccgccctgt gcttctcttc catgctgtat ggaatgaaga ttgcaaatct ggcctacgtc 360
 accaagactc gggtcagggtt ctccagactc gaccgctggg ccgacgtgcg gttccacgaa 420
 aagaggagaa tgaagctggg gtcagatata agcaaacacc caagtcact gctagccaag 480
 atctttttat acagggctga gtattctcat gggaacatcg gtgtggatgt ggaagtccag 540
 gggcccccag aagcccgaga tgggcagctc ctattccgcc tggatttgaa ccgcaaaag 600
 gtgtgtgacc tgaggcttcg gaatggcgga acccagctcg ttacctctac tcacctcttc 660
 ccactctgcc ggacacccca gtttgctttc tacaatgaag accaggagtt gccctgtcca 720
 ctggggcccc gtgaatgcta tgaactccat gtccattgta agaccagctt tgtgggctac 780
 ttccccagca cagtctctcg ggagctgctg ggacctgggg agtcgggttc agaaggagcc 840
 ggacacattct acattgcccg ctctctgggt cccctcgccc acagccccct ggctgcacag 900
 ctgaagccca tgaactccct caagcggacc cggatcaccc gaaacctgtt ggttcacagt 960
 cggatagagg aaggagagag acctgaccgc gctaagggct atgacctgga gtttaagtat 1020
 cgctgtggga cactactacc acctccccgc ctacaggcag tgcctcccat gcttcttcag 1080
 ggaacaagta tcttctactgc ccttaaggag atcgacagaa tcaaggccca gctggagaca 1140
 gccctgaagt ggaggaaacta tgaggtgaag ctgcggtcg tgcctgcacct ggaggaaact 1200
 cagatggagc atgatatacg gcaactatgac ctggagtcgg tgccatgac ctgggacctt 1260
 gtggaccaga accccaggct gctcacgctg gagggttcct gagtgaactg gagccgcccc 1320
 tcagtgctac gggggcgacca cctgtttgcc ctlttgtctt cggagacaca ccaggaggac 1380
 cccatcacat ataaggcctt tgtgcacaag gtggaaattg accgtgtcaa gctgagcttt 1440
 cccatgagcc tctgtggat ctgttggaat gggctgacct tcaaggtgaa ctttaccttc 1500
 aaccgccagc cgctcgaggt ccagcacctg gccctggagc tgacaggggc ctggctgctg 1560
 ttggcccatg tcttctctgt ggcacctcgg gacgtcccgc agctgcacct agatgtgaaa 1620
 ctcaagctgt acgacggcag tctggagtca aaccacagag agctgcaggc catgaggcac 1680
 atgtttacg gcacaccccg tccagccccc tacatcatct ttgggcccc aggcacccgg 1740
 aagacttgga cgttagtgga ggcatttaag cagggtgtgt agcactttgc caagccccac 1800
 atcttggtct gcgtccatc caactcaggg gctgacctac tctgtcaaa gctccgggtc 1860
 caccctcta gctccatcta ccgctcctg gcccccagca gggacatccg catggtacct 1920

gaggacatca agccctgctg caactgggac gcaaagaagg gggagtagtg atttcccgcc 1980
 aagaagaagc tgcaggaata ccgggtctta attacacccc tcatcactgc cggcaggttg 2040
 gtctcggccc agtttcccat tgatcacttc acacacatct tcatcgatga ggctggccac 2100
 tgcactggagc ctgagagtct ggtagctata gcagggtctga tggaaagtaa ggaacagggt 2160
 gatccaggag ggcagctggt gctggcagga gacccctcggc agctggggcc tgtgctgcgt 2220
 tcccactctga cccagaagca tggactggga tactcactgc tggagcgctg gctcatctac 2280
 aactccctgt acaagaaggg cctctgatggc tatgaccccc agttcataac caagctgctc 2340
 cgcaactaca ggtctcatcc caccatcctg gacattccta accagctcta ttatgaaggg 2400
 gagctgcagg cctgtgcctga tgtcgtggat cgagaacgct tctgcccgtg tctgagcccta 2460
 cctcgacagg gctttcccat catctttcac ggcgtaatgg gcaaaagatga gctggaaggc 2520
 aacagcccat cctctctcaa cctgaagag gctgcacacag tgaactccta cctgaagctg 2580
 ctctcgccct cctcctccaa gaaggggcaaa gctcgcttga gccctcgaag tgtggggctc 2640
 atctccccgt accggaataa ggtggagaaa atccgttact gcatcaccaa acctgacagg 2700
 gagctctcag gactggatga catcaaggac ttgaaggctgg gttcagtaga agaattccaa 2760
 ggccaagaac gaagcgtcat cctcatctcc accgtgcgaa gcagccagag ctttctgcag 2820
 ctggatctgg actttaatct ggggtttcctt aagaacccca agaggttcaa tgtagctgtg 2880
 acccgggcca aggcctctgt catcatcgtg gggaaaccccc tctcctcggg catggaacct 2940
 gactggaag gattctctga gttctgtaaa gaaaacggag ggtataccgg gtgtcccttc 3000
 cctgcctaaa tggacctgca acagggacag aatttactgc aagggtctag caagctcagc 3060
 cccctctact caggggcccca cagccatgac tacctcccc agggagcggga ggggtgaagg 3120
 ggctctgtct tgcgaagtgc gccagagtgg aggaatgagc tctgaagaca cagcagccag 3180
 cctctctcga ccagccaagc cttaactgcc tgccctgacc tgaaccagaa cccagctgaa 3240
 ctgccctctc aaggacacag aaggctgggg gaggagggtt acaacccaa ccatccaac 3300
 cctctccctg ctggggagaa tgacacatca agctgctaac aattggggga aggggaagga 3360
 agaaaactct gaaac 3375

<210> 31

<211> 2434

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1259937CB1

<400> 31

caaggtatccg atgggtatat ggagtgtgag gtaatggatc attcatgtgg aaggatgcag 60
 ggggtttttg agaccagggt ttggaagaga gttcagcact gctggtagtt ttgggaataca 120
 cccatgtgca ggcgacacat gaggcagtaa ggaactctgc aggggtccct gagatttggga 180
 aatgttagga agagcaatgg attgaggtcc gaacctggag gatctgctat acgcagagct 240
 ggggagaggg acagagtcag taccagagtc ggaataaagg ggggtgggaa ggggaacctg 300
 agtcaggaga cttgcctggc aggcgctgcc ctgccagcag aggcctgaca gtgggtttcca 360
 tgaactgcac cctcgtctgt ggctgggaca gggccactga cacagtatcg gagcacagaa 420
 ggggaaagga gcagagggga ttccaaactct gccagttaga agctgtgtgg ctttgggcat 480
 gtctcttaac cctctcgagc ctcaattatt tcatccataa aaataataac 540
 ttttgtcaaa ggcgcatgtt gaattatttag atcctcagaa taatgccttg cttgtagcaa 600
 atgttagctg gaggaaaagg aagagaaaaa agctgcagc agctgaagga ttttcatatt 660
 agaactgtct tggacctatc tggcagatgc agaagcacac acacacggag gggcatggat 720
 ttgcccgcgc cttagacatg ttgtgtcttc tectggatcc ttgggtccag gtgcctttacc 780
 tgagctcagg tgaagtgtgc aagcagagcc ctctggtgtg ttggaatgct ttgtggcgccc 840
 gtgctcctgg tgacacaggg acctcacaat cctccctccc acggtctcct ctcatgtcct 900
 cccagctcta tttctcgtt cctcttctcc ccaggcccgag aactgtccct ttgtggtccc 960
 caaccaggac gagccccctc ctggcagcag ctgtgccatc caagttgggg ataaagctcc 1020
 ctatgacato tgcggccag accactcagt gttgacacct cagctgcctg tgacagctc 1080

```

cgtgagagag gtgatggcag cgttggccca ggaggatggc tggaccaagg ggcaggtgct 1140
ggtgaagggtc aattctgcag gtgatgccat tggcctgcag ccagatgccc ggtgtgtggc 1200
cacatctctgt gggctcaatg agcgtctctt tgttgtcaac ccacaggaaag tgcattgagct 1260
gatcccaacac cctgaccagc tggggccccc tgtgggtctct gctgagggggc tggacctggt 1320
gagtgcccaag gacctggcag gccagctgac ggaccacgac tggagcctct tcaacagtat 1380
ccaccagggtg gactgtatcc actatgtgct gggcccccag catctgcggg atgtcaccac 1440
cgccaacctg gagcgtctca tgcgcgcgtt caatgagctg cagtactggg tggccaccga 1500
gctgtgtctc tgcctcgtgc ccggcccccg gggccagctg ctcaggaaagt tcattaagct 1560
ggcggcccat ccaaggagc agaagaatct caattccttc tttgccgtca tgtttggcct 1620
cagcaactcg gccatcagcc gcttagccca cacttgggag cggctgcctc acaaagtccg 1680
gaagctgtac tccgcctctg agaggctgct ggatccctca tggaaaccac ggggtataccg 1740
actggccctc gccagctctt cccctcctgt catccctctc atgccccctc ttctcaaaaga 1800
catgaccttc attcatgagg gaaaccacac actagtggag aatctcatca actttgagaa 1860
gatgagaatg atggccagag ccgcgcggat gctgcaccac tggcgaagcc acaacctgtg 1920
gcctctctca ccactcagaa gccgagtttc ccacctccac gaggacagcc aggtggcgag 1980
gatttccaca tgcctggagc agtccctgag caccggaggt ccagccagca cctgggctta 2040
tgtccagcag ctgaaggatca ttgacaacca gggggaactc tccgcctctc cccgagagct 2100
ggagccatga ggagggggctg ggactggagc tggagcaggc acttgacgct ggggaaagcca 2160
gggtgtggcg ggcccaagata ctccacaggct ggccacagct gggcgaagct ctccgtggag 2220
tggactcgag tctctggagc aggcagtggt gaggcagcca tccctgtgta tgactggagc 2280
ctaaggagga ccctggagtg gacccgagcc aggaataacg aatgacccaa ggccaaggaa 2340
gggaggacag agaggcccca ggagtgggtg gagagtggag tgcgctggga cgttgtgtgc 2400
aatagagagg tctccacacc agaaaaaaa aaaa 2434

```

<210> 32
 <211> 892
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1452285CB1

```

<400> 32
cagcctcttc tctgcacctc caattgcgct ccccaagtct ctctcgtgcy cagagccagc 60
gctgcgcttc cctggtcagc cagggcacgt ctggcccgcc gccaggatgc aggcccccga 120
caaggagcac ctgtacaagt tgctgtgat tggcgacctg ggcgtgggga agaccagtat 180
catcaagcgc tacgtgcacc agaacttctc ctgcactac cgggcccaca tcggcctgga 240
cttcgcgctc aaggtgctcc actgggagcc ggagactgtg gtgcgcctgc agctctggga 300
tatcgacagt caagaaagat ttggaaacat gacgagggtc tattaccgag aagctatggg 360
tgcatattat gtcttcgatg tcaccaggcc agccacattt gaagcagtgg caaagtggaa 420
aaatgatttg gactccaagt taagtctccc taattggcaa ccggtttcag tggttttgtt 480
ggccaacaaa tgtgcacagg ggaaggatgt gctcatgaac aatggcctca agatggacca 540
gttctgcaag gagcacggtt tcgtagggatg gtttgaaaca tcagcaaaagg aaaataaaa 600
cattgatgaa gcctccagat gccctggtgaa acacatactt gcaaatagst gtgacctaat 660
ggagtctatt gagcggagcy tcgtgaagcc ccactctaca gcaaccaagg ttgccagctg 720
ctctgctgtg gccaaatctc agtaggcacc ttgtctgggt tctggttaga atgacctcat 780
tgttccacaa attgtgcctc tatttttacc attttgggta aacgtcagga tagagatacc 840
acatgtggca agccaaagat ctatgcctcc atatgtgctt tctctgttagc tt 892

```

<210> 33

<211> 2288
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1812894CB1

<400> 33
 ccagatggag gcaacagtag taggttgac gacttgctaa gccggtaggt 60
 ggctctgggt aagagatggg aagagaggct gatccctgtc cccacacagca ctggaggagct 120
 ccgggtccca ggaaggggca aatggtgagg ccgagtcagg tgagctcagg ccgcttcggg 180
 ctgagctggc agggcgccctg gcagaaatgg aaaccatgaa ggctgtggga gaggtgagcg 240
 agagcacgaa ggccgaggct gtggctgcgg tcgacggcca gtgccaagag gaggtggcct 300
 cgctcgaggg catctgaaa gactccatca gcagctatga agcccagatc accgccttga 360
 agcaggagcg acagcagcag cagcaggact gtgaggagaa ggagcgggag ctggggccgc 420
 tgaagcagct gctgtcccg gcctacccc tggactccct ggagaagcag atggaaaaagg 480
 ccacagagga ctcgagagaag ctgcgggaga tcgtactgcc catggaaaag gagatcgagg 540
 agctgaaggc gaagctgctg agggccgagg agctgattca ggagatccag agacgtcccc 600
 ggcatgcccc ttccctgcac ggctccacgg agttgctgcc cctgtcccg gatccatgcg 660
 ccccgctgga gcccttgagg gagctgagcg gagatggggg tccagccgct gaggccctcg 720
 ctcaacaact cgatgacagc gccctccatct cctccttctc ccttgccggg ggggtcgcca 780
 gcagctctct cctgccccaa agccgcaggg gccctgagccc tgaacaggaa gagacggcct 840
 cgctgggtgtc tacggggcacc ctgggtcccg agggcatcta cctgccccct cctggctacc 900
 agctcgtccc agacactcag tgggagcagc tgcagacaga gggccgacag ctgcagaagg 960
 acctggagag cgtcagtcgc gagcgggacg agctccaaga gggcctgaga cggagcaatg 1020
 aggactgtgc caagcagatg cagggtgctcc tggcccagggt ccagaactca gacgagctcg 1080
 tgcggaccct gcaagggacc gtgagccagg cccaggagcg ggtgacagctg catatggcgg 1140
 agctgggtcac caccacaag tgccctgcacc atgaggtaaa gcggttgaat gaggaaaacc 1200
 aagggtcccg ggcggagcag ctgccatcct cagcccccca gggctcgacg caggagcagg 1260
 gcgagagagca atcactgcc accctctgtgc cagagctgca gcagctgctg tctgcagcgc 1320
 ggcaagaggc gagggcccg ctgcaggccc agggagcacgg ggccgagcgc ctgcggatcg 1380
 agatcgtgac gctgcgggag gctctggagg agggagacagt ggccaggggc agcctggagg 1440
 ggcagctgag ggtgcagcgg gagggagacag aggtgctgga ggctccctg tgcagcctga 1500
 ggacagagat ggagcgggtg cagcaggaaac agagcaaggg ccagctccca gacctcctct 1560
 cagaacagag gagccaagtg ctgcggctgc aggcagagct ggagaccagt gacgaggtgc 1620
 agagggattt cgtgcgactg tcccaggccc tgcagggtgc cctagagcgg atccgccagg 1680
 ctgagacccg gtagcaagtg cgcagcatca tggatgaggg ccactcacg gacgtcaggg 1740
 acatcaagga cactgaggg gtcaggatat cccaccccc accctgggaa agacgccttt 1800
 cccactcctt gaaacatgag gccctgctct ggggtcttgg atggcttttc caccgctcct 1860
 gagactgggg ttgaggggac tgacgggggc caccacccgc cgcctccctc gcgcctcctc 1920
 ccagggtgct tgggcctcct gttctcaggg atcacacctg ggtgaggggc ccaagccctc 1980
 ccgggaacca aaggtgcagg ctacggcctg cggctttctg gctcgtgtgc tgctcctgg 2040
 gctccagccc tccctgccc ccagcccgtc cctgcccag ggcacagcgg agccatgggg 2100
 gctgggagtc cccatcagc gcagttaggt gggcccggc cctgggacag gcagctgctc 2160
 tctggtctgc atgacactaa gacgctgtgc cacagcggcg acccaggctc ccaagcttgc 2220
 acagaggcaa ggccagactt ttccgtggtt tattttcaat aaataagcag ctcagcgcaa 2280
 aaaaaaaa

<210> 34
 <211> 1813
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> 1708, 1711, 1713, 1715
 <223> Incyte ID No: a o r g o r c o r t, unknown, or other

<220>
 <221> misc_feature
 <223> Incyte ID No: 3074884CB1

<400> 34
 gcacgaagga ggcggcgctt caggtggccc tggcggagat gccggaagat gcggacgaga 60
 acgcgcagga ggaagctctg cggggagagc ctctgctgcc ggccgggacc cagcgctgt 120
 gtctggttca ccttgacgtc aagtggggcc cggggaagtc gcagatgact cgaagccagt 180
 ggcaggtggc ggagggccaca gcgctgggtg acacgcctga cggctggctc gtggtgcaga 240
 caatggtcgt gtccaccaa acgcgggaca ggaagctcat ctttggcaaa gggaactttg 300
 agcacctgac agaaaagatc cgagggtctc cagacgtcac gtgcgtcttc ctgaacgtgg 360
 agaggatggc tgccccgacc aagaaagaac tggaaagccg ctggggcgctg gaggtgtttg 420
 accgcttcac ggtcgctctg cacatcttcc gctgtaacgc ccgcacgaag gaggcccgcc 480
 ttacagtggtc cctggcgagg atgcgctgc acaggtcgaa cttgaaaagg gacgtcgccc 540
 acctgtaccg aggagtcggc tcgcgtaca tcatggggtc aggaataatc ttcatgcagc 600
 tgacacagcg tctcctgaga gagaaggagg ccaagatcag gaaggccttg gacaggcttc 660
 gcaagaagag gcacctgctc cgcggcgagc ggaacgaggc ggaagtcccc gtgatctccg 720
 tgggtggggtt caccactgc ggaagaacca cgctgatcaa ggcactgacg ggcgctcacc 780
 ccattccagcc acgggaccag ctgtttgcca cgctggagct caccggccacc gccggcacgc 840
 tgccctcacg catgacgcgt ctgtacgtgg acaccatcgg ctctctctcc cagctgccc 900
 accgctcatc cagatcttct tccgccacc tggaaagacgt ggcccactcg gatctcatct 960
 tgacagtgag ggaagtcagc caccgcgagg cggagctcca gaaatgcagc gttctgtcca 1020
 cgctcgctgg cctgcagctg cccgccccgc tcttggatc catggtggag gttcacaca 1080
 aggtggacct cgtgcccggg tacagcccca cggaaacgaa cgtcgctgcc gtgctctgcc 1140
 tgcggggcca cgggctccag gagctgaaag ctgagctcga tgcggcggtt ttgaaggcga 1200
 cggggagaca gatcctcact ctccgtgtga ggctcgagc ggcgcagctc agctggctgt 1260
 ataaggagcg cacagttcag gagtggaagc tgatcctga ggaacggcg gcgcagctga 1320
 gggctcatcat cagcaactca gcctacggca aattccggaa gctcttccca ggaatgaacg 1380
 acgcccacag aggcctgcgg ggtgggggca tcgctgcctg gggagctgag gcgttacccg 1440
 tgtgtttggg gcagcttggg gtcaggtgca gcagggtctc ccttgtctgg ttctgacccc 1500
 gtctcgctcc cagccatttg ctgggatgac cgtgcagggc ggtgacacgg ccgcacctgc 1560
 cccaaagcgg gcgccccgag cgtccaactc aagcctgagc atccacacaa ttcagctggg 1620
 ccttcggctg ctgctgtgaa ctgctttccc tcggaatgtt tccgtaacag gacattaaac 1680
 ctctgttttt acttcctgta aaaaaaanac nngnnaaaaa aaaaagggcg gccccgctcc 1740
 tagaggattc caagccttac cgtaacgcgt tgcattggcg agcggctata agctctcttc 1800
 aatagggggc cac 1813

<210> 35
 <211> 1733
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3452277CB1

<400> 35
 ctacggcctg gaccgagtga ccaatccgaa tgaagtcaag gtaaacgaga aacaaacagt 60
 cgttgctgtc aaaaaagaga tcattgtatta ccaacaggcc ttgatgaggt ccacagtga 120

gtcttctgtg tccctgggag ggattgtgaa atacagtggag cagtttctcat ccaacgatgc 180
catcatgtca ggtgtccctcc ccagcaaccc ctggatccacc gatgacaccc agttctctggga 240
cttaaaatgcc aaattgggtg aaatcccaac caagatgcga gtggaacgat gggcctctcaa 300
cttcagcgaa ttgatccgag accccaagg tcgacagagc ttccagtact tctcaagaa 360
agaattcagt ggagagaatc tgggattctg ggaagcctgc gaggatctga agtatggaga 420
tcagtccaaa gtcaaggaga aagcagagga gatttacaag ctgttctctgg ccccgggggc 480
gaggcgctgg atcaacatag atggcaaaac catggacatc acagtgaagg ggctgaagca 540
ccccaccgcg tatgtctgg acgcccacaa aaccacatt tacatgtcca tgaagaagga 600
ttcttatgtc cgtattttaa aatctccgat ctataaggac atgtggcca aagctattga 660
acctcaggaa accaccaaga aaagctccac cctccctttt atcgcgctc acctgcgctc 720
cagcccaagc cctgtcatcc tgagacagct ggaagaggaa gccaaagccc gagaagcagc 780
caacactgtg gacatcaccc agccgggcca gcacatggct ccagccccc acttgacctg 840
gtacacgggg acctgcatgc ccccgctctc ttctagcccc ttctctctct cctgcgctc 900
ccccaggaa cctttcgccct caccagcgcg ctctatccgg cgacccagca caccatctg 960
ccccctaccc atcagagtgg ccttggagag ctcatcgggc ttggagcaga aaggggagtg 1020
cagcgggtgc atgcgcccc gtgggccttc tgtcacccag agcagcgagc cctccctcga 1080
cacctctggt cctgcagccc gccccagggc cccctcctaag gcccgcatgg cctctgctct 1140
cagcaggttt ctgagacagc gctgtctggc ctacactgtc ttggccaggc tctcacccaa 1200
gtgcccctgt gtgtcccaag ggagggtgca gcccttgggg gacgtggggc agcagctggc 1260
acgattgaaa tccaagagag tagcaaaact ttccagatc aaatggatg tggccacggg 1320
gagcgggac tgcttgatgg actcggagga tgcctggaaca ggagagtcgg gtgaccgggc 1380
cacagaaaag gaggctcatc gcccttggga gagcctgtaa ggaagagggc aggctgagct 1440
gggggctctg gaccaggaa atgctctgac agatgccatg gtatgggcca caggacacac 1500
tgctcgaga accaaagtgc atttgggtga catttgaaga ttggggagac aagatggggg 1560
agatttggcg aagaaatgct ctggctgggt accagggggc aaactctctt cctctctctg 1620
accctccctc ccttggcgag aagaaacgca tgtggaccag agactcttct ctgtgctctt 1680
aaaaccaata aaaggttaac tttaagtctt ttggaaaaaa aaaaaaaaag ggg 1733

<210> 36

<211> 1776

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<223> Incyte ID No: 4203832CB1

<400> 36

cccagccgag cccagcctag cccagagccca gcccgagcga acccgagacc ccaagcccga 60
gcgcgcgccca gcccgagcag agccctccagc ccgctcacc cgcgtgcacc cccagcgacc 120
ctcagccgct ctctgcctct ctctcgcccc cgcgcgccgc ctgcgcggcc cctctgccaa 180
tgaaactgac cgcgatgatc aagaagatgt gcccgagcga ctcggaagct agtatcccg 240
ccaagaactg ctatcgatg gtcatcctcg gctcgctcca gttgggccaag acggccatcg 300
tgtcgcgctt cctcaccggc cgcctcgagg acgctacac gcctaccatc gaggacttcc 360
accgcaagtt ctactccatc cgcggcgagg tctaccagct ccacatcctc gacacgtctc 420
gcaaccaccc gttccccgcg atcggtgtgc tctccatcct cacaggagac gttttcatcc 480
tggtgttcag tctggacaac cgcgactcct tcgaggaggt gcagcgctc aggcagcaga 540
tctctgcac caagtcttgc ctcaagaaca aaaccaagga gaacgtggag gtgcccctgg 600
tcactctgcy caacaagggt gaccgcgcat tctaccggca ggtggaccag cgcgagatcg 660
agcagctggt gggcgacgac cccagcgct cgcgctact ctgatctctg gccaaagaa 720
acagcagcct ggaccagatg ttccgcgcgc tcttcgccat gcaccaagct cccagcgaga 780
tgagcccgaa cctgcaccgc aaggtctcgg tgcagtact cgacgtgctg cacaagaag 840
cgctcgcgaa caagaagctg ctgcggcgcc gcagcgcgcc cgcgcggcgc gacccggcgc 900
acgcctttgg catcgtggca cctctcgccg gccggccccc cgtacacagc gacctcatgt 960

```

acatccgcga gaagccagc gccgcagcc aggccaaagga caaggagcgc tgcgtcatca 1020
gctaggagcc ccgccgcgct ggccagacaaa cctaaaggagg acctttttgt taagtcaaat 1080
ccaaacggccc ggtgcgcccc aggcggggag cgccgcgcgga ctggcgctctc cccctccggc 1140
gatccgcccc cagcactggg gagcgccac tgaaccgaga agggacggtc atctgctccg 1200
gaaggaaga gaacggggcca agactgggac tattccccac ccccggtccc ccattgaggc 1260
ccgccacccc cataactttg ggagcgaggg cccagccgag ggtggattta tcttctcaaa 1320
gactaaagag tgagcgcggg gctggggagg gatgtgaagt tatccagcct ctgctaggct 1380
tcaagaaccc gtctgcgccg ctgagggttc aggacccacg gggcattatc ttgtctgtga 1440
ttccgggttg ctgtgacgag cggtagagcc tctgcctccc cgaaactaag cggggggggc 1500
tgggtcaaat catagccaa tgacttgttt acatgtgagt gaaactgcac aaagggaacac 1560
aaaaaacaac ttgcacttta acggtagttc cgggtgtcaac atggacacga acaaaacctt 1620
accagggtgt ttactactgt gtgtgtgtga gcttttaaa attttgcttt attttggttt 1680
ttaatatata ataaaaaat ttaaaatgga aaaaaaata aaaaaaaaa aaaaaaaaa 1740
aaaaaaaaa aaaaaaaaaa gaaaaaaaaa aaaaaa 1776

```

```

<210> 37
<211> 1316
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 104368CB1

```

```

<400> 37
cggacgcgtg gggcggttt tctcttttag gctcaggggc ctgaccact gctcttctc 60
ttaagaaggt gctccattcc ttccggcgcc cggagctgct ggcccaaaag gatccggagc 120
gagctagggc agacacccatg accacccttg atgataagtt gctggggagg aaactgcagt 180
actactatag cagcagtgag gatgaggaca gtgaccacga ggacaaggac cgaggcagat 240
gtgccccagc cagcagttct gtgcctgcag aggtgagct ggcaggcgaa ggcattctcag 300
ttaacacagg cccaaaaaggt gtgatcaatg actggcgccg cttcaagcag ttggagacag 360
agcagagggg ggagcagtg cgggagatgg aaaggctgat caagaagctg tcaatgactt 420
gcaggtccca tctggatgaa gaggaggagc aacagaaaca gaaagacctc caggagaaga 480
tcagtgaggaa gatgactctg aaggagtttg ccataatgaa tgaggacca gatgatgaag 540
agtttctgca gcagtaccgg aagcagcgaa tggaaagatg gcggcagcag cttcacaagg 600
ggccccaact caagcaggtt ttgagatct ccagtggaga agggttttta gacatgattg 660
ataaagaaca gaaagcatt gtcatcatgg ttcattttta tgaggatggc attccaggga 720
ccgaagccat gaatggttgc atgatctgcc ttgccgcaga gtaccacgct gtcaagttct 780
gcaaggtgaa gagctcagtt attggcgcca gcagtcagtt caccaggaat gcccttctct 840
ccctgctgat ctataagggg ggtagaatga tcggcaattt ttgtctgtgt actgaccagc 900
tgggggatga ttctcttctg gtggaccttg aagcttttct ccagggaatt ggattactcc 960
cagaaaaagg agtcttggtg ctgacatctg tgcttaactc tgccacgtgt cacagtgaag 1020
atagcgacct ggaatatgat tgaactgata gtctagtgtc atagatttct cattgtttgg 1080
gttggaatac acgtcattgt ttatttttgt tcttttgtct ctgtgtttct cagctgttct 1140
ttgtagtccc ttttatatg cataaaataa agaaattctt agattaaatc agaagtctga 1200
ataactgtta acctagcaat aaggtgactt acagttgtat aaacagggaag ccaggctttt 1260
gaactgttta tctaagattc tgtggtgtga catctctgtt attgtttcca gtcact 1316

```

```

<210> 38
<211> 1554
<212> DNA
<213> Homo sapiens

```

<220>

<221> misc_feature

<223> Incyte ID No: 1441680CB1

<400> 38

```

gtgggttgtg  gagtggcgac  cgtagtgtag  gcggttgctg  agacagacgc  tggagcggtt  60
aggaggagcc  cgagcggtaa  gggaaagccg  gatgaggggc  gtgttgacgt  ggagagataa  120
agccgagcac  tgtataaatg  acatcgcat  taagcctgat  ggaactcaac  tgattttggc  180
tgccggaagc  agtatattcg  tttatgacac  ctctgatggc  accttacttc  agccctctaa  240
gggacacaaa  gacactgtgt  actgtgtggc  atatgcaag  gatggcaagc  gctttgtctc  300
tggatcagct  gacaaaaagc  ttattatctg  gacatcaaaa  ctggaaggca  tttctgaagta  360
cacgcacaat  gatgtctaac  aatgtgtctc  ctacaatcct  attactcttc  aactggcatc  420
ttgttctctc  agtgactttg  ggttgtgtgc  tcttgaacag  aagtctgtct  ccaaacacaa  480
atcaagcagc  aagatcatct  gctgcagctg  gacaaatgat  ggtcagttac  tggcgctggg  540
gatgttcaat  gggatcatca  gcatacggaa  caaaaatggc  gaggagaaa  taaagatcga  600
gcggccgggg  ggctccctct  cgccaatatg  gtccatctgc  tggaaacctt  caagagagga  660
gcggttgctg  attcctggct  tggctgactg  gggacagaaa  gtttccctct  accagctgag  720
acgtaatgac  atcggaagag  atcgggcact  gaactttgac  cctctgtgca  tcagctactt  780
tgaaaaaacg  attggaagag  atcgggcact  gaactttgac  caagtatctc  ttttaccaca  840
tactaaagcg  gagtacattt  tgctgggggg  ttcagacaa  tgggtgtgga  cgtgtcaagc  900
ggatggagtg  cggcttgagg  ctgttgggga  gcagaactcc  gggcaccatt  ccttctacca  960
gaaacccgat  tccaactatg  tgggtgtcgg  ctgccaggac  tatgccatta  gggatagcat  1020
gcttattttc  agcagcatcc  atggagttta  caaggaccgc  tatcggatta  aatgcagaaa  1080
gactgacgtc  atttgcagc  acctgatcac  tgagcagaaa  atcccaactc  cagagaaaaa  1140
gcttgtcaag  aagattgcc  tctacagaaa  tcgattggct  cattaccggg  taaaggagaa  1200
ccctcatct  agtgtgtatt  cagaggactt  atcagacatg  gccaatcaca  tccctctgtg  1260
gattatcaag  aagtttgagt  gcaacctctg  ggtgtgtgt  aaggagcggg  agtggcagat  1320
caggagaaa  cggctgcagt  gctgtcctt  cagcggagtg  ccttggacct  aaggctctct  1380
ggagtctctc  attcgttaca  tcaagtgat  cgggtggccct  ctgggaagag  aaggctctct  1440
agtggggctg  aagaagatgt  acttgttagt  gtattcattc  atattgatg  taaaggatta  1500
tttttcaact  agtactgatg  tcttggaaa  tcttacctgg  aaacatgttt  gcaaaaaaca  1554
ttattgggtc  tttcatcttt  tttcttggt  ttacatattt  gttcaataaa  aata

```

<210> 39

<211> 2320

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1494955CB1

<400> 39

```

gggcgcgcgc  gtactcacta  gctgaggtgg  cagtgggttc  accaactatg  agctctcgca  60
gatgtcggag  ctcatggggc  tgtcgttgtt  gcttgggctg  ctggccctga  tggcgacggc  120
ggcgttagcg  cgggggtggc  tgcgcgcggg  ggaggagagg  agcggccggc  ccgctctcca  180
aaaagcaaat  ggatttccac  ctgacaaatc  ttccggatcc  aagaagcaga  aacaatatca  240
cgggattcgg  aaggagaagc  ctcaacaaca  caacttcacc  cacgcctcc  tggctgcagc  300
tetgaagagc  cacagcggga  acatatcttg  catggacttt  agcagcagtc  gcaaatacct  360
ggctacccgt  gcagatgatc  gcaacctcgc  catctggagc  accaaggact  tctctcagcg  420
agagcacgcg  agcatgagag  ccaacgtgga  gctggaccac  gccacctctg  tgcgtctcag  480
cctgtactcg  agagccttca  tctcttggtc  ggcacacggg  gacacctctc  gtgtcttcaa  540
gatgaccaag  cggggaggtg  ggggctacac  cttcacagcc  accccagagg  acttccctaa  600
aaagcacaag  gcgcctgtca  tcgacattgg  catgtctaac  acaggggaag  ttatcatgac  660
tgctecagtg  gacaccactg  tctcatctg  gagctgaag  ggtcaagtgc  tgtctaccat  720

```

caacaccaac cagatgaaca acacacacgc tgcgttatct cccctgtggca gatttgtagc 780
 ctgcgtgtggc ttaccccagc atgtgaaggt ttgggaagtc tgctttggaa agaaggggga 840
 gttccaggag gtggtgcgag ccttcgaact aaagggccac tccgcggctg tgcactcgct 900
 tgctttctcc aacgactcac ggaggatggc tctctgtccc aaggatggta catggaaact 960
 gtgggacaca gactgggaat acaagaagaa gcaggacccc tactgtctga agacaggccg 1020
 ctttgaagag gcggcggggt cgcgcgcgtg ccgcctggcc ctctccccc acgcccaggt 1080
 cttggccttg gccagtgcca gtagtattca tctctacaat accggcgggg gcgagaagga 1140
 ggagtgcttt gagcggttcc atggcagatg tatcgccaac ttgtcctttg acatcactgg 1200
 ccgctttctg gcctctctg gggacggggc ggtgcggctg ttccacaaca cctctggcca 1260
 ccgagccatg gtggaggaga tgcagggccca cctgaagcgg cctcccaacg agagaccoccg 1320
 ccagaggctg cagcagcagc tgacccaggc ccaagagacc ctgaagagcc tgggtgcccct 1380
 gaagaagtga ctctgggagg gcccgggcca gaggattag gaggagggat ctggcctcct 1440
 catggcgctg ctgccatctt tctctccagg tggaaagcct tcagaaggag tctcctggtt 1500
 ttctactggg tggccctgct tcttccatt gaaactaact ttgtctactt aggtctctct 1560
 ctcttctgct gcctgtgaet cctcctgaat agtggccaag gtgcttttct tctcccgagg 1620
 cccagtggtt ggaatctgtc ccdacctggc actgaggaga atggtagaga ggagaggaga 1680
 gagagagaga atgtgatttt tggcctctgt gcagcacatc ctacacacca aagaagtttg 1740
 taaatgttcc agaacaacct agagaacacc tgagtactaa gcagcagttt tgcaggagtt 1800
 ggagactggg atagcttccc atcacagaac tgtgttccat caaaaagaca ctaagggatt 1860
 tctctctggg cctcagttct atttgtaaga tggagaataa tctctctctg gaactccttg 1920
 caaagatgat atgaggttaa gagaatatca agtccccagg tctggaagaa aagtagaaaa 1980
 gagtagtact atgtccaat gtcataaag tggtaaaagt ggggaaccagt gtgctttgaa 2040
 accaaattag aaacacattc cttgggaagg caaagtttct tgggacttga tcatacattt 2100
 tatatggttg ggaacttctc cttcgggaga tgatatcttg tttaaaggaga cctcttttca 2160
 gttcatcaag ttcatcagat atttgagtgc ccaactctgt cccaaaataa tatgactggt 2220
 ggattaaaat tgaataagac atggttctgt ccatcaaaag tggctggttg gagagagaga 2280
 tacaccotta ttaagtgtct tgtgttagtt tattcatagc 2320

<210> 40

<211> 879

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1508161CB1

<400> 40

gaagaatttg ttcaggcgtt cgtgcagaag gacccttttg ataatgacaa gagttgtctac 60
 agtgaacgga agaaaaacac aaacttagaa gcttacgtgg aatgggttaa tcgctcagc 120
 tactttggtt ctacagaaat ctgtatgcct gttacgaaaa aacaccgagc aagaattgat 180
 gagtatttca ttgacgtagc tcgggagtggt ttttaacattg gcaacttcaa cctcttgatg 240
 gcgataactc ctggtagtaa tatgagccca gtctctcgac taaaaaaaac ttgggccaaa 300
 gtgaagactg caaaatttga cattcttgag catcagatgg acccttcaag caatttctat 360
 aattatcgaa cagctctctg tggggcgaga caaaggtctt taactgtcca tagtagtaga 420
 gaaaagattg tgataccatt cttcagttct ttaactcaag atatttattt cctcaatgag 480
 ggttgtgcca accgccttcc caatggccat gtcaattttg agaattttg ggaactggcc 540
 aaacaagtga ctgaatttat gacatggaaa atcagtgaggt gtccatttga gagggaccgg 600
 aagatctctg agtatctgct cacagatcca gtctctcagtg aagatgtcct ctacttggct 660
 tcttatgaga gtgaaggacc tgaaaaatcat atagagaaag acagatggaa gtctttaagg 720
 tcgagcctct taggcagagt ttaacacatg ggagctgcct gcctgtcgtc gctgtgctt 780
 cctgcatgat atggaggggc tggcctttgt tttctggcat ctcgtaccac gaacatctac 840
 gaagaccctg cagtcattgg agcaccgggg tcagcaaaag 879

<210> 41
 <211> 2248
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> misc_feature
 <223> Incyte ID No: 1811877CB1

<400> 41
 ggaggaccag gaggacatca ctgcccttga cctcagccct gacaacgagg tgctgggtgac 60
 agccagtcgg gcattgctgc tggctcagtg ggccctggcaa gaggggcagcg ttaccgcgct 120
 gtggaaggcg atacacacg ccccccgtggc caccatggcc ttgcgacccea cctccactct 180
 gctagccaca gttggctgtg atggggccgt gcgcgtctgg gacatcgtag ggcactacgg 240
 gacacaccac ttccgaggct cgcggcggtgt cgtgcacctg gtggccttcc acccgagacc 300
 tacacgcctg ctgctcttct cctcggccac ggatgccgccc atccgcgtgt ggtcactgca 360
 ggaccggtca tgctctggctg tgctgaactgc ccactacagc gccgtcacct cactggcctt 420
 cagcgccgac ggccacacca tgctcagctc cggccgtgac aagatatgta tcatctggga 480
 ccttcagaag tcgcaggcca cgaggaccgt gccgtgtttt gagagcgtagg aggcctgctgt 540
 gctgttgcca gaggagccag tgtcccagct ggggtgtgaag tcccacgggc tgtactttct 600
 gacagctggc gaccaaggca ctctgcgcgt gtgggaggca gcttctgggc agtgtgtgta 660
 cagcgaggcc cagccgcggg gccctgggca ggagctgacc cactgcaccc tggcacacac 720
 cgccggctgt gtccctacgc ccaccgccga ccacaacctg ttgctctacg aggcctcgctg 780
 cctgcggctg cagaaaacgt tcgctggcta cagtgaaggag gttttggatg tccggtttct 840
 tgggcccagc gactccccag ttgtcgtggc ctccaatagc cccctgctaa aagtgtttga 900
 gctgcagacg taccgctgccc agatcctcca cggccacacg gatatcgctc tggccctgga 960
 tbtgttcogg aagggttggc tctttgccag ctgtgccaaag gatcagagcg tccgtatctg 1020
 gagaatgaac aaggctggccc aggtgatgtg cgtgggtcag ggttccggtc acacacacag 1080
 tgtgggcacc gtctgctgct ctaggctgaa ggagtccttc ctggtgacag gcagccagga 1140
 ctgcactgtg aagcttggtg ctcttcccaa agccttctgt tccaagaaca cagccccaga 1200
 caacggccct atctctctgc aggccacgac cactcagcgc tgcctatgata aggcacatca 1260
 cagcgtggct attgccccca acgacaagct gctggccaca ggctcacagg accgcacggc 1320
 caagctctgg gccctgccac agtgccagct gctgggtgtc ttctcaggcc accggcgctg 1380
 cctctggtgt gtcccagttc ctcccattgga ccaggtgctg gccacggcct cagctgatgg 1440
 caccatcaag ctctggggac tccaggactt cagctgtctc aagacatttg aggggacaga 1500
 tgctctctgt ctgaagggtg cctttgtgag ccgtggccag cagctgctgt ccagcggttc 1560
 ggatggcctc gtgaagctct ggaccatcaa gaacaacgag tgtgtgcgga cgctggatgc 1620
 ccacgaggac aaggtctggg ggtcgcactg cagccggctg gacgaccagc cctcactgtg 1680
 ggccactgac tcccagatca tctctggaa ggatgtgacc gaggcgagagc aggcagagga 1740
 cgaggccagg caagaggagc aggtggtcag gcagcaagag ctggacaacc tgctgcata 1800
 gaagcggtac ctgcggggcg tgggcctggc catctccctg gatcggcccc acaccgtgct 1860
 gactgtcatc gaggccatcc ggagggaacc tgaggcctgc tgaggactgg aagccaccat 1920
 gctccagact cggcgcgacc agaaagaggg cctgctgcgc ttctgctgca cgtggaacac 1980
 caactcgcgg cactgccacg agggccaggc cgtgctgggt gtgctcttga ggccagaggg 2040
 ccccgaggag ctgctggcct acgaaggcgt gcgggacgct cttgaggccc tgctgcctca 2100
 cactgtagcg cactttcagc ggctcagcag gaccctccag gccgcgctt tcttggactt 2160
 cctgtggcac aacatgaagc tccctgtgccc ggccgcccgc cccaccccct ggggaaaccca 2220
 taaaggcgca ctgccctaaa aaaaaaaaaa 2248

<210> 42
 <211> 2146
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1848674CB1

<400> 42
 gttattggca agttcccctg cagttgtttg ggctgtccct gtggctgggt ctggggtgtg 60
 cggccagcca tggagcgctc tggggccacg gaagtgcagc gctcagagcg gtcgggacgg 120
 gaccocgacg ttgcgggtcac catgggtctc acgggggttcg gtaaaaaagc tcgcacattt 180
 gacttggaaag caatgtttga acaaaactga aggacagctg tggaaaaaag tcgcaaaaca 240
 ctggaagcaa gaaaaaaga ggaagaatg aacagagaga aagaattaa aagacaaaat 300
 gaagatattg agcccaacatc ctcaagatca aatgtggtea gagattgctt caaatcatct 360
 tccagggata cgagcagcag tgaagtgaac cagagttctg actctcttga tgatgagtta 420
 attggccctc cttttacccc taaaaatggtt ggaaaaccag ttaattttat ggaggaagat 480
 atctcctggtc cttttacccc acctcttaat gaagaagaag aagaagcaga ggaagaagaa 540
 gaggaagagg aggaagagga aaatcctgtt cacaagattc ctgactcgca tgagataaac 600
 ctgaagcatg gcactaaaaa agtgtctgct tttgggtctgg atccctcagg tgcccgtttg 660
 gtgacaggag gatattgacta tgatgttaag ttttgggtt ttgtctgaat ggatgtcttc 720
 ttttaaggcat ttccgacctg tcagccctgt gagtgcctac agatcaagtc attacagtat 780
 agtaaacacg gagacatgat tcttgttgta tctggaagct ctcaggccaa ggtgatggac 840
 agagatgggt ttgaagtaat ggaatgtata aaaggagacc agtatattgt ggacatggcc 900
 aacacacagg gtcatacagc aatgcttcac actggctcat ggcattccaa aataaaggga 960
 gaatttatga cttgtc caaa tgatgcgact gtgaggacgt ggggaagtga aaatccaaag 1020
 aagcaaaaaa gtgtgttttaa accacggacg atgcaagcca aaaaagtcac tccactaacg 1080
 tgcacatata gttagagatgg aaacctcata gcagctgctt gccagaatgg aagcatacag 1140
 atctgggacc gaattttgac tgttcatcct aagttccact ataaacaggc tcatgactcg 1200
 ggacacagca cttctgtcgt gactttttcc tatgatggta atgtccttgc ctctgttga 1260
 ggtgacgatt cattaataat atgggacatc cgacaattta ataaacactt tttttcagcc 1320
 tcgggtcttc ccacatgttt ccaaatgact gactgctgtt tcagtcacga tgataagctc 1380
 atagtcactg gtacatctat tcaaaaggga tgtggcagcg gcaaatgtt tttttttgag 1440
 cgtaggactt tccaaagggt gtatgaaata gacatcacag atgctgagtt tgttcgcctg 1500
 ctgtggcatc caaagctgaa ccagatcatg gtgggaactg gaaatggatt ggctaaagtc 1560
 tattacgacc ccaacaagag tcagagggga gcaaaattat gtgtggttaa aaccacagcg 1620
 aaggcaaaac aagctgagac tctaactcag gactacatca tcacccctca tgccttgctt 1680
 atgtttccgt agccccgcca acggagtaca aggaacacg tggagaagga cagactggat 1740
 cccctgaagt cgcataaac tgaacctcct gtgacaggcc caggttcgtg tggccagatt 1800
 ggaacccacg ggggcatctt ctcttctat atgtggaaga acattgcttt ggacaagacc 1860
 gatgacagta atcttcggga agccattttg cgtcatgcca aagcagcaga agacagacca 1920
 tattgtgggt ctcacagata ttcaagact cagcccaaaa ccatgttttg caagttgaa 1980
 tctgtgatg aggaagcaaa gaatgagcca gaatggaaaa aacgtaaaaa ttgaagaatc 2040
 tcatttgaga gctgtttgca tgagttggga gggatggga caggttttgg tttttttttt 2100
 atgctcatga aattaaaaat tcatttttat gaaaaaaaaa aaaaaa 2146

<210> 43
 <211> 714
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2012970CB1

<400> 43
 ggatggcgag cagcggaggc gaggcgggtg cgagagcagc ggctccgcca ttgacgagg 60


```

aggcctgagg gacggggccag cgggtgcacaa gaagagaccg agggcggttg ccccgagaga 120
gccaggggcca tggaggcccaa catgccgaag cggaaaggagc ctggcaggtc tctccgtatc 180
aaagtctact ccatggggcaa cgccgaagtg gggaaaagct gtattataaa gcgataactgt 240
gagaaaaagat tctgtgtctaa atacctggca acaattggaa ttgactatgg agtcacaaaag 300
gtacacgtca gagacagaga aatcaaaagt aacatctttg atatggctgg accatcccctc 360
ttctatgagg ttcgaaatga gttttacaag gacacacagg gtgtgatact ggctctatgat 420
gttgggcaga aagactcctt tgacgccttt gatgcgtggc tggcagaaat gaagcaagag 480
cttggacctc atggaacatc ggaataatc atattttag ttgtgtccaa caagatgtat 540
tgtaccaaac atcgctctgt agatgaagt gaaggacgtc ttgtggctga aagcaaaagg 600
tctctgtact ttgaaacttc agcacaact ggagaaggca ttaattgagt gttccagata 660
catcttggat agaactaatg gataaattag tctgtttaaa aaagaaaaaa aaaa 714

```

<210> 44

<211> 1779

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2254315CB1

<400> 44

```

caggagggaag atggcgccgt ccgcagctgc cgctgagctc caggcttctg ggggtccgcg 60
gcacccagtg tctctgttgg tgttgggaat ggcgggatcc gggaaaaaca cttttgtaca 120
gaggtctcaca ggacacactgc atgcccaagg cactccaccc tatgtgatca acctggatcc 180
agcagatcat gaagtccctt tctctgccaa tattgatatt cgtgatactg taaagtataa 240
agaagtaagt aaacaatatg gacttggacc caatggcgcc atagtacct cactcaatct 300
ctttgtctac agatttgatc aggtgatgaa atttattgag aaggcccaga acatgtccaa 360
atatgtgttg attgacacac ctggacagat tgaggtatcc acctggtcag cttctgggac 420
aattatcact gaagcccttg catcctcatt tccaacagtt gtcactctat taatggacac 480
atcgagaagt accaacccag tgaccttcac gtccaacatg cctatgcct gcacatcct 540
atacaaaacc aagctgcctt tcatttgtgt cactgaataaa actgacatca ttgaccacag 600
ctttgcagtg gaattggatgc aggtatttga ggcttccaa gatgccttga atcaagagac 660
tacatactgc aataaacctga ctggttcaat gagcctgggt ttgatgagt ttacagctc 720
actcagggtg gtgggtgtct ctgctgtct gggactctga ttgatgaac ctttgtgtga 780
agttaccagt gctgccgaag aatatgaag ggagtatcgt cctgaatatg aacgtctgaa 840
aaaatcactg gccaacgcag agagccaaca gcagagagaa caactggaac gccttcgaaa 900
agatatgggt tctgtagctt tggatgcagg gactgcacaa gacagcttat ctctgtgtct 960
gcaccccttc gatttgatcc tgactcgagg aaaccttggat gaagaggatg aggaagcaga 1020
cagcgatact gatgacattc accacagagt tacagaggaa agccatgaag agccagcatt 1080
ccagaatttt atgcaagaat cgtatggcaca atactggaa agaaaaaata aataggagac 1140
tttagcacac ttcaacttgtt tctagaagtc cagaatttgg gacctccacg tgaaagaact 1200
ttcttacct ctgaaactgg ggtccccata agggataatt ttccctcag tagcaaaagt 1260
tctctctaa gagaaattct gtgactcaga tgaagtccag gatagaagac ccttggacct 1320
ggcaggttat tctgtattat tcttggcct tctcctgtga tttatgtcga gaaggatata 1380
ctgagctgat actcttccaa gctcacaact tcaagtttta tcaattgaac tcaagtaact 1440
ttgtgctgta ggaattggaat caaaagaac tagtctctct gtgaccacct cagatctcta 1500
ttattaggct agatgtatag cctctactcc cccagctctc tgcctctgac cctgcaactgt 1560
aagttgccct tctatttaga gccaaaggaa agggaaacat gagcttatcc agaagcgggtg 1620
cagagtctcc ttggcaatca accaacgttg ctatgaata tcgctcacac tgtatagctc 1680
attataggac gtcagggttg ttgaaaaaag tgggcaagac atgattaatg aatcagaact 1740
ctgtttcatt ggtgacttgg ataaagactt ttttaattt 1779

```

<210> 45
 <211> 2234
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2415545CB1

<400> 45
 cccctctctga aaaatgggttt caccccgctcc ttctctagaa cgggtcgggc cgtaccact 60
 gtccggcccg gaggggaact gttttctccg gaagtgaaca cagcctgact agggaaagga 120
 ggaggcgggg cagtggggcc ttcggcgccg actatggaag gagccggcta cagggtgggtg 180
 tttgagaagg gcggagtgtta cctgcacacc agcgctaaga agtatcagga ccgagactct 240
 ctcatcgctg gtgtcatccg tgtcgtgaaa aaggacaatg acgtcctcct gactctgggtc 300
 cctgtagagg aggtcggaga ttccacccaa atcctcttct ccaagaagga ctccagtggg 360
 ggtgactcat gtgcttctga ggaggaaaca acctttgacc ccggtatga accctgactgg 420
 gctgtcatca gcaactgtgc gccacagccc tgccactcag agccccagag aggtgcagag 480
 cccagctgcc cccagggtcc tcagtgagtc tgggggagct aaagtccatc 540
 cgccgctcca agcaggcct cagctgggcc tacctgggtc tggtgaccca ggctggaggt 600
 tccctgcccg catcgcaact ccaccgctgg ggcaccgctg cctctcctcg cgtcctcagc 660
 cgctacctgc tgttggccag ctcccgagc gactcccgcc tctacctgt ctccccacc 720
 gaactcctctg ctctctccaa ctctctccac cactctgcag tctttgacca ggacagctcc 780
 aatgtggtgt cacgcttctc ccaggatccc tactccacca ccttcagcag ctctcccgga 840
 gtgaccaact tcttcggggg tgccctgcag ccacagcctg agggagccgc ctccgacctt 900
 cccccccaa ccgacgatga gcccgagcct ggattcgagg tcatttctct tgtggagctg 960
 gggcctcggc caaccgtgga gccgggcccct ccagttacag aggaggagtg ggcagccacc 1020
 gtgggcccctg aaggtcgctt cgagcaggtc cctgagctga agaaccgagt ctctcgggg 1080
 ggtctgagcc ccagcctgcg gcgcgaggcc tggaaagtcc tctagggta cctcagctgg 1140
 gaaggcacag ctgaggagca caaggcccaa atacgcaaga aaacggatga gactttccgc 1200
 atgaagctgc agtgaaatc tgtgagccct gagcaggagc ggagaaactc acttctgcat 1260
 ggataccgca gctcatcga aaggatgtg agccgcactg acaggaccaa caagtctctac 1320
 gagggctccg agaaccggg gctgggccc gctgaacgata tctctctccc ctactgcatg 1380
 tatcacttgc acctcggcta cgtccagggc atgagtgatc ttctctcccc gatcctctac 1440
 gtcattcaga accaggttga tgccttcttg tgttctcttg gcttcatgga gctggtgcaa 1500
 gggaactttg aagagagcca ggagaccatg aagcggcaac tccggcgact cgtgctgctc 1560
 ctgagggtgc tggacccccct gctctgcgac ttctctggat cccaggactc ggctctctc 1620
 tgcctctgtt tccggtggct cctcatctgg ttcaaggagg aattccccct cccgagtgct 1680
 ctctggctgt gggaggtgct gtggacaggg ctccctggcc ccaatctgca cctctggtg 1740
 gcttcgcca tctgggacat ggagaggggac acctcatgc tgcctcggtt cggctccaat 1800
 gagatcctca agcacatcaa cagctgact atgaagctga gcgtggagga cgtcgtgacc 1860
 cgcgcggagg cctgcaccc ccagctaac gccctgaccc gagctgcct ccaacgtgca 1920
 ggagatcctg gggctggccc cgccacgcag agccccacag ccctcgccc accgctctcc 1980
 cgctgctctc gtaagccac ccgggcccgc ccaccccgc ccctctcac ggacacagcc 2040
 ccgacgccc acagcagcct ggagatcctg cccgaggagg aggcagaggt cgcgactcct 2100
 taaccocgoc aggcagctc gttctgcaca ggcaactttag cccgagccag gcaacactgc 2160
 gagggggcag gtgtgtctcg ccgccctgct gataagctgg ctctattaaa ctgacacttc 2220
 tcaaaaaaaa aaaa 2234

<210> 46
 <211> 3150
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> 96, 97, 99, 3070-3072, 3074, 3078, 3080-3082, 3085-3087, 3091,
 <222> 3099, 3100, 3103, 3107, 3110-3112, 3114, 3115, 3121, 3123, 3125,
 <222> 3128, 3136, 3138, 3140, 3141, 3143, 3145, 3147, 3149
 <223> Incyte ID No: a o r g o r c o r t, unknown, or other

<220>
 <221> misc_feature
 <223> Incyte ID No: 2707969CB1

<400> 46
 acacaggagc aatgcaaaat ttgataaagc atttttctgt cagatcagct gagccctact 60
 gcccttccct tcaagattcc tggagaccca gatgtnnngna tcttttcattg acaacaaaat 120
 aatgtgtcat gatgatgatg ataaagaccc tgtactccgg gtattttgatt cccgagttga 180
 caagctcagg ctgttgtaag ttcggacacc tactctccgt acatccatgt accagaagtg 240
 taccactgtg gatgaagcag agaagacaaat tgagctgcgt ctggcaaaaa ttgaccatac 300
 tgcacattcc ccaactttac ttgacatgaa gattggacaa gggaaaatgt agccgggctt 360
 ctccctctaa ctgcagctctg atgtactttc cactggggcca caggcaacaa agtggacgaa 420
 aaggaatgcc cctgccacgt ggagggcgaa agatcggcag aagcagcaca cagaaacact 480
 gcgttttagt aatgacaga gggagaaagta catccaggaa gccaggcata tgggcagcac 540
 tatccgccag cccaactctg ccaactctct tccatcagtg attgccaga ccaattgaaa 600
 gtttgttagg gccctgctga aggaatgccg caataagacc aagaggatgc tgggtgaaaa 660
 gatggggcca gaagctgtgg agctagggca tggggaggtg aacatcacag ggggtgaaag 720
 gaacaccctg atttgccagc ttttgtatct cctggaaaag atctggagtc atggactaca 780
 agtgaacacg gttgaatcac ccttatggtc ccactctgta cattatcagg acaaccgcca 840
 gagaaaaact acatcaggaa gccctcagta ctccaggata ctctttgatt cagaaactag 900
 gaagctctga gccagctcac tcatgcctcc cctgaggatc cctctgattc aggatagtag 960
 gccatccagc aacatcgggg aaatcaagac tgaatgtggg aaggccagag catgggtgag 1020
 actgtcccatg gaaaaaaagt tactttccag acacctgaag cagctcctct cagaccatga 1080
 gctcaccaaa aagttatata agcgtatgct ctccctgcgc tgtgatgacg agaaggagca 1140
 gttcctctat cactcctctg ctttcaaatg cgtcgattac tttgtcttca ccaatgtctt 1200
 cacaactatc ctgatccogt accacattct gatcgtaacca agcaaaaagc tggggggctc 1260
 catgtttact gccaacccat ggaatctgtat acaggagaaa ttgggtgaga cacagattac 1320
 cgagattccc aggaatgtgc tagagatgac ctcgagtgcc cagaactggg ggaagcttac 1380
 tactgtccag attgcccagc ataactctgg gctgtatgcc aaatgggtgt tggagtatgt 1440
 gatggtcagg attagatcca caggacatac ctacaagttc ccgtgtggcc tgggttaggt 1500
 gaaggcgcat gatgatgaa gccctggagcg gatcctagtt ggggagctgc tcatatcccc 1560
 gccctaggctg ctgagagggc catgcggcag cccgcgcgtg cagcagctcc ccagtgctat 1620
 ccggaggcct gttaccatct caccacaaca caagcccaag ctgaacactg ggcagatcca 1680
 ggagctccatc ggggaggcag tcaatggcat tgtgaagcag ttccataaag ctgaaaaaga 1740
 gcgagggcgt ctgacgcgtg tgctctgtgg agagtgtggc cttgtctcgg ccttggaaac 1800
 ggctttccag catggtatta aatcgcccc gctctcaaaa aatgtcttca tttgggattt 1860
 gctggaaaaa gcacaaaact attatgagac attagagaa aatgaagtag tccctgagga 1920
 aaactggcat ccaagagccc ggaactcttg cgaatttgtc actgcaatca acaatctccc 1980
 ttctacacat gccaaagtga gcaagtttca gatctgggtg tgcttgggag ccagagatca 2040
 cctctacac cactggattg cctcgtctgc tgactgccc ttgattctgt tgcctgagac 2100
 ggatgtggca ctgatcaag accatacact tgccaattcc ttgattctgt tgcctgagac 2160
 attgcaggag ttcaacatca cgtctggagc gtcccttgtc aagggcctag acatctgacc 2220
 tccagcacc agccagcagc aggaactgaga aagaactcacc ctgcagctct gaccttttt 2280
 cccaaaaggga cttaagcgat tgtgcaggag taggagacaa aatgtacact cactgtaaaa 2340
 agagaactag aggaattttg gaataaataa tctattttag agtttatttg ctgatttgtt 2400
 ttttaccac tttcatgtga aagagtata gggagaggga gcaggctgtg tgcgccttat 2460
 tttgaagctg gtgcctccc tcgcccgtgg cactgctggt aagcctgagg cctccctgga 2520
 ctgagcctgt ggcactgcgt gcgggacagt tatgttctct tgccccgtg cattaattag 2580

```

gccccccac atcattttta aactaatgtt ttcttatatt aacattatta tggatatattg 2640
gctttcatag gccacacaca ggtgtgtctgc gcgggaagcc ccatgctcca atcaaggga 2700
tttttagtag tgcctctaag caagcaccga tgaagtcagtc ccacgtatatt tcttttttgt 2760
cagtatgttt tggggaaggag acatgcgggg atgtgtcatc gtgcccataa cccattttcc 2820
tgtgggcaca gtttcacaga agtaaacata agcatgtttt aacaggtttt tctttttctt 2880
ttcttttttt aaaaattttt atttatttaa ccogccattg tgtgttttta agtattttct 2940
ttttttaagg aaaggaaaag cttgtcacaa tctaactggc tatgttatta ttataaaatt 3000
tatgttttgc aacttagaaa ccagctacag tatggcccac ttaataaaac acctgaataa 3060
aaaaaaaaagn nngnngngnng nngtnnnngag naggagggnn gngngngngn nngnngggag 3120
ntnantntgt ggggtgngnng nngnngnngnt

```

<210> 47

<211> 1806

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2817769CB1

<400> 47

```

gtctgcgcgc aggtgcccgt cggcgccggc gccgcccgtt ccgcggctgt cgcgcgcgtc 60
gtgcgtgcgc ctgcggggag gggacggggc tgcgttctct cctccttctt cccgcctctc 120
agctgcgcgc ccgaaccttt tctcgctgcc gctgggagcc cgtgtcatcg cccaggccga 180
gcacgatgcc ccgtaaaaag ggaggtgatg gaattaaacc acccccaatc attgggaagt 240
ttggaacctc actgaaaaat ggtattgtgt gattgccaaa tgttgggaaa tctactttct 300
tcaatgtgtt aaccaatagt caggcttcag cagaaaaactt ccgcttctgc actattgatc 360
ctaattgagag cagagtacct gtgccagatg aaaggtttga ctctctttgt caataccaca 420
aaccagcaag caaaaattcct gcctttctaa atgtggtgga tattgtctgt cttgtgaaag 480
gagctcacaa tgggcagggc ctggggaatg cttttttatc tcatattagt gcctgtgatg 540
gcattcttca tctaacacgt gcttttgaag atgatgatat caagcacgtg gaaggaaagt 600
tagattcctat tcgagatata gaaataatac atgaagagct tcagcttaaa gatgaggaaa 660
tgattgggccc cattatagat aaactagaaa aggtggctgt gagaggagga gataaaaaac 720
taaaaacctga atgtgcaaa gtaaaatcct gggttatagt caaaagaaac 780
ctgttcgctt ctatcatgat tggaaatgaca aagagattga agtgttgaat aaacacttat 840
ttttgacttc aaaaccaatg gtctacttgg ttaactcttc tgaaaaagac tacattagaa 900
agaaaaacaa atggttgata aaaattaaag agtgggtgga caagtatgac ccaggtgctt 960
tgggtcattcc ttttagtggg gccttggaa ccaagttgca agaattgca gctgaggaga 1020
gacagaagta cctggaagcg aacatgacac aaagtgcctt gccaaaagatc attaaggctg 1080
gggtttgcagc actccaacta gaatactttt tcaactgcagg ccagatgaa gtgcgtgtcat 1140
ggacatcagc gaagggaact aaggctcttc aggcctgcagg aaagattcac acagattttg 1200
aaaaggagat cattatggct gaagttaatg aatcacgaaga tttttaaagat gaaggttctg 1260
aaaatgcagt caaggctgct ggaaagtaca gacaacaagg cagaattatg attgttgaag 1320
atggagatatt tcatctcttc aaatttaaca cactcaaca ccggaagaag aatacaaaat 1380
tagtctatgc tcagataaac atacaacctt caaaaggcat ctgattttta aaaaatttaa 1440
attttctgaaa accaatgcga caaataaaat tggggagatg ggaattcttt accaaacaaat 1500
tattttttatt tgttttaaaa ttaaaatact gtgtaccccc ccccaactca tgaattgcag 1560
gttcaactaaa tgtgaacagc ttgtcttttc acgtgattaa gacctactc caaattgtag 1620
aagcttttca ggaacacatc tactctcatg atacttcatt aatctccatc atgtatgcaa 1680
agcctgacac atttgacagt gaggacaatg tggcttgctc ctttttgaaat ctacagataa 1740
tgcatgtttt acagtactcc agatgtctac actcaataaa acatttgaca aaacccaaaa 1800
aaaaaa

```

<210> 48
 <211> 2880
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2917557CB1

<400> 48
 gagaggaggagg agaggagaaga agaggagaaga gaagatgaag aaagtgaaga agaggaggaa 60
 gagaggaggagg aaagtgaagg cagtgaaggat gatgagggaag atgaaaagggt gtcagatgag 120
 aaggatttcag ggaagacatt agataaaaaag ccaagttaag aaatgagctc agattctgaa 180
 tatgactctcg atgatgatcg gactaaagaa gaaaggggctt atgacaaaagc aaaacggaggg 240
 attgagaaaac ggcgacttga acatagtaaaa atgtgtaaaac ccgaaaaagct aagagccctc 300
 attatctcgcg tacttgggca tgtggacaca ggggaagacaa aaattctaga taagctccgt 360
 cacacacatg tacaagatgg tgaagcaggt ggtatcacac aacaaattgg ggccaccaat 420
 gttccctcttg aagctattaa tgaacagact aagatgatta aaaaatttga tagagagaat 480
 gtacggattc caggaaatgct aattattgat actcctgggc atgaatcttt cagtaatctg 540
 agaaaatagag gaagctctct ttgtgacatt gccattttag ttgttgatatt tatgcatggt 600
 ttggagcccc agacaattga gtctatcaac ctctcctaat ctaaaaaatg tcccttcatt 660
 gttgcactca ataagattga taggttataat gattggaaaa agagtctctga ctctgatgtg 720
 gctgctactt taaagaagca gaaaaagaat acaaaaagatg aatttgagga gcgagcaaaag 780
 gctattattg tagaatttgc acagcaggggt ttgaatgctg ctttgtttta tgagataaaa 840
 gatccccgca cttttgtgtc ttgggtacct acctctgcac atactcgtga tggcatggga 900
 agtctgatct acctcttctg agagttaact cagaccatgt atagcaagag acttgacacac 960
 ttgtaagagc tctgagacaca ggtgatggag gttaaagctc tcccggggatg gggcaccact 1020
 tagatgtgca tcttgatcaa tggcgctttg aaggaaggag atacaatcat tgttctctgga 1080
 gtagaagggc ccattgtaac tcagattcga ggcctcctgt tacctcctcc tatgaaggaa 1140
 ttacagtgta agaaccatg tgaaaagcat aaagaagtag aagcagctca gggggtaaaag 1200
 attcttggaa aagacctgga gaaaaacattg gctggtttac cctcctctgt ggctataaaa 1260
 gaagatgaaa tccctgttct taaagatgaa ttgatccatg agttaaagca gacactaaat 1320
 gctatcaaat tagaagaaaa aggagtctat gtccaggcat ctacactggg tcttctggaa 1380
 gctctactgg aatttctgaa aacatcagaa gtgccctatg caggaaatga cattggccca 1440
 gtgcataaaa aagatgttat gagggcttca gtgatgttgg aacatgacct tcgatgtgga 1500
 gtaattttgg cttctgatgt gagaattgaa cgagatgcac aagaaatggc tgatagttaa 1560
 ggagttagaa ctttttagtg agaaattatt tatcattttat ttgatgcctt tacaaaatat 1620
 agacaagact acaagaaca gaaacaagaa gaattttaagc acatagcagt atttccctgc 1680
 aagataaaaa tctcctccca gtacattttt aattctcag atctcgatagt atccgatagg gaatgggggt 1740
 acggtggaag cagggtcaggt gaaacagggg acacccatgt gtgtcccaag caaaattttt 1800
 gttgacatcg gaatagttaac aagtattgaa ataaaccata aacaagtgga tgttgcaaaa 1860
 aaaggacaag agtttttgggt aaaaatagaa cctatccctg gtgagtcacc caaaatgttt 1920
 ggaagacatt ttgaagctac agatattctt gttagtaaga tccagccgca gtccattgat 1980
 cgcactaaaag actggttcaag agatgaaatg cagaaggagt actcgccagt tatgtggag 2040
 ctgaagaaaag tatttgaaat catctaattt tttcacatgg agcaggaact ggagtaaatg 2100
 caatactgtg ttgtaaatc ccaacaaaaa tcagacaaaa aatggaacac agctatttgg 2160
 acactatggg acttaagtat ggaaggaaag aaaaatgggt tataaaatgt tttccatgag 2220
 aaaccaaaga acttaacactg gtttgacagt ggtcagttat atgtccccac agttccaatg 2280
 tgccctgttca ctcaactctc ccttccccca ccttctctca ctctgctctc gttttaaagt 2340
 tgccctctcc ccaaaatttg atttttatta cagatctaaa gctcttttga ttttataactg 2400
 attaaactcg tactgcagta ttgattaac caagctctcg cagattttgt tatgtggag 2460
 acttttttga cgttaagaaat acttctttat ttatgcatat tcttcccaca gtgatttttc 2520
 cagcattctt gtgccattg ctttttagggc ttttataaaa tagaaaaata ggcattctga 2580
 tattttttta gctgctttgt gtgaaaccaa ggtgtgaaaag cacagctggc tgctttttac 2640
 tgcttgtgta gtcacgagtc catgttaatc atcacattc taacccaacac taccataaaa 2700

gaaaacagac atccaccagt aagcaagctc tgttaggctt ccatgttagt gtagcttctc 2760
 tcccacaagt tgcctccta ggacaagaat tatcttaca catcacacta 2820
 cctgtgatgc cagcacctgg taacagtaga gatttttata cattaatctt gatctgtttt 2880

<210> 49

<211> 1109

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3421335CB1

<400> 49

cccacgcgtc cgctcgctct tgggtcatgc ctggccagca gaaagcagct ccatagggga 60
 ggagagccac cgaggtatct acagctcgag tctaatagta acacagagga ttccagcagt 120
 gccaccatgg gtctctgtgaa tccagaggt cacaaggcgg aagcccaggt ggtgatgatg 180
 ggctctggact cggcggggcaa gaccacgctc ctttacaagc tgaaggggca ccagctgggt 240
 gagacccctg ccaactgttg tttcaacgtg gagcctctga aagctcctgg gcacgtgtca 300
 ctgactctct gggacgttgg ggggcaggcc ccgctcagag ccagctggaa ggactatctg 360
 gaaggcacag atatcctcgt gtacgtgctg gacagcacag atgaagcccg cttacccgag 420
 tcggcggctg agctcacaga agtccctgaa gaccccaaca tggctggcgt ccccttcttg 480
 gtgctggcca acaagcagga ggcacctgat gcacttcgcg tgcctaaagt cagaacacag 540
 ctgagcttag agagattcca ggaccactgc tgggagctcc ggggctgcag tgcctcact 600
 ggggaggggc tggccgaggg cctgcagagc ctgtggagcc tectgaaatc tcgcagctgc 660
 atgtgtctgc aggcagagc ccatgggggt gagcgcggag acagcaagag atcttgatcc 720
 agacagagca gcatattctt gctcatacaa actagaagaa ccagctgac ttctgagaaat 780
 ttacgccttag tctatcaaac aagaaatgct ggcttggccc ggtggctcat gcctgtaatc 840
 ccagcatgtt gggagaccac ggtgggggaa tcccttgagc ccaggagtgt gagagcaaca 900
 tcacaacacc ccaatttctac taataatcaa aaaaattggc gggcatggtt gcactgtgct 960
 ttagtccacg ctacttggga ggctgaggca ggagaatcgc ttgagcccaa gaggtagagg 1020
 ttgcagtgag ccaagatcgc gccactgcac tccagctctg gcaacagagt gagacctgt 1080
 tctagtgggt ataataataa tgatgtagt 1109

<210> 50

<211> 2407

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 605761CB1

<400> 50

ctttcgtctg cgccattacc tgagttctcc tccagcgctt ccgaccctc tccgattagc 60
 ggtcccagga gtttccaagg taaccgcgca gtaggcgga tctcattagg cggaaagcga 120
 aacccggaag tgacgtctct accgggtgtc agcagcgaga ggggtcgagg atggcggcgc 180
 gcaagggtcg ggtgcgcagc tgtgaaacgg ggaagcccat ggaagccgag tccggcgaca 240
 caagttccga gggcccgccc caggtctacc tgcccgccc gggcccgccg ctacgcgaag 300
 gggaggagctt ggtcatggac gaggaggcct atgtgctcta ccacagagcg cagactggcg 360
 cccctgtctc cagctttgac atagtcgggg atcacctggg agacaacogg acagagcttc 420
 ctcttacctt ttacttctgt gctgggaccc aggtctgagag cgccacagagc aacagactga 480
 tgatgcttcg gatgcacaat ctgcatggga caaagccccc accctcagag ggcagtgatg 540

```

aagaagaaga ggaggaagat gaagaggatg aagaagagcg gaaacctcag ctggagctgg 600
ccatggtgcc ccaatctggt ggcatcaacc gagttcgggt gtcattggctg ggtgaagagc 660
ctgtggctcg ggtgtgtgtca gagaagggcc agstggaggt gtttgccgtg cgccggcttc 720
tgacagtggt ggaggagccc caggccctcg gctggacaca cagccttctt cgtggatgag caggcccaaa 780
tgaagcccat ctctcctctt ctgaccggtg actgtcaaaa gaacatccac ctctggacac 900
cccgggtgac cgggtgcctg cactgtggac acgggccatt cgtgggccac acacgctctg 960
ctacggcagcg cgggtccttg cactgtggac acgggccatt cgtgggccac acacgctctg 1020
tggagagacct cagatgtgtca ccgactgaga acacgggtgt tgcctcctgc tcagctgacg 1080
cctccatcgc catctgggac atccgggcag cccccagcaa ggcctgcatg ctcaccacag 1080
ccaccgcccc tgatggggac gtcaatgtca tcagctggag ccgcccggag cctctcctgc 1140
tcagtgccgg ggtatgatgg gccctcaaga tctgggacct tctggcagtt aagtctggtt 1200
ccccagtgcc cacttcaag cagcacgtgg ccccgtgac cctcgtcgag tggcaccccc 1260
aggacagcgg ggtcttttga gctcgggtg cagaccacca gatcacacag tgggacctgg 1320
cagtggagcg ggaacctgag gcggggcagc tggaggccga ccccgagctg gccgaccttc 1380
cgcacgagct gctgttcctg caccaggggc agaccgagct gaaggagctg cactggcacc 1440
cgactgtccc agggctcctg gtcagcacgg cgtgtcagg ctccaccatc ttcgcacca 1500
tcagcgtctg aggcgtccca ctgggtctga tcttctcttc tgcctggaaa ctgaagtcca 1560
attgggtccc cctggaaggg gtctcatcag gtcctgttgc tgagacttgc cggcctgtgg 1620
gctgcccgtg tgatctctgt ttgacttctg gttctctaga aggcctggct ctgatccagt 1680
gacctctctc accaagaagc tcggtttaac cagggtcctg taagaccact cccaccaga 1740
gacttgtgtg gctgtgtgtg gctgtgtgtg cggattcctt cctgtcagct gtgacctatt 1800
tgactgtgtg ccccaagacc cagt.tttttg tttgtttgtt tgagacggag tctgtgtctg 1860
tcgcccagcg tggagtgcag tagcacgac tcggctcact gcaacctccg cctcctgggt 1920
taaaagtgat ctctcagctc agtctccagg gtatctggga ttacaggcat gtgccaccac 1980
accccgtaaa tttttgtatt tttagttag acggggtttc accatgttgt cagggtgggt 2040
ctcaaatctt tgattctcag tgactctgtc gccccggcct cccagagtcg tgggttggga 2100
ttacaggcgt gaggcacgcg tctccgctca ggaccagatt ttggctctcg gtccccagea 2160
ggggactcgg gggatataca gtggctgcac caaatggag gctgtggggt gttccacaca 2220
attgctctgc gaccttgttc tctctgccag ctgggtttgg ccagagattc tcggttgggt 2280
ggctacatct gaccctcttc cctcctcct gactttagag cgtgggtgag tgcggggagg 2340
aaggtcaggg ctctcagcga gcaataaagg accaggaaga ggctgtgaggt gtaaaaaaaa 2400
aaaaaaa
2407

```

```

<210> 51
<211> 1158
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 483862CB1

```

```

<400> 51
ggaagaccgt cccggatggc ctccgggact gccagtggt ggaggtgagc tcggggattg 60
ccggcatctc cgtctctgct ggttgcctca tgcctcagcg tgcggcgctc agccctcgct 120
cgcatgtgtg gcgctgaggt gccggggcag caagtgcacat gtcgtccggc ccccgccgct 180
ctgacttccc ccgctggaag cgcacacatc cggagcaact gaggcgccgg gaccggctgc 240
agagacagcg gtcgaggag atcactcctgc agtataacaa attgctggaa aagtcagatc 300
ttcattcagt gttgcccag aaactacagg ctgaaaagca tgacgtacca aacaggcacg 360
agataaagtc cggacatgat ggcacatgga atgacaatca gctacaagaa atggcccaac 420
tgagagctaa gcaccaagag gaactgactg aattacacaa gaaacgtggg gagttagctc 480
aactgttgat tgacctgaat aaccaaagtc agcggaagat caggagagatg cagatgaatg 540
aagcaaaaat tgacagaatg ttgcagacta tctctgacct ggagacggag tgcctagacc 600
tgcgactaa gctttgtgac cttgaaagag ccaaccagac cctgaaggat gaatatgatg 660

```

```
cctctgcagat cacttttact gccttgagg gaaaactgag gaaaactacg gaagagaacc 720
aggagctggt caccagatgg atggctgaga aagccacgga agccaatcgg cttaatgcag 780
agaaatgaaa agactccagg aggcggcaag cccggctgca gaaagagcct gcagaagcag 840
caaaaggaaac tctaccagtc gaacaggatg atgacattga ggtcatttgt gatgaaactt 900
ctgatcacac agaagagacc tctcctgtgc gagccatcag cagagcagcc acgtaagttag 960
gcaggttttgg gccagggaaa agacagcttg aggagcaata tgaaggcaca tctgtggaca 1020
tgacaaagaa tgcagtcaga tgcacccaac cccttactcc tttctggga caccagcgt 1080
cgaacacacc acagaggtgt ctagtcttcc tcagttcacc tctgtctaata gggagggaag 1140
cagaacacgg gtggcttc 1158
```

<210> 52

<211> 1026

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1256777CB1

<400> 52

```
ctgcgggctt ctctcgtcg ccatggaaac gaaagcggcc aagtagagct cgtcctgac 60
gcgcgccttc cgttgggctc cggccggcta agccgcggcg gacaactatg ctgaaagcca 120
agatcctctt cgtggggcct tgcgagagtg gaaaaactgt tttggccaac tttctgacag 180
aatcttctga catcactgaa tacagcccaa cccaaggagt gaggatccta gaatttgaga 240
accgcgatgt taccagcaac aacaaaggca cgggctgtga attcgagcta tgggactgtg 300
gtggcgatgt taagtgttag tcttgcctgg cggccctgat gaaggatgt catggagtgg 360
tgatcgctct caatgctgac atcccaagcc acgggaagga aatggagatg tggattctcc 420
gttttgtcca acagcgttc ttacaggaca cacagtgtat gctaattgca caccacaac 480
caggctctgg agatgataaa ggaagcctgt ctttgtcgcc acccttgaac aagctgaagc 540
tggtgcaact aaacctggaa gatgaccttg aggagatcgg gatggaattc ataaagtatt 600
taaaaaagcat aatcaactcc atgtctgaga gcagagacag gaggagatg tcaattatga 660
cctagccagc ctccacttgg gactgccaca tccccagtg aatcagcatg tttctcgggt 720
cagatctgaa atcacatoca gctcctgatg ttttctctc cctctgacg cagaggaagt 780
gttctactct gcaggaaagg acctgtcaca cagggcgctt actcagacca tctgtctct 840
gccctgagtt cagttgagaa aatcctatta tcaaatattg atttctctgg ccagaaactt 900
cccaagagacc tgtaaaatgg agggatttac cactccacat atgtccagtt aaacagtttg 960
tggacttgta accgtcgag cccaatgata caacagtagt ttaatcacgt gaaaaaaaaa 1020
aaaaaa 1026
```

<210> 53

<211> 2456

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2198779CB1

<400> 53

```
gacgagcgac gagatgacgg aggagcgggt ggccaacgca tgcggcgagt cgggtgtaaac 60
aaggcctcgc gccgcgtcgg gtccgtgcag cgtccctggc tgggatttgc attggctgg 120
cagagagggt acctggaat ccaacaccgc ccaacacccc tcccgtctcc cagtcgggg 180
acttcgatag gattggagaa ggagttgaca ggaggagccc ccgcacagga cctaagaagt 240
```


ctgtgaccag aagatgggat cgcggaacag cagcagtgca ggatccgggt ccggagacc 300
 ctcccgagggc ttgccccgaa gaggggctgg cctgcgtcgg agtgaggaag aggaagaaga 360
 ggatgaagat gtggatccgg cccagggtact ggctctatct ctcgcgcagag ccgaagtggag 420
 gttggtgcag ggaggaggat cagcaaaatt acaattcatt caggccctct tggactcaga 480
 ggaagagaat gacagacctt gggatggtcg tcttggggat cgatacaacc cacctgtgga 540
 tgctaccctt gacaccgggg agctggaatt caatgagatc aagacaacag tggaaactggc 600
 cacaggcgag ctggggctta ggcgggcgcg ccagaagcac agctttctcc gaatgttgca 660
 ccagagagaa cggggcctct gccatcgggg aagcttctcc cttggagaac agtctcgagt 720
 gatatctcac tctttggcca atgactctgg cttcaactgat agtactctc caaaggcttt 780
 ctgtggcatc tacagcaaa agtggtcaaat attcatgtct gcttgccaa accagacaat 840
 ccgactctat gactgcccag atggccgttt ccgtaaaatt aagagcatca agggcccgca 900
 tgtaggctgg agcgtcttgg atgtggcctt caccctgtat gggaaccact tctctactc 960
 tagctggtct gattacattc atatctgcaa tatctatggt gagggagata cacacactgc 1020
 cctgggtctc aggccagatg agcgtcgctt tgctgtcttc tccattgctg tctctcaga 1080
 tggacagaaa gtactaggag gggccaatga tggctgctg tatgtcttt accagaaca 1140
 gaaccggcgc accctctagg ttgagtcoca tgaggatgat gtgaatgcag tggcctttgc 1200
 tgatataaag tcccaaatcc tgtttctctg gggagatgat gccatctgca aagtgtggga 1260
 tcgacgcacc atcggggagg atgaccccaa gcctgtgggt gcactggctg gacaccagga 1320
 tggcatcacc ttcattgaca gcaagggtga tgcccggtat ctgatctcca actctaaaga 1380
 ccagacatc aaactctggg atatccgacg cttttccagc cgggaaggca tggaaactc 1440
 acgcccaggct gccacacagc aaaactggga ctatcggtg ctacgaatgc ccataaaagg 1500
 gtttaactct catccctacc cagcctggcg gaagctgaag ctcccagggg acagctcctt 1560
 gatgacctac cggggccacg gagtgtgca caccctcacc cgctgcgggt tctcccocat 1620
 tcatagcact gggccagcag tcatctacag tggctgtccc actggcaagg tgggtgtgta 1680
 cgacctctta agtggccaca ttgtgaagaa gctgaccaac cacaaggcct gtgtgctgta 1740
 cgtcagttgg caccctttg aagagaagat tgcagcagat tctgtgggag ggaacctgcy 1800
 tctgtggcag tacgcggagg ctgagtaact ccaggatgac atgccagaat ctgaggaaatg 1860
 tgccagcgcc ctgcggccag tgccccaatc ctctacaccc ttttctcctg cccagtagat 1920
 ccaacctcca gccccatata ggggtgaacct ctgtataagg tctctgcctc ctctccctt 1980
 tctcccttgt ggggaatgtt tggaggaatc actggcatt gatggggaat aacataagcc 2040
 tgggctctga gccctacgtg agccctggaa gattctcccc atggggcaga gtgtctctct 2100
 tacgtgctca caccacgta gcttgggtcc ctatctctgg ccagagtttg gcaggactgc 2160
 cattatctgg ggtgtggcct ctgcacgcaa gagaagtgtc ctgggtgttt ttaactatgt 2220
 ttgaatgta ggggttggat cctagagtag atgcctgagg ccacatctga acagacctgt 2280
 cagccagctc tgccagggtc tcaagttgag gattcaactg gccaatcaca ggaaggtgt 2340
 cctggccttt ctctcaggg tctctagggg aggggcatgg gtaagggtgt ttctccagca 2400
 cctcctctgg gtggggataa tgtctgctgt catgtctggg tctttaaggat aggaca 2456

<210> 54

<211> 1771

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2226116CB1

<400> 54

cggctcgagt taatatcttc gttgagcgga accttgctat tccataagag gatgtgtcca 60
 gtgtgtgtag agatttcatg ttttaaatcc tttgtacaga aatcctgctc ccaagtcca 120
 gataggctga cgggtcagag ggcaagacgt gacccagggc cgagaggggt agtgaccagg 180
 aaaaactgat tcaatcagat actgtttgtg ttcaagaaag tgacaaaaga cctgctgcat 240
 gaggccctcg tcttcagttt ctgtttcatg cccagcatta aaccaagtat ctcatcttgc 300
 caatttgact tctgtagggg ccatggcacc tgcaaggtgt ttctcagcaa gattgaggac 360

```

cgtgttttcag ggcgtggggc attgggcttt gtccacatgg gctggcctga agcccagccg 420
gctactgcga cagcgggctt ctcccaggct gctctcggtc ggccgtgccc accctgccaa 480
gcactcaggaa ctcccgggga agaagctgct ctctgagaaa aagctgaaaaa ggtactttgt 540
ggactatcgg agagtgtcgt tctgtggagg aaacggaggc gctgggggcaa gctgcttcca 600
cagtgcagccc cgcaaggagt ttggaggccc tgatggaggg gacggaggca acggtggaca 660
cgtcattctg agagttagcc agcaagtcaa gtccctgtcg tcggtctcgt cgcggtacca 720
gggtttcagt ggagaagatg gaggagagta aaactgcttc gggcgagtg gcgcgtcct 780
ctacatccgg gtcccctggg gcacgtggtg gaaggaggga ggcagagattg tggccgacct 840
gtcttcgctg ggagatgagt acattgccc gctggggggg gcaggaggga aaggcaaacg 900
cttcttctctg ggaacaaca accgtgcccc tgtgacctgt acccctggagc agccaggaca 960
gcagcgagtt ctccactcgg agctcaagac ggtggcccac gccggaatgg tgggattccc 1020
caacgcgggg aagtctcac tgctccgggc catttcaaac gccagaccgc ccgtggcttc 1080
ctaccggttc accaccctga agccccaact cgggatcgtc cactacgaag gccacctaca 1140
aatagcagtg ccgacatccc ccggcatcat acgaggcgcc caccagaaca ggggtctggg 1200
gtccgctctc ctcaggcaca tcgagcgctg ccgctttctc ttgttcgtgg tggatcttc 1260
tcacgctgag cctgtggactc aagttgacga tttaaaatat gaactggaga tgtatgaaaa 1320
gggctgtctc gcgaggcccc acgcaatcgt cgcaaacaa attgacctcc ctgaagacca 1380
agccaatctg tcccagctcc gggatcactt gggacaggag gtcatctgtc tgtcggcgtt 1440
gacggcgagc aacctggagc agctgctgtt ggcctgaag gtgctgtaag acgcctacgc 1500
ggaggccgag ctgggccaag gcccccagcc gctcagtggt tagccacgcc agagcggggt 1560
cgctctgggg cctctgtctg agcaaacctg ggtgtgaatt cgggtgtttt gaatgcataa 1620
agtgccttgt ggacacgggg gagtgtgtgt gcttctgggt ctctggggcc cgccctgctg 1680
cctgagatgc cctcatgttg ggaagcattc cgtgcccccc accccgcctg cctccgata 1740
ttctgcacc tgtcagcctg cgctgactga t 1771

```

<210> 55

<211> 2724

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2504472CB1

<400> 55

```

gctgaccagt tggcgacatg gtggcaccgc tgctggagac ttctcacgtg ttttctgtcc 60
caaacccggg gcggggagtc ctgaaactgga gctctgggcc cagaggactt ctggcctttg 120
gcacgtctcg ctccgttggt ctctatgacc cctgaaaaag ggttgttggt accaaactga 180
atggtcacac gcgccgagtc aattgcatac agtggatttg taaaaggagt ggctccctct 240
ctactgaatt agtttctgga ggatctgata atcaagtgat tcaactggaa atagaggata 300
atcagctgtt aaaaagcagt catcttcaag gccatgaagg acctgtttat gcggtgcatg 360
ctgtttacca gaggaggaca tcagatcctg cattatgtac actgatcgtt tctgcagctg 420
cagattctcg ttttcgactc tgggtctaaa agggctcoaga agtaattgtc cttaagactt 480
taaaactttg aaatgagatt gctttggctc tctgcttctc ttttttgcca aatactgatg 540
taccataatt agcatgtggc aatgatgatt gcagaattca catatttgct caacaaaaatg 600
atcagtttca gaaagtgtct tctctctgtg gacatgagga ttgatttaga ggagtggaa 660
gggcagcctt tggtagagat cttttctcag caagctgttc acagagattc ctgataagaa 720
tatggaagct gtatatcaag tcaacatctt tagaaaacta ggaatagcat aacataagac 780
tgaaagaaaa tacttttacc atagaaaatg aaagtgttaa aatagcattt gctgttactc 840
tggagacagt gtagccgggt catgaaaact gggtaaatgc agttcactg caacctgtgt 900
tttcaaaaag tgggtctcca cagcagccag tgagattatt atctgtctcc atggataaaa 960
ccatgtattc ctgggctcca gatgaagagt caggagtttg ctgagaacag gttcgagtag 1020
gtgaagttag tgggaatact ttgggatttt atgattgccca gttcaatgaa gatggctcca 1080
tgatcattgc tcatgtcttc accggagcgt tgcaccttgg gaaacagaat acagttaacc 1140

```

caagagagtg gactccagag attgtcatt caggacactt tgatgggtgc caagacctag 1200
 tctgggagtc agaaggagaa tttattatca ctgttggtac tgatcagaca actagacttt 1260
 ttgtctccatg gaagagaaaa gaccaatcac agtggtacttg gcatgaaatt gcaaggccct 1320
 agatacacgg gtagtacctg aaatgttttg caatgattaa tegggtttcag tttgtatctg 1380
 gagcagatga aaaagtctctt cgggtttttt ctgcacctcg gaattttgtg gaaaaatttt 1440
 gtgccattac aggacaatca ctgaatcatg tgctctgtaa tcaagatagt gatcttccag 1500
 aaggagccac tgtccctgca ttgggattat caaataaagc tgtctttcag ggagatatag 1560
 cttctcagcc ttctctgaga gaggagctgt taactagtac tgggttttgag tatcagcagg 1620
 tggcctttca gccctccata ctactgagc ctcccactga ggatcatctt ctgcagaata 1680
 ctttgtggcc tgaagttcaa aaactatag ggacggttta tgaatatatt tgtgttact 1740
 gtaacagttc aaagactctg cttgectcag cttgtaaggc agctaagaa gagcatgcag 1800
 ctatcattct ttggaacct acatcttgga aacaggtgca gaatttagtt tccacagtt 1860
 tgacagtcac gcagatggcc ttctcaccta atgagaagtt ctacttagct gtttccagag 1920
 atcgaaacctg tgcattgttg aaaaagcagg atacaatctc acctgagttc gagccagttt 1980
 tttagtcttt ttgcctcacc aacaaaatta ctctgtgca cagtagaatt atttggttct 2040
 gtgattggag tctgcagcag aagtatttct tcactgggag tcgagacaaa aagtggtgtg 2100
 tctgggtgta gtgcgactcc actgatgact gtattgagca caacattggc cctgtctct 2160
 cagtcctgga cgtgggtggg gctgtgacag ctgtcagcgt ctgcccagtg ctccaccttt 2220
 ctcaacgata cgtggttgca gtaggattgg agtgtggaaa gatttgccta tatacctgga 2280
 aaaaagctga tcaagttcca gaaataaat actggaccca ctgtgtagaa caagctcaa 2340
 gccaaagtca tacactggct atcagaaaaat tatgctggaa gaattgcagt ggaaaaactg 2400
 aacagaaagg agcagaaggt gctgagtggt tacacttgcc aagctgtggt gaagatcaca 2460
 ctgtgaagt acacagagtc aataaatgtg cactgtaatg gacttaataa gactatgct 2520
 gcagtcactg gtatcttaaa atattatcat gtaaacaggt catctttacc ttcataactg 2580
 aattgagttt ctgggttttt tttttttttg agatggagtc ttgttttgc ccaacctcca 2640
 cctccaggt tcaagcgatt ctctttcttc agcctctcga gttagctggga ctagaggcac 2700
 accaccatgc ccggctaatt ttgt 2724

<210> 56

<211> 2963

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3029920CB1

<400> 56

ggccgaagag gctggcagggt ggcgcctggt ggtgggtgct cctggtgaga ggagtcact 60
 ccgtgcgtgc gggcgagggt cggccccga gagccgcga catgaagaa gacgtgcga 120
 tctcgtgtgt gggggaacct agagttggga agacatcat gattatgct ctggtcagtg 180
 aagaatttcc agaagaggtt cctcccggg cagaagaaat caccattcca gctgatgtca 240
 cccgagagag agttccaaca cacattgtag attactcaga agcagaacag agtgaatgac 300
 aactctatca agaaatatct caggctaagt tcatctgtat agtgtatgct gtttaaca 360
 agcatctat tgataattat acaagtcgat ggattcctct cataaagtga agaacagaca 420
 aagacagcag gctgccttta atattgggtg ggaacaaatc tgatctgggt gaactataga 480
 gtatggagac catccttctt attatgaacc agtatacaga aatagaacac tgtgtggagt 540
 gttcagcgaa aaactgcgaag aacatatcag agctcttcta ttacgcacag aaagctgttc 600
 ttcatcttcc agggccctgt tactgcccag aggagaagg gatgaacaca gctgtatata 660
 aagcccttca tctgtatatt aaaaatatct atcaagataa tgatggtaact ctcaatgatg 720
 ctgaactcaa cttctttcag aggatttgtt tcaacactcc attagctcct caagctctgg 780
 aggatgtcaa gaattagtc agaaaacata taagtgtgtg ttgtggctga agtgggtg 840
 cctggaagg ttttctctt ttacacacac tttttatcca gagagggaga cagcaacta 900
 cttggactgt gcttcagaca ttgtgtatg atgatgacct ggatttgaca cctgaatatt 960

```

tggtccccct gctgaaaata cctcctgatt gcaactactga attaaatcat catgcatatt 1020
tatttctctca aagcacccttt gacaagcatg atttggatag agactgtgct ttgtcacctg 1080
atgagcttaa agatttattt aaagtcttcc ctacacatcc ttggggggcca gatgtggaata 1140
acacagtttg taccaatgaa agaggctgga taacctacca gggatctcct tccoagtgga 1200
cgctcacgac ttttttagat gtacagcggt gccctggaata ttgggctat ctaggctatt 1260
caatattgac tgagcaagag tctcaagctt cagctgttac agtgacaaga gataaaaaga 1320
tagacctgca gaaaaaacaa accaaaagaa atgtgttcag atgtaatgta attggagtgga 1380
aaaaactgttg gaaaagtggg gttctctcagg ctctctcttgg aagaaactta atgaggcgaga 1440
agaaaaattcg tgaagatcat aaatcctact atgcgattaa cactgtttat gtatatggac 1500
aagagaaaata cttgtgtgtg catgatattc cagaatcgga atttctcaact gaagctgaaa 1560
tcatttgtga tgttgtatgc ctgggtatat atgtcagcaa tcccaaatcc ttgtaact 1620
gtgccaggat ttttaagcaa cactttatgg acagcagaat acccttgctta atcgtagctg 1680
caaatgcaga cctgcgatga gttaaaacag aatacagtat ttcacctact gattttctgca 1740
ggaaaacaaa aatgcctcca ccacaagcct tcacttgcaa tactgctgat gcccccagta 1800
aggatattct ttgttaattg acaacaatgg ccactgtatcc gcacgtgaca caagctgacc 1860
tcaagagctc cagcttttgg cttcagagcaa gttttgtgtg tactgttttt cagtttttgg 1920
gctttgctat gtacaaagca ttattgaaac agcgatgata taaaagaaaa tactgtccct 1980
accaaaaaca aataacttta tgtacattta gaatgcttta agttctgctta gaattattga 2040
gatatttata catcgagagt tactttatta atattgttaa ttcacgcata agagtatttt 2100
aatgatagtt ataactgcag tattggctag catatggaaa gaaaacagct aacagccaaa 2160
ctaaaatggc taaattccag aggccaaaag ggaatatatt gtaaatatgt gtacatatcc 2220
aggcaagata tggctctcca agctgagttc tagaaatgat gttcttagac atttctaagt 2280
ggttattgta gtgtcactt ggctcactct tctaggttta agttagccca gagattgtat 2340
ttactcatgg atcactttat ttatttcaca ttactcaga atgatccttt gggttctata 2400
aggacataag gtacaaattt ccattgtctc tccattttta aaaaacataca acacagttgc 2460
agcttaccaa catgacattt ttccagtcag ttgtggtagg ccagccttga agccatcgca 2520
cagcttagaa acttgtgtag ctgagttggt agctcacctt taagggtgaa tttaggtaaa 2580
agcaaatagc agaggcggtta tctatgtgat tatgttgtct cctgtgcagt atgttgaatt 2640
ttatagccct tccaatgaaa taiaaaaaaa atttgtatat taccoatgtt tttagtttaa 2700
ataaagagct acccttaact ctgttgaatt tcatcccaag tgtaaatcat tctataatgg 2760
ctgtgtctgt tatagtatat tacagtaact gcactgtgtc caaagtgttc tatatcaggg 2820
taggataacc tagagcgagt aattttttta atgataaaat aaatctaatt atataaact 2880
ctcatgataa accatatttt tccatcatca gccctttcaa gtatttaaat aaataactgc 2940
tgtgtactgt gaaaaaaaaa aaa 2963

```

<210> 57

<211> 3332

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3332415CB1

<400> 57

```

gcttggcaga ggctggcggg catcgtgccg gtccctgccg gtctccgggg caecgggcca 60
ccgcccacac cctctctccc tgccatggag ccgagctggt acgctcagaa gcagctctga 120
ccgcggaggc gaagccgcgg ggccctctggg ctacgacagg agggagcgac ggggctctcg 180
gcgcagacct ccgggtcgga gctggacggg agatgttccc ttccggagagg cagctccttc 240
acattcttaa caactggccc caactgggac ttcaacttga aaagaaaaacg cagagagaaa 300
gatgatgatg ttgtaagcct tagcagcctt gatctgaagg agccaagcaa taaaagagtt 360
cgactctgtg ctcgtgtcac gtccttggca aattttaact ctctgtgaag aaatggagct 420
gtcagacgtt ttggtcaaac aatacagtc tttacccttc gtggtgacca cagatcccca 480
gcctctgccc agaagtttct tagcaggtca acagtcacca caccgcgcaa gagaaggagc 540

```

agtgcatctgt ggtcagagat gctggacatc accatgaagg agtctctcac caccaggagg 600
 atcagacggc agggaggcaat atatgaaatg tcccgagggt aacaggagattt aattgaggat 660
 ctcaaaacttg caagaaaggc ctaccatgac cccatgttaa agttgtccat catgtcagaa 720
 gaggaactca cacatatatt tgggtgatctg gactctttaca tacctctgca tgaagatttg 780
 ttgacaagaa taggagaagc aaccaagcct gatggaaacag tggagcagat tggtcacatt 840
 ctctgtgagct ggtttaccgcg cttgaatgcc tacagaggtt actgtagtaa ccagctggca 900
 gccaaagctt ttcttgaatca aaagaacacg gatccaagag tccaagactt cctccagcga 960
 tgtctcagat ctcccttcag tcgaaaaacta gatctttgga gtttcttaga tatccctcga 1020
 agtcgcctag tcaaatacc cttactgtta aaagaatttc ttaaacacac tccaaaagag 1080
 caccctgatg ttacgtctct ggaggatgct atattgtata tacaggagatt cctctctgat 1140
 atcaacttga agaaaggtga atccgagtg cagtattaca tcgacaagct ggagtagctg 1200
 gatgaaaagc agagggaccc cagaatcgaa gcgagcaaaag tgctgtctgt ccagtgaggag 1260
 ctgctggagca agagtggaca taaactttac attttctctg ttcaagacat cttggttctg 1320
 actcggcccg tcacacggaa cgaaacggcac tcttaccagg tttaccggca gccaatccca 1380
 gtccaagagc tagctcctaga agacctcgag gatggagatg tgagaatggg aggtctcttt 1440
 cgaggagctt tcaagaaact agagaaagct aaaaatatct ttagaattcg cttccatgac 1500
 cctctctccg cccagctctca cactctgcaa gccaatgacg tgttccaca gcagcagtg 1560
 ttcaactgtg ttcgagcgcc cattgccccc ttccagtcgg caggcagctcc acctgagctg 1620
 caggggcctgc cggagctgca cgaagagtgt gaggggaacc acccctctgc gagggaactc 1680
 acagcccaga ggagggcatc cacagtcttc agtgttactc aggtagaagt cgtatgaaa 1740
 gcttacagat tgggctctgg catgcagatg gcagaggaca gcaagagctt aaagacacac 1800
 cagacacagc cggcctccg aagagcgagg gacaaaagcc ttctctgttg caaacggaaa 1860
 gagacttttg tctagagaaag gctctgtgtg ttaactgagt ggagagactg tttgtttata 1920
 aatgtgtaca gttttgtttt ctctgtaagg gagcatcata gggttacttt ataccagttg 1980
 taacattttt attgtttttg gttgttcttt ttcttttttt taatggcagc taagatata 2040
 cagattactg ttaaatgtca gtcccttttt tttaaaagt attttcttga gttatttaga 2100
 acatggttag cctggtattt tttaatacaa caaaatattt atgaaatggg ttttctctta 2160
 attctggatt catcatggct ttctaatacc aattgtataa tttaacatat tcccaaaaac 2220
 ttgaattttt gcaaatgtct gaattctgcc agtgtttctt tgctaacgct tgcattgcaa 2280
 atttgaaatt ttaacattgg caccctaaac ctacatggaa tgtatgtctg gagtatttca 2340
 aactttacat tgaacataaa ttctcttgga aaacaaacca taagcctgag gaggttttca 2400
 tcaactggct tgccttatat tagttttgtt ttactgtac attctctcat ttacattcat 2460
 ttaacctgcc gattatttaa tttttttatt gtaagtagt ttttagcatt tgcctttatt 2520
 tttttacttt gatgcctttt caaattggca tgccttttaa gtatttttct tctctgattaa 2580
 aaatgtgtgt gtgtgtgtgt gtgtgtgtgt atatatatat atttttttaa atcacattaa 2640
 ttttaccagc tgaaccaaag ccatactggt tttaggccaa ttaagaaaaat tgcatttttt 2700
 aaagtgtagc atttcagggt aaagacccat gaaatggctt gatgtattct agactactga 2760
 aagaaaaaca cttcaaaagt tttgttgaaa gttttagtgt tgcctgaaat cgaagaggga 2820
 aggtgtattg tagtgattta aaagaaaaag agaggaaaaag agagtatttt tgccttcaag 2880
 taaattgtct ggttctgcca gacatttcc aagtgtgaaa ggagatagga gaagctcaac 2940
 ttagggcgct gtagtgaatt gtagaaggct cgaggggacg tggacttatt tgccttggtt 3000
 tgcaataacc gcaaatcaatg agtttgaaaa gaaacaatga aatgtgttaa aaatttgacc 3060
 atattagata aatttttggtg gatttagtca taagatggaa aaagactcgt gaattcttta 3120
 ttacaaaaat ttctctgtaa aatgggatca tcatctttga aaggggggag gagggatgaa 3180
 agcccgatta taatgggtgat caattccaag cagtgttgac tattctgtga aatattttt 3240
 gccagtggaa atgataatca gaaaagactg taaatagatc catccaaagt atttctctgt 3300
 acaaatgaat gatactatta aaaaaaaaaa aa 3332

<210> 58
 <211> 2617
 <212> DNA
 <213> Homo sapiens
 <220>

<223> Incyte ID No: 4031536CB1

<400> 58
 tttgtaaatg tgccgtgtatt acatgtagag agtattctgc aaccaagagg agttttaaaa 60
 tgtcaaaaacc gggaaaaacct actctaaacc atggccttgg tccgtgtgat cttaaaagtg 120
 caaaagagcc tctaccacat caaactgtga tgaggatatt tagcattagc atcattgccc 180
 aaggccctccc tttttgtcga agacggatga aaagaaagt ttgagggtcc 240
 gctctttttc tttgggaaa aaacatgtca aagtctcaga atatacaagt accactgggc 300
 ttgtaccatg ttcagcaaca ccaacaactt ttggggacct cagagcagcc aatggccaag 360
 ggcaacaacg agccgaatt acatctgttc agccacctac aggcctccag catatggcta 420
 aaatgtttca gagctggagt ggaccagaga aattgtctgc tttagatgaa ctcattgata 480
 gtgtgtgaac aacacaagta aaacatatga tgcaagtgat agaaccctac ttccaacgag 540
 acttcatttc attgctccct aaagagttag cactctatgt gctttcattc ctggaaacca 600
 aagacctgct acaagcagct cagacatgtc gctactggag aattttggct gaagacaacc 660
 tctctggag agagaaatgc aaagaagagg ggattgatga accattgac atcaagagaa 720
 gaaaagtaat aaaaccaggt ttcatacaca gtccatggaa aagtgcatac atcagacagc 780
 acagaattga tactaactgg aggcgaggag aactcaaatc tctaaagtg ctgaaaggag 840
 atgatgatca ttgatgtaca tgcttacagt tttgtggtaa ccgaatagt agtgcgtctg 900
 atgacaacac tttaaaagtt tgggtcagcag tcacaggcaa atgtctgaga accattagtgg 960
 gacatacagg tggagattgg tcatcacaaa tgagagacaa catcatcatt agtggatcta 1020
 cagatcggac actcaaatg ttggaatgcag agactggaga atgtatacac accttatatg 1080
 ggcatacttc cactgtcggt tgtatgcatt ttcataaaaa aagagtgttt agcggttctc 1140
 gagatgccac tcttaggttt tgggaattg agacaggcca gtgtttacat gttttgatgg 1200
 gtcatgttgc agcagtcgcg tgtgttcaat atgatggcag gagggtgttt agtggagcat 1260
 atgattttat ggtaaaagtg tgggatccag agactgaac ctgtctcac acgttgcagg 1320
 ggcatactaa tagatctat tcattacagt ttgatggtat ccatgtggtg agtggatctc 1380
 ttgatacatc aatccgtgtt tgggatgtgg agacagggaa ttgcattcac acgttaacag 1440
 ggcaccagtc gttaaacaagt ggaatggaac tcaaaagacaa tattcttgtc tctgggaatg 1500
 cagatctctac agtttaaaatc tgggatatca aaacaggaca gtgtttacaa acattgcaag 1560
 gtcccaacaa gcatcagagt gctgtgacct gtttacagtt caacaagaac ttgttaatta 1620
 ccagctcaga tgatggaact gtaaaactat gggacttgaa aacgggtgaa ttatttcgaa 1680
 acctagtca attggagagt ggggggagtg gggggagtgt gtggcgagtc agagcctcaa 1740
 acacaaagct ggtgtgtgca gttgggagtc ggaatgggac tgaagaaacc aagctgctgg 1800
 tgctggagct tgatgtggac atgaagtga gacagaaaa gatgaattg tocaattgtg 1860
 tagacgatat actccctgcc ctccccctg caaaaagaaa aaaagaaaaa aaaaagaaa 1920
 aaatcccttg ttctcagtg tgccaggtgt tggcttgggg caacagattg aaaagacct 1980
 cagactaaga aggaaaagaa gaagagatga caaacctaa ctgacaagag aggcgtctgc 2040
 tgtctcatca cataaaaggc ttacattttt actgagggca gctttgcmaa atgagacttt 2100
 ctaaatcaaa ccaggtgcaa ttatttcttt attttctct ccagtggttc ttgggagctc 2160
 ttaatgctga aacatcatta cagattctgc tagcctgttc ttttaccact gacagctaga 2220
 cacctagaaa ggaactgcaa taatatcaaa acaagtactg ttgactttc taattagaga 2280
 gcatctgcga caaaaagtca tttttctgga tgggaaaaagc ttaaaaaaat tactgtgaat 2340
 tggttttgta cagtatatcat gaaaagcttt ttttttttt ttggccaacc attgccaatt 2400
 tcaatcaatc acagtattag cctctgttaa tctatttact gttggtcca tatacattc 2460
 tcaatgcata tgtgtgtcaa aggtggcaag ttgtcctgg tctgtgagt cctgagatgg 2520
 atttaattct tgatgctgg gctagaagta ggtcttcaaa tatgggattg ttgtcccaac 2580
 cctgtactgt actccagtg gccaaaacta tttatgc 2617